

# **INTRO to DATA SCIENCE**

## **LECTURE 5: NAIVE BAYESIAN CLASSIFICATION**

## **LAST TIME:**

- CLASSIFICATION PROBLEMS**
- TRAINING/TEST SETS & CROSS-VALIDATION**
- KNN CLASSIFICATION**

**QUESTIONS?**

# HOW'S THE HOMEWORK GOING?







**I. INTRO TO PROBABILITY**

**II. NAÏVE BAYESIAN CLASSIFICATION**

**EXERCISES:**

**III. NAÏVE BAYES CLASSIFICATION IN PYTHON**

# **I. INTRO TO PROBABILITY**



*Q: What is a probability?*

*Q: What is a **probability**?*

*A: A number between ? and ? that characterizes the likelihood that some event will occur.*

*Q: What is a **probability**?*

*A: A number between **0** and **1** that characterizes the likelihood that some event will occur.*

*Q: What is a **probability**?*

*A: A number between 0 and 1 that characterizes the likelihood that some event will occur.*

*The probability of event  $A$  is denoted  $P(A)$ .*

*Q: What is the set of all possible events called?*

*Q: What is the set of all possible events called?*

*A: This set is called the **sample space**  $\Omega$ . Event  $A$  is a member of the sample space, as is every other event.*

*Q: What is the set of all possible events called?*

*A: This set is called the **sample space**  $\Omega$ . Event  $A$  is a member of the sample space, as is every other event.*

*The probability of the sample space  $P(\Omega)$  is 1.*



*Q: Consider two events  $A$  &  $B$ . How can we characterize the intersection of these events?*



*Q: Consider two events  $A$  &  $B$ . How can we characterize the intersection of these events?*

*A: With the joint probability of  $A$  and  $B$ , written  $P(AB)$ .*

*Q: Suppose event  $B$  has occurred. What quantity represents the probability of  $A$  **given** this information about  $B$ ?*

*Q: Suppose event  $B$  has occurred. What quantity represents the probability of  $A$  **given** this information about  $B$ ?*

*A: The intersection of  $A$  &  $B$  divided by region  $B$ .*

*Q: Suppose event  $B$  has occurred. What quantity represents the probability of  $A$  **given** this information about  $B$ ?*

*A: The intersection of  $A$  &  $B$  divided by region  $B$ .*

**NOTE**

*This information about  $B$  transforms the sample space.*

*Take a moment to convince yourself of this!*

*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B?*

*A: The intersection of A & B divided by region B.*

*This is called the **conditional probability** of A given B, written  $P(A|B) = P(AB) / P(B)$ .*

**NOTE**

*This information about B transforms the sample space.*

*Take a moment to convince yourself of this!*



*Q: Suppose event B has occurred. What quantity represents the probability of A **given** this information about B?*

*A: The intersection of A & B divided by region B.*

*This is called the **conditional probability** of A given B, written  $P(A|B) = P(AB) / P(B)$ .*

**NOTE**

*This information about B transforms the sample space.*

*Take a moment to convince yourself of this!*

*Notice, with this we can also write  $P(AB) = P(A|B) * P(B)$ .*

*Q: What does it mean for two events to be independent?*

*Q: What does it mean for two events to be **independent**?*

*A: Information about one does not affect the probability of the other.*



*Q: What does it mean for two events to be **independent**?*

*A: Information about one does not affect the probability of the other.*

*This can be written as  $P(A|B) = P(A)$ .*

*Q: What does it mean for two events to be independent?*

*A: Information about one does not affect the probability of the other.*

*This can be written as  $P(A|B) = P(A)$ .*

*Using the definition of the conditional probability, we can also write:*

$$P(A|B) = P(AB) / P(B) = P(A) \rightarrow P(AB) = P(A) * P(B)$$

*Probably the only calculation in the whole course:*

*Probably the only calculation in the whole course:*

$$P(AB) = P(A|B) * P(B)$$

*from last slide*

*Probably the only calculation in the whole course:*

$$P(AB) = P(A|B) * P(B)$$

$$P(BA) = P(B|A) * P(A)$$

*from last slide  
by substitution*

*Probably the only calculation in the whole course:*

$$P(AB) = P(A|B) * P(B)$$

*from last slide*

$$P(BA) = P(B|A) * P(A)$$

*by substitution*

*But*  $P(AB) = P(BA)$

*since event AB = event BA*

*Probably the only calculation in the whole course:*

$$P(AB) = P(A|B) * P(B)$$

*from last slide*

$$P(BA) = P(B|A) * P(A)$$

*by substitution*

*But*  $P(AB) = P(BA)$

*since event AB = event BA*

$$\rightarrow P(A|B) * P(B) = P(B|A) * P(A)$$

*by combining the above*

*Probably the only calculation in the whole course:*

$$P(AB) = P(A|B) * P(B)$$

*from last slide*

$$P(BA) = P(B|A) * P(A)$$

*by substitution*

*But*  $P(AB) = P(BA)$

*since event AB = event BA*

$$\rightarrow P(A|B) * P(B) = P(B|A) * P(A)$$

*by combining the above*

$$\rightarrow P(A|B) = P(B|A) * P(A) / P(B)$$

*by rearranging last step*



*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

*Some facts:*

*- This is a simple algebraic relationship using elementary definitions.*

*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

*Some facts:*

- This is a simple algebraic relationship using elementary definitions.*
- It's interesting because it's kind of a "wormhole" between two different "interpretations" of probability.*

*This result is called **Bayes' theorem**. Here it is again:*

$$P(A|B) = P(B|A) * P(A) / P(B)$$

*Some facts:*

- This is a simple algebraic relationship using elementary definitions.*
- It's interesting because it's kind of a "wormhole" between two different "interpretations" of probability.*
- It's a very powerful computational tool.*

*Briefly, the two interpretations can be described as follows:*

*Briefly, the two interpretations can be described as follows:*

*The frequentist interpretation regards an event's probability as its limiting frequency across a very large number of trials.*

*Briefly, the two interpretations can be described as follows:*

*The frequentist interpretation regards an event's probability as its limiting frequency across a very large number of trials.*

*The Bayesian interpretation regards an event's probability as a "degree of belief," which can apply even to events that have not yet occurred.*

*If this sounds crazy to you, don't worry...we won't dwell on the theoretical details.*



*If this sounds crazy to you, don't worry...we won't dwell on the theoretical details.*

*If this sounds interesting, there are plenty of resources available to learn more about Bayesian inference.*

*If this sounds crazy to you, don't worry...we won't dwell on the theoretical details.*

*If this sounds interesting, there are plenty of resources available to learn more about Bayesian inference.*

*This a good direction to head if you like math and/or if you're interested in learning about cutting-edge data science techniques.*

# **II. NAÏVE BAYESIAN CLASSIFICATION**

*Suppose we have a dataset with features  $x_1, \dots, x_n$  and a class label  $C$ .  
What can we say about classification using Bayes' theorem?*

*Suppose we have a dataset with features  $x_1, \dots, x_n$  and a class label  $C$ . What can we say about classification using Bayes' theorem?*


$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*Bayes' theorem can help us to determine the probability of a record belonging to a class, given the data we observe.*

*Each term in this relationship has a name, and each plays a distinct role in any Bayesian calculation (including ours).*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*This term is the **likelihood function**. It represents the joint probability of observing features  $\{x_i\}$  given that that record belongs to class C.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$



*This term is the **likelihood function**. It represents the joint probability of observing features  $\{x_i\}$  given that that record belongs to class C.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*We can observe the value of the likelihood function from the training data.*



*This term is the **prior probability** of  $c$ . It represents the probability of a record belonging to class  $c$  before the data is taken into account.*

$$P(\text{class } C | \{x_i\}) = \frac{P(\{x_i\} | \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$



*This term is the **prior probability** of  $c$ . It represents the probability of a record belonging to class  $c$  before the data is taken into account.*

$$P(\text{class } C | \{x_i\}) = \frac{P(\{x_i\} | \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*The value of the prior is also observed from the data.*

*This term is the **normalization constant**. It doesn't depend on  $C$ , and is generally ignored until the end of the computation.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$




*This term is the **normalization constant**. It doesn't depend on  $C$ , and is generally ignored until the end of the computation.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*The normalization constant doesn't tell us much.*

*This term is the **posterior probability** of  $c$ . It represents the probability of a record belonging to class  $c$  after the data is taken into account.*

$$P(\text{class } C | \{x_i\}) = \frac{P(\{x_i\} | \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$



*This term is the **posterior probability** of  $c$ . It represents the probability of a record belonging to class  $c$  after the data is taken into account.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*The goal of any Bayesian computation is to find (“learn”) the posterior distribution of a particular variable.*

*The idea of Bayesian inference, then, is to **update** our beliefs about the distribution of  $C$  using the data (“evidence”) at our disposal.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*Then we can use the posterior for prediction.*

*Q: What piece of the puzzle we've seen so far looks like it could intractably difficult in practice?*



*Remember the likelihood function?*

$$P(\{\mathbf{x}_i\} | C) = P(\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\} | C)$$

*Remember the likelihood function?*

$$P(\{\mathbf{x}_i\} \mid C) = P(\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\} \mid C)$$

*Observing this exactly would require us to have enough data for every possible combination of features to make a reasonable estimate.*

*Q: What piece of the puzzle we've seen so far looks like it could intractably difficult in practice?*

*A: Estimating the full likelihood function.*

*Q: So what can we do about it?*

*Q: So what can we do about it?*

*A: Make a simplifying assumption. In particular, we assume that the features  $x_i$  are conditionally independent from each other:*

*Q: So what can we do about it?*

*A: Make a simplifying assumption. In particular, we assume that the features  $x_i$  are conditionally independent from each other:*

$$P(\{x_i\} | C) = P(x_1, x_2, \dots, x_n | C) \approx P(x_1 | C) * P(x_2 | C) * \dots * P(x_n | C)$$

*Q: So what can we do about it?*

*A: Make a simplifying assumption. In particular, we assume that the features  $\mathbf{x}_i$  are conditionally independent from each other:*

$$P(\{\mathbf{x}_i\} | C) = P(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n | C) \approx P(\mathbf{x}_1 | C) * P(\mathbf{x}_2 | C) * \dots * P(\mathbf{x}_n | C)$$

*This “naïve” assumption simplifies the likelihood function to make it tractable.*

$$P(\text{class } C \mid \{x_i\}) = \frac{P(\{x_i\} \mid \text{class } C) \cdot P(\text{class } C)}{P(\{x_i\})}$$

*the **training phase** of the model involves computing the likelihood function, which is the conditional probability of each feature given each class.*

*the **prediction phase** of the model involves computing the posterior probability of each class given the observed features, and choosing the class with the highest probability.*



## ***Advantages:***

- *Fast to train (single scan). Fast to classify*
- *Not sensitive to irrelevant features*
- *Handles real and discrete data*
- *Handles streaming data well*

## ***Disadvantages:***

- *Assumes independence of features*

# **LAB**

## **III. NAIVE BAYESIAN CLASSIFICATION**