

## Appendix B: SMBKC Stock Assessment Input Files

## The data file:

```
## #=====
## # Gmacs Main Data File Version 1.1: SM17 example
## # GEAR_INDEX DESCRIPTION
## # 1 : Pot fishery retained catch.
## # 1 : Pot fishery with discarded catch.
## # 2 : Trawl bycatch
## # 3 : Fixed bycatch
## # 4 : Trawl survey
## # 5 : Pot survey
##
## # Fisheries: 1 Pot Fishery, 2 Pot Discard, 3 Trawl by-catch, 3 Fixed by-catch
## # Surveys: 4 NMFS Trawl Survey, 5 Pot Survey
## #=====
##
## 1978 # Start year
## 2017 # End year
## 2018 # Projection year
## 5 # Number of seasons
## 5 # Number of distinct data groups (among fishing fleets and surveys)
## 1 # Number of sexes
## 1 # Number of shell condition types
## 1 # Number of maturity types
## 3 # Number of size-classes in the model
## 5 # Season recruitment occurs
## 5 # Season molting and growth occurs
## 4 # Season to calculate SSB
## 1 # Season for N output
## # size_breaks (a vector giving the break points between size intervals with dimension nclass+1)
## 90 105 120 135
## # weight-at-length input method (1 = allometry i.e.  $w_l = a \cdot l^b$ , 2 = vector by sex, 3 = matrix by sex)
## 3
## # weight-at-length allometry  $w_l = a \cdot l^b$ 
## 4.03E-07
## # b (male, female)
## 3.141334
## # Male weight-at-length
## 0.000748427 0.001165731 0.001930510
## 0.000748427 0.001165731 0.001688886
## 0.000748427 0.001165731 0.001922246
## 0.000748427 0.001165731 0.001877957
## 0.000748427 0.001165731 0.001938634
## 0.000748427 0.001165731 0.002076413
## 0.000748427 0.001165731 0.001899330
## 0.000748427 0.001165731 0.002116687
## 0.000748427 0.001165731 0.001938784
## 0.000748427 0.001165731 0.001939764
## 0.000748427 0.001165731 0.001871067
## 0.000748427 0.001165731 0.001998295
## 0.000748427 0.001165731 0.001870418
## 0.000748427 0.001165731 0.001969415
## 0.000748427 0.001165731 0.001926859
## 0.000748427 0.001165731 0.002021492
```

[illegible]

```

##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.1800    0.0000    0.4500    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
##      0.0000    0.4400    0.0000    0.1900    0.3700
## # Fishing fleet names (delimited with : no spaces in names)
## Pot_Fishery:Trawl_Bycatch:Fixed_bycatch
## # Survey names (delimited with : no spaces in names)
## NMFS_Trawl:ADFG_Pot
## # Number of catch data frames
## 4
## # Number of rows in each data frame
## 28 16 26 26
## ## CATCH DATA
## ## Type of catch: 1 = retained, 2 = discard
## ## Units of catch: 1 = biomass, 2 = numbers
## ## for SMBKC Units are in number of crab for landed & 1000 kg for discards.
## ## Male Retained
## # year seas fleet sex obs cv type units mult effort discard_mortality
## 1978 2 1 1 436126 0.03 1 2 1 0 0
## 1979 2 1 1 52966 0.03 1 2 1 0 0
## 1980 2 1 1 33162 0.03 1 2 1 0 0
## 1981 2 1 1 1045619 0.03 1 2 1 0 0
## 1982 2 1 1 1935886 0.03 1 2 1 0 0
## 1983 2 1 1 1931990 0.03 1 2 1 0 0
## 1984 2 1 1 841017 0.03 1 2 1 0 0
## 1985 2 1 1 436021 0.03 1 2 1 0 0
## 1986 2 1 1 219548 0.03 1 2 1 0 0
## 1987 2 1 1 227447 0.03 1 2 1 0 0
## 1988 2 1 1 280401 0.03 1 2 1 0 0
## 1989 2 1 1 247641 0.03 1 2 1 0 0
## 1990 2 1 1 391405 0.03 1 2 1 0 0
## 1991 2 1 1 726519 0.03 1 2 1 0 0
## 1992 2 1 1 545222 0.03 1 2 1 0 0
## 1993 2 1 1 630353 0.03 1 2 1 0 0
## 1994 2 1 1 827015 0.03 1 2 1 0 0
## 1995 2 1 1 666905 0.03 1 2 1 0 0
## 1996 2 1 1 660665 0.03 1 2 1 0 0
## 1997 2 1 1 939822 0.03 1 2 1 0 0
## 1998 2 1 1 635370 0.03 1 2 1 0 0
## 2009 2 1 1 103376 0.03 1 2 1 0 0

```

## 2010	2	1	1	298669	0.03	1	2	1	0	0
## 2011	2	1	1	437862	0.03	1	2	1	0	0
## 2012	2	1	1	379386	0.03	1	2	1	0	0
## 2014	2	1	1	69109	0.03	1	2	1	0	0
## 2015	2	1	1	24407	0.03	1	2	1	0	0
## 2016	2	1	1	24.407	0.03	1	2	1	0	0
## # Male discards Pot fishery										
## 1990	2	1	1	254.9787861	0.6	2	1	1	0	0.2
## 1991	2	1	1	531.4483252	0.6	2	1	1	0	0.2
## 1992	2	1	1	1050.387026	0.6	2	1	1	0	0.2
## 1993	2	1	1	951.4626128	0.6	2	1	1	0	0.2
## 1994	2	1	1	1210.764588	0.6	2	1	1	0	0.2
## 1995	2	1	1	363.112032	0.6	2	1	1	0	0.2
## 1996	2	1	1	528.5244687	0.6	2	1	1	0	0.2
## 1997	2	1	1	1382.825328	0.6	2	1	1	0	0.2
## 1998	2	1	1	781.1032977	0.6	2	1	1	0	0.2
## 2009	2	1	1	123.3712279	0.2	2	1	1	0	0.2
## 2010	2	1	1	304.6562225	0.2	2	1	1	0	0.2
## 2011	2	1	1	481.3572126	0.2	2	1	1	0	0.2
## 2012	2	1	1	437.3360731	0.2	2	1	1	0	0.2
## 2014	2	1	1	45.4839749	0.2	2	1	1	0	0.2
## 2015	2	1	1	21.19378597	0.2	2	1	1	0	0.2
## 2016	2	1	1	0.021193786	0.2	2	1	1	0	0.2
## # Trawl fishery discards										
## 1991	2	2	1	3.538	0.31	2	1	1	0	0.8
## 1992	2	2	1	1.996	0.31	2	1	1	0	0.8
## 1993	2	2	1	1.542	0.31	2	1	1	0	0.8
## 1994	2	2	1	0.318	0.31	2	1	1	0	0.8
## 1995	2	2	1	0.635	0.31	2	1	1	0	0.8
## 1996	2	2	1	0.500	0.31	2	1	1	0	0.8
## 1997	2	2	1	0.500	0.31	2	1	1	0	0.8
## 1998	2	2	1	0.500	0.31	2	1	1	0	0.8
## 1999	2	2	1	0.500	0.31	2	1	1	0	0.8
## 2000	2	2	1	0.500	0.31	2	1	1	0	0.8
## 2001	2	2	1	0.500	0.31	2	1	1	0	0.8
## 2002	2	2	1	0.726	0.31	2	1	1	0	0.8
## 2003	2	2	1	0.998	0.31	2	1	1	0	0.8
## 2004	2	2	1	0.091	0.31	2	1	1	0	0.8
## 2005	2	2	1	0.500	0.31	2	1	1	0	0.8
## 2006	2	2	1	2.812	0.31	2	1	1	0	0.8
## 2007	2	2	1	0.045	0.31	2	1	1	0	0.8
## 2008	2	2	1	0.272	0.31	2	1	1	0	0.8
## 2009	2	2	1	0.635	0.31	2	1	1	0	0.8
## 2010	2	2	1	0.363	0.31	2	1	1	0	0.8
## 2011	2	2	1	0.181	0.31	2	1	1	0	0.8
## 2012	2	2	1	0.100	0.31	2	1	1	0	0.8
## 2013	2	2	1	0.400	0.31	2	1	1	0	0.8
## 2014	2	2	1	0.100	0.31	2	1	1	0	0.8
## 2015	2	2	1	0.100	0.31	2	1	1	0	0.8
## 2016	2	2	1	0.100	0.31	2	1	1	0	0.8
## # Fixed fishery discards										
## 1991	2	3	1	0.045	0.31	2	1	1	0	0.5
## 1992	2	3	1	2.268	0.31	2	1	1	0	0.5
## 1993	2	3	1	0.500	0.31	2	1	1	0	0.5

## 1994	2	3	1	0.091	0.31	2	1	1	0	0.5
## 1995	2	3	1	0.136	0.31	2	1	1	0	0.5
## 1996	2	3	1	0.045	0.31	2	1	1	0	0.5
## 1997	2	3	1	0.181	0.31	2	1	1	0	0.5
## 1998	2	3	1	0.907	0.31	2	1	1	0	0.5
## 1999	2	3	1	1.361	0.31	2	1	1	0	0.5
## 2000	2	3	1	0.500	0.31	2	1	1	0	0.5
## 2001	2	3	1	0.862	0.31	2	1	1	0	0.5
## 2002	2	3	1	0.408	0.31	2	1	1	0	0.5
## 2003	2	3	1	1.134	0.31	2	1	1	0	0.5
## 2004	2	3	1	0.635	0.31	2	1	1	0	0.5
## 2005	2	3	1	0.590	0.31	2	1	1	0	0.5
## 2006	2	3	1	1.451	0.31	2	1	1	0	0.5
## 2007	2	3	1	69.717	0.31	2	1	1	0	0.5
## 2008	2	3	1	6.622	0.31	2	1	1	0	0.5
## 2009	2	3	1	7.530	0.31	2	1	1	0	0.5
## 2010	2	3	1	9.571	0.31	2	1	1	0	0.5
## 2011	2	3	1	1.800	0.31	2	1	1	0	0.5
## 2012	2	3	1	1.600	0.31	2	1	1	0	0.5
## 2013	2	3	1	0.8	0.31	2	1	1	0	0.5
## 2014	2	3	1	1.1	0.31	2	1	1	0	0.5
## 2015	2	3	1	1.600	0.31	2	1	1	0	0.5
## 2016	2	3	1	3.600	0.31	2	1	1	0	0.5

## ## RELATIVE ABUNDANCE DATA

## ## Units of abundance: 1 = biomass, 2 = numbers

## ## for SMBKC Units are in crabs for Abundance.

## ## Number of relative abundance indices

## 2

## ## Number of rows in each index

## 40 9

## # Survey data (abundance indices, units are mt for trawl survey and crab/potlift for pot survey)

## # Year, Seas, Fleet, Sex, Abundance, CV units

## 1978	1	4	1	6832.819	0.394	1
## 1979	1	4	1	7989.881	0.463	1
## 1980	1	4	1	9986.830	0.507	1
## 1981	1	4	1	6551.132	0.402	1
## 1982	1	4	1	16221.933	0.344	1
## 1983	1	4	1	9634.250	0.298	1
## 1984	1	4	1	4071.218	0.179	1
## 1985	1	4	1	3110.541	0.210	1
## 1986	1	4	1	1416.849	0.388	1
## 1987	1	4	1	2278.917	0.291	1
## 1988	1	4	1	3158.169	0.252	1
## 1989	1	4	1	6338.622	0.271	1
## 1990	1	4	1	6730.130	0.274	1
## 1991	1	4	1	6948.184	0.248	1
## 1992	1	4	1	7093.272	0.201	1
## 1993	1	4	1	9548.459	0.169	1
## 1994	1	4	1	6539.133	0.176	1
## 1995	1	4	1	5703.591	0.178	1
## 1996	1	4	1	9410.403	0.241	1
## 1997	1	4	1	10924.107	0.337	1
## 1998	1	4	1	7976.839	0.355	1
## 1999	1	4	1	1594.546	0.182	1

```

## 2000 1 4 1 2096.795 0.310 1
## 2001 1 4 1 2831.440 0.245 1
## 2002 1 4 1 1732.599 0.320 1
## 2003 1 4 1 1566.675 0.336 1
## 2004 1 4 1 1523.869 0.305 1
## 2005 1 4 1 1642.017 0.371 1
## 2006 1 4 1 3893.875 0.334 1
## 2007 1 4 1 6470.773 0.385 1
## 2008 1 4 1 4654.473 0.284 1
## 2009 1 4 1 6301.470 0.256 1
## 2010 1 4 1 11130.898 0.466 1
## 2011 1 4 1 10931.232 0.558 1
## 2012 1 4 1 6200.219 0.339 1
## 2013 1 4 1 2287.557 0.217 1
## 2014 1 4 1 6029.220 0.449 1
## 2015 1 4 1 5877.433 0.770 1
## 2016 1 4 1 3485.909 0.393 1
## 2017 1 4 1 1793.760 0.599 1
## 1995 1 5 1 12042.000 0.130 2
## 1998 1 5 1 12531.000 0.060 2
## 2001 1 5 1 8477.000 0.080 2
## 2004 1 5 1 1667.000 0.150 2
## 2007 1 5 1 8643.000 0.090 2
## 2010 1 5 1 10209.000 0.130 2
## 2013 1 5 1 5643.000 0.190 2
## 2015 1 5 1 2805.000 0.180 2
## 2016 1 5 1 2378.000 0.186 2
## ## Number of length frequency matrices
## 3
## ## Number of rows in each matrix
## 15 40 9
## ## Number of bins in each matrix (columns of size data)
## 3 3 3
## ## SIZE COMPOSITION DATA FOR ALL FLEETS
## ## SIZE COMP LEGEND
## ## Sex: 1 = male, 2 = female, 0 = both sexes combined
## ## Type of composition: 1 = retained, 2 = discard, 0 = total composition
## ## Maturity state: 1 = immature, 2 = mature, 0 = both states combined
## ## Shell condition: 1 = new shell, 2 = old shell, 0 = both shell types combined
## ##length proportions of pot discarded males
## ##Year, Seas, Fleet, Sex, Type, Shell, Maturity, Nsamp, DataVec
## 1990 2 1 1 0 0 0 15 0.1133 0.3933 0.4933
## 1991 2 1 1 0 0 0 25 0.1329 0.1768 0.6902
## 1992 2 1 1 0 0 0 25 0.1905 0.2677 0.5417
## 1993 2 1 1 0 0 0 25 0.2807 0.2097 0.5096
## 1994 2 1 1 0 0 0 25 0.2942 0.2714 0.4344
## 1995 2 1 1 0 0 0 25 0.1478 0.2127 0.6395
## 1996 2 1 1 0 0 0 25 0.1595 0.2229 0.6176
## 1997 2 1 1 0 0 0 25 0.1818 0.2053 0.6128
## 1998 2 1 1 0 0 0 25 0.1927 0.2162 0.5911
## 2009 2 1 1 0 0 0 50 0.1413 0.3235 0.5352
## 2010 2 1 1 0 0 0 50 0.1314 0.3152 0.5534
## 2011 2 1 1 0 0 0 50 0.1314 0.3051 0.5636
## 2012 2 1 1 0 0 0 50 0.1417 0.3178 0.5406

```

```

## 2014 2 1 1 0 0 0 50 0.0939 0.2275 0.6786
## 2015 2 1 1 0 0 0 50 0.1148 0.2518 0.6333
## ##length proportions of trawl survey males
## ##Year, Seas, Fleet, Sex, Type, Shell, Maturity, Nsamp, DataVec
## 1978 1 4 1 0 0 0 50 0.3865 0.3478 0.2657
## 1979 1 4 1 0 0 0 50 0.4281 0.3190 0.2529
## 1980 1 4 1 0 0 0 50 0.3588 0.3220 0.3192
## 1981 1 4 1 0 0 0 50 0.1219 0.3065 0.5716
## 1982 1 4 1 0 0 0 50 0.1671 0.2435 0.5893
## 1983 1 4 1 0 0 0 50 0.1752 0.2726 0.5522
## 1984 1 4 1 0 0 0 50 0.1823 0.2085 0.6092
## 1985 1 4 1 0 0 0 46.5 0.2023 0.2010 0.5967
## 1986 1 4 1 0 0 0 23 0.1984 0.4364 0.3652
## 1987 1 4 1 0 0 0 35.5 0.1944 0.3779 0.4277
## 1988 1 4 1 0 0 0 40.5 0.1879 0.3737 0.4384
## 1989 1 4 1 0 0 0 50 0.4246 0.2259 0.3496
## 1990 1 4 1 0 0 0 50 0.2380 0.2332 0.5288
## 1991 1 4 1 0 0 0 50 0.2274 0.3300 0.4426
## 1992 1 4 1 0 0 0 50 0.2263 0.2911 0.4826
## 1993 1 4 1 0 0 0 50 0.2296 0.2759 0.4945
## 1994 1 4 1 0 0 0 50 0.1989 0.2926 0.5085
## 1995 1 4 1 0 0 0 50 0.2593 0.3005 0.4403
## 1996 1 4 1 0 0 0 50 0.1998 0.3054 0.4948
## 1997 1 4 1 0 0 0 50 0.1622 0.3102 0.5275
## 1998 1 4 1 0 0 0 50 0.1276 0.3212 0.5511
## 1999 1 4 1 0 0 0 26 0.2224 0.2214 0.5562
## 2000 1 4 1 0 0 0 30.5 0.2154 0.2180 0.5665
## 2001 1 4 1 0 0 0 45.5 0.2253 0.2699 0.5048
## 2002 1 4 1 0 0 0 19 0.1127 0.2346 0.6527
## 2003 1 4 1 0 0 0 32.5 0.3762 0.2345 0.3893
## 2004 1 4 1 0 0 0 24 0.2488 0.1848 0.5663
## 2005 1 4 1 0 0 0 21 0.2825 0.2744 0.4431
## 2006 1 4 1 0 0 0 50 0.3276 0.2293 0.4431
## 2007 1 4 1 0 0 0 50 0.4394 0.3525 0.2081
## 2008 1 4 1 0 0 0 50 0.3745 0.2219 0.4036
## 2009 1 4 1 0 0 0 50 0.3057 0.4202 0.2741
## 2010 1 4 1 0 0 0 50 0.4081 0.3371 0.2548
## 2011 1 4 1 0 0 0 50 0.2179 0.3940 0.3881
## 2012 1 4 1 0 0 0 50 0.1573 0.4393 0.4034
## 2013 1 4 1 0 0 0 37 0.2100 0.2834 0.5065
## 2014 1 4 1 0 0 0 50 0.1738 0.3912 0.4350
## 2015 1 4 1 0 0 0 50 0.2340 0.2994 0.4666
## 2016 1 4 1 0 0 0 50 0.2255 0.2780 0.4965
## 2017 1 4 1 0 0 0 50 0.0849 0.2994 0.6157
## ##length proportions of pot survey
## ##Year, Seas, Fleet, Sex, Type, Shell, Maturity, Nsamp, DataVec
## 1995 1 5 1 0 0 0 100 0.1594 0.2656 0.5751
## 1998 1 5 1 0 0 0 100 0.0769 0.2205 0.7026
## 2001 1 5 1 0 0 0 100 0.1493 0.2049 0.6457
## 2004 1 5 1 0 0 0 100 0.0672 0.2484 0.6845
## 2007 1 5 1 0 0 0 100 0.1257 0.3148 0.5595
## 2010 1 5 1 0 0 0 100 0.1299 0.3209 0.5492
## 2013 1 5 1 0 0 0 100 0.1556 0.2477 0.5967
## 2015 1 5 1 0 0 0 100 0.0706 0.2431 0.6859

```



[illegible]

The base model control file:

```
## ## ----- ##
## ## LEADING PARAMETER CONTROLS ##
## # Controls for leading parameter vector theta
## # LEGEND FOR PRIOR:
## #          0 -> uniform #          1 -> normal #          2 -> lognormal
## #          3 -> beta
## #          4 -> gamma
## # ntheta
##    12
## ## ----- ##
## # ival      lb      ub      phz    prior      p1      p2      # parameter      #
##    0.18      0.01      1      -4      2      0.18      0.02      # M
##    14.3      -7.0      30      -2      0      -7      30      # log(R0)
##    10.0      -7.0      20      -1      1     -10.0      20      # log(Rini)
##    14.13979    7.0      16      1      0      7.0      16.      # log(Rbar)
##    80.0      30.0      310     -2      1     72.5      7.25     # Recruitment size distribution
##    0.25      0.1      7      -4      0      0.1      9.0      # Recruitment size scale (varian
##    0.2      -10.0      0.75     -4      0     -10.0      0.75     # log(sigma_R)
##    0.75      0.20      1.00     -2      3      3.0      2.00     # steepness
##    0.01      0.00      1.00     -3      3      1.01      1.01     # recruitment autocorrelation
##    14.9      10.00     15.00      3      0      5.00     20.00     # logNO vector of initial number
##    14.5      10.00     15.00      3      0      5.00     20.00     # logNO vector of initial number
##    14.3      10.00     15.00      3      0      5.00     20.00     # logNO vector of initial number
## ## GROWTH PARAM CONTROLS ##
## ## Two lines for each parameter if split sex, one line if not ##
## ## number of molt periods
```

```

## 1
## ## Year(s) molt period changes (blank if no changes)
##
## ## ----- ##
## # ival      lb      ub      phz  prior    p1    p2      # parameter      #
## 14.1      10.0    30.0     -3     0    0.0  999.0    # alpha males or combined
## 0.0001     0.0     0.01    -3     0    0.0  999.0    # beta males or combined
## 0.45       0.01     1.0     -3     0    0.0  999.0    # gscale males or combined
## 121.5      65.0    145.0    -4     0    0.0  999.0    # molt_mu males or combined
## 0.060      0.0     1.0     -3     0    0.0  999.0    # molt_cv males or combined
##
## ## ----- ##
## ## SELECTIVITY CONTROLS                                     ##
## ##      Each gear must have a selectivity and a retention selectivity. If a uniform      ##
## ##      prior is selected for a parameter then the lb and ub are used (p1 and p2 are      ##
## ##      ignored)                                           ##
## ## LEGEND                                                  ##
## ##      sel type: 0 = parametric, 1 = coefficients, 2 = logistic, 3 = logistic95,      ##
## ##                  4 = double normal (NIY)                                           ##
## ##      gear index: use +ve for selectivity, -ve for retention                             ##
## ##      sex dep: 0 for sex-independent, 1 for sex-dependent                             ##
## ## ----- ##
## ## ivector for number of year periods or nodes                                     ##
## ## POT      TBycatch FBycatch  NMFS_S  ADFG_pot
## ## Gear-1    Gear-2    Gear-3    Gear-4    Gear-5
## 2          1          1          1          1      # Selectivity periods
## 0          0          0          0          0      # sex specific selectivity
## 0          3          3          0          0      # male selectivity type
## ## Gear-1    Gear-2    Gear-3    Gear-4    Gear-5
## 1          1          1          1          1      # Retention periods
## 0          0          0          0          0      # sex specific retention
## 3          2          2          2          2      # male retention type
## 1          0          0          0          0      # male retention flag (0 -> no, 1 -> yes)
## ## gear  par  sel
## ## index index par sex  ival      lb  ub  prior    p1    p2  mirror period period  ##
## # Gear-1
## 1      1      1  0    0.4    0.001 1.0  0      0      1      3    1978  2008
## 1      2      2  0    0.7    0.001 1.0  0      0      1      3    1978  2008
## 1      3      3  0    1.0    0.001 2.0  0      0      1     -2    1978  2008
## 1      1      1  0    0.4    0.001 1.0  0      0      1      3    2009  2017
## 1      2      2  0    0.4    0.001 1.0  0      0      1      3    2009  2017
## 1      3      3  0    1.0    0.001 2.0  0      0      1     -2    2009  2017
## # Gear-2
## 2      7      1  0    40     10.0 200  0     10     200  -3    1978  2017
## 2      8      2  0    60     10.0 200  0     10     200  -3    1978  2017
## # Gear-3
## 3      9      1  0    40     10.0 200  0     10     200  -3    1978  2017
## 3     10      2  0    60     10.0 200  0     10     200  -3    1978  2017
## # Gear-4
## 4      8      1  0    0.7     0.001 1.0  0      0      1      4    1978  2017
## 4      9      2  0    0.7     0.001 1.0  0      0      1      4    1978  2017
## 4     10      3  0    0.9     0.001 1.0  0      0      1     -2    1978  2017
## # Gear-5
## 5     11      1  0    0.4     0.001 1.0  0      0      1      4    1978  2017

```

```

##      5      12      2      0      0.7      0.001 1.0      0      0      1      4      1978      2017
##      5      13      3      0      1.0      0.001 2.0      0      0      1     -2      1978      2017
## ## Retained
## # Gear-1
##     -1      14      1      0     120     100     200      0      1     900     -1      1978      2017
##     -1      15      2      0     123     110     200      0      1     900     -1      1978      2017
## # Gear-2
##     -2      16      1      0     595      1     700      0      1     900     -3      1978      2017
##     -2      17      2      0      10      1     700      0      1     900     -3      1978      2017
## # Gear-3
##     -3      18      1      0     590      1     700      0      1     900     -3      1978      2017
##     -3      19      2      0      10      1     700      0      1     900     -3      1978      2017
## # Gear-4
##     -4      20      1      0     580      1     700      0      1     900     -3      1978      2017
##     -4      21      2      0      20      1     700      0      1     900     -3      1978      2017
## # Gear-5
##     -5      22      1      0     580      1     700      0      1     900     -3      1978      2017
##     -5      23      2      0      20      1     700      0      1     900     -3      1978      2017
##
## ## ----- ##
## ## PRIORS FOR CATCHABILITY
## ##      If a uniform prior is selected for a parameter then the lb and ub are used (p1 ##
## ##      and p2 are ignored). ival must be > 0 ##
## ## LEGEND ##
## ##      prior: 0 = uniform, 1 = normal, 2 = lognormal, 3 = beta, 4 = gamma ##
## ## ----- ##
## ## LAMBDA: Arbitrary relative weights for each series, 0 = do not fit.
## ## SURVEYS/INDICES ONLY
## ## ival      lb      ub      phz      prior      p1      p2      Analytic?      LAMBDA
## ## 1.0        0        2      -1        0        0      9.0      0          1      # NMFS trawl
## ## 0.00411135867487 0 5      1        0        0      9.0      0          1      # ADF&G pot
## ## ----- ##
## ##
## ## ----- ##
## ## ADDITIONAL CV FOR SURVEYS/INDICES
## ##      If a uniform prior is selected for a parameter then the lb and ub are used (p1 ##
## ##      and p2 are ignored). ival must be > 0 ##
## ## LEGEND ##
## ##      prior: 0 = uniform, 1 = normal, 2 = lognormal, 3 = beta, 4 = gamma ##
## ## ----- ##
## ## ival      lb      ub      phz      prior      p1      p2
## ## 0.0000001      0.00000001 10.0      -4      4      1.0      100 # NMFS
## ## 0.0000001      0.00000001 10.0      -4      4      1.0      100 # ADF&G
## ## ----- ##
## ##
## ## ----- ##
## ## PENALTIES FOR AVERAGE FISHING MORTALITY RATE FOR EACH GEAR
## ## ----- ##
## ## Mean_F      STD_PHZ1      STD_PHZ2      PHZ
## ## 0.2          0.05          50.0          1 # Pot
## ## 0.001        0.05          50.0          1 # Trawl
## ## 0.001        0.05          50.0          1 # Fixed
## ## 0.00         2.00         20.00         -1 # NMFS
## ## 0.00         2.00         20.00         -1 # ADF&G

```

```

## ## ----- ##
##
## ## ----- ##
## ## OPTIONS FOR SIZE COMPOSTION DATA (COLUMN FOR EACH MATRIX)
## ## ----- ##
## ## LIKELIHOOD OPTIONS
## ## -1) Multinomial with estimated/fixed sample size
## ## -2) Robust approximation to multinomial
## ## -3) logistic normal (NIY)
## ## -4) multivariate-t (NIY)
## ## -5) Dirichlet
## ## AUTOTAIL COMPRESSION
## ## pmin is the cumulative proportion used in tail compression.
## ## ----- ##
## # 1 1 1 # Type of likelihood
## # 2 2 2 # Type of likelihood
## # 5 5 5 # Type of likelihood
## # 0 0 0 # Auto tail compression (pmin)
## # 1 1 1 # Initial value for effective sample size multiplier
## # -4 -4 -4 # Phz for estimating effective sample size (if appl.)
## # 1 2 3 # Composition aggregator
## # 1 1 1 # LAMBDA
## ## ----- ##
##
## ## ----- ##
## ## TIME VARYING NATURAL MORTALIIY RATES
## ## ----- ##
## ## TYPE:
## ## 0 = constant natural mortality
## ## 1 = Random walk (deviates constrained by variance in M)
## ## 2 = Cubic Spline (deviates constrained by nodes & node-placement)
## ## 3 = Blocked changes (deviates constrained by variance at specific knots)
## ## 4 = Time blocks
## ## ----- ##
## ## Sex-specific? (0=no, 1=yes)
## # 0
## ## Type
## # 3
## ## Phase of estimation
## # 4
## ## STDEV in m_dev for Random walk
## # 10.0
## ## Number of nodes for cubic spline or number of step-changes for option 3
## # 2
## # 0 # Females (ignored if single sex...)
## ## Year position of the knots (vector must be equal to the number of nodes)
## # 1998 1999
## # 1976 1980 1985 1994 # Females (ignored if single sex...)
## ## ----- ##
##
## ## ----- ##
## ## OTHER CONTROLS
## ## ----- ##
## # 3 # Estimated rec_dev phase

```

```

## 3      # Estimated rec_ini phase
## 0      # VERBOSE FLAG (0 = off, 1 = on, 2 = objective func)
## 2      # Initial conditions (0 = Unfished, 1 = Steady-state fished, 2 = Free parameters)
## 1978   # First year for average recruitment for Bspr calculation
## 2016   # Last year for average recruitment for Bspr calculation
## 0.35   # Target SPR ratio for Bmsy proxy
## 1      # Gear index for SPR calculations (i.e. directed fishery)
## 1      # Lambda (proportion of mature male biomass for SPR reference points)
## 1      # Use empirical molt increment data (0 = FALSE, 1 = TRUE)
## 0      # Stock-Recruit-Relationship (0 = None, 1 = Beverton-Holt)
## ## EOF
## 9999

```