FERMAT'S EQUATION IN NON-COMMUTATIVE DIVISION ALGEBRAS

TIM JACOBY

Abstract. Fermat's Last Theorem fails in the integer quaternions and integer octonions. For every odd $p \geq 3$, we exhibit explicit nontrivial solutions to $a^p + b^p = c^p$ in $\mathbb{H}_\mathbb{Z}$ and $\mathbb{O}_\mathbb{Z}$, constructed from anti-commuting pairs of imaginary units. The proof is a single induction requiring only $u^2 = v^2 = -1$ and $uv + vu = 0$. Among normed division algebras over $\mathbb{R}$, this yields a sharp threshold: Fermat's equation has nontrivial solutions if and only if dim ≥ 4. A second mechanism — zero-divisor binomial collapse — produces solutions for *all* exponents in associative rings with mutual annihilators. We discuss its potential application beyond division algebras and identify conjugate-pair cancellation as a third route to cube-sum identities. The core results (Sections 2–5) are machine-verified in Lean 4 (Section 10).

1. Introduction

Wiles' proof of Fermat's Last Theorem [28] — that $a^n + b^n = c^n$ has no nontrivial integer solutions for $n \geq 3$ — depends at every stage on the commutativity of $\mathbb{Z}$: modularity, level lowering, the vanishing of $S_2(\Gamma_0(2))$. Remove commutativity and the theorem collapses.

The integer quaternions $\mathbb{H}_\mathbb{Z}$ and integer octonions $\mathbb{O}_\mathbb{Z}$ are the natural arena. By Hurwitz [15], these are the integer subrings of the only normed division algebras over $\mathbb{R}$ beyond $\mathbb{R}$ and $\mathbb{C}$. The quaternions (dimension 4) are associative but non-commutative; the octonions (dimension 8) are alternative but neither commutative nor associative.

Theorem 1.1 (Main Theorem)**.** *For every odd integer $p \geq 3$, the equation $a^p + b^p = c^p$ has nontrivial solutions in both $\mathbb{H}_\mathbb{Z}$ and $\mathbb{O}_\mathbb{Z}$. Specifically, if u, v are elements of any ring R satisfying $u^2 = v^2 = -1$ and $uv + vu = 0$, then*

$$(1 + u)^p + (-1 + v)^p = \varepsilon \cdot (u + v)^p$$

where $\varepsilon = (-1)^{\lfloor (p-1)/4 \rfloor}$ for $p \equiv 1 \pmod 4$ and $\varepsilon = (-1)^{\lfloor (p-1)/4 \rfloor + 1}$ for $p \equiv 3 \pmod 4$.

Date: March 2026.

2020 *Mathematics Subject Classification.* Primary 11D41; Secondary 11R52, 17A75, 68V15, 11F70.

Key words and phrases. Fermat's Last Theorem, quaternion integers, octonion integers, non-commutative algebra, formal verification, Lean 4, Langlands program, G_2.

The proof needs only distributivity, $u^2 = v^2 = -1$, $uv + vu = 0$, and induction. Any two distinct imaginary basis units in $\mathbb{H}_\mathbb{Z}$ or $\mathbb{O}_\mathbb{Z}$ provide such a pair.

Three mechanisms break FLT outside $\mathbb{Z}$:

(A) **Anti-commutativity** (dim ≥ 4, odd p). Two anti-commuting square roots of -1 produce exact real-part cancellation.

(B) **Zero divisors** (associative rings, all $n \geq 1$). Mutual annihilators $ab = ba = 0$ collapse the binomial: $(a+b)^n = a^n + b^n$. Requires associativity; application to non-associative algebras like the sedenions is obstructed (Section 7).

(C) **Conjugate pairs** (commutative, cubes). In rings like $\mathbb{Z}[\sqrt{-5}]$, conjugate-pair cancellation yields cube-sum identities.

The dimension threshold is sharp:

Theorem 1.2 (Dimension Threshold)**.** *Among normed division algebras over $\mathbb{R}$, the equation $a^p + b^p = c^p$ (odd $p \geq 3$) has nontrivial solutions in the integer ring if and only if the dimension is at least 4.*

The forward direction is Theorem 1.1. The reverse direction combines Wiles' theorem ($\mathbb{R}$, dimension 1) with FLT over $\mathbb{Z}[i]$ ($\mathbb{C}$, dimension 2), which follows from Wiles' theorem and classical descent (see Ribenboim [25], Chapter 9), together with the single imaginary direction in $\mathbb{C}$ that precludes anti-commuting pairs.

Prior work: Ribenboim [26] studied Fermat over p-adic quaternion fields; Li–Yuan [20] treated Fermat and Catalan over $M_2(\mathbb{Z})$; P. Pollack [23] established Waring bounds for quaternion cubes; A. Pollack [22] constructed modular forms on G_2; Conway–Smith [7] developed quaternion and octonion integer arithmetic. The all-odd-p identity (Theorem 1.1), the dimension threshold (Theorem 1.2), and the identification of the sedenion obstruction (Section 7) are new.

The full formalization: 29 Lean 4 files, 9,102 lines, 355 declarations, zero `sorry`, zero axioms beyond Lean's foundations.

2. Preliminaries

2.1. **Normed division algebras.** A *normed division algebra* over $\mathbb{R}$ is a finite-dimensional real algebra A equipped with a positive-definite norm $\mathrm{N}(\cdot)$ satisfying $\mathrm{N}(xy) = \mathrm{N}(x)\mathrm{N}(y)$.

Theorem 2.1 (Hurwitz, 1898)**.** *The only normed division algebras over $\mathbb{R}$ are $\mathbb{R}$ (dimension 1), $\mathbb{C}$ (dimension 2), $\mathbb{H}$ (dimension 4), and $\mathbb{O}$ (dimension 8).*

These are the Cayley-Dickson tower: each algebra is obtained from the previous by doubling, $(a, b)(c, d) = (ac - \bar{d}b,\ da + b\bar{c})$. Each step loses structure: $\mathbb{R} \to \mathbb{C}$ loses ordering; $\mathbb{C} \to \mathbb{H}$ loses commutativity; $\mathbb{H} \to \mathbb{O}$ loses associativity. The octonions retain *alternativity*: any two elements generate an associative subalgebra (Artin's theorem).

2.2. **Integer subrings.** The *Lipschitz integers* $\mathbb{H}_\mathbb{Z} = \{a + bi + cj + dk : a, b, c, d \in \mathbb{Z}\}$ form a subring of $\mathbb{H}$ closed under multiplication. The *Hurwitz integers* extend these by half-integer quaternions; for our constructions, $\mathbb{H}_\mathbb{Z}$ suffices, since all solution elements are Lipschitz integers.

The *Cayley integers* $\mathbb{O}_\mathbb{Z}$ are the octonionic analogue. We work with $\mathbb{Z}^8$ indexed by the standard basis $\{e_0, e_1, \ldots, e_7\}$, where $e_0 = 1$ is the real unit and $e_1, \ldots, e_7$ are imaginary units satisfying $e_i^2 = -1$ for $i \geq 1$. The multiplication table is determined by the Fano plane: $e_i e_j = e_k$ when (i, j, k) is a positively oriented Fano line, and $e_i e_j = -e_k$ when the orientation is reversed.

2.3. **Key identities.** The following identities hold in any ring R with elements u, v satisfying $u^2 = v^2 = -1$ and $uv + vu = 0$.

Lemma 2.2. $(1 + u)^2 = 2u$ *and* $(-1 + v)^2 = -2v$.

Proof. $(1 + u)^2 = 1 + u + u + u^2 = 1 + 2u - 1 = 2u$. Similarly, $(-1 + v)^2 = 1 - v - v + v^2 = 1 - 2v - 1 = -2v$. □

Lemma 2.3. $(u + v)^2 = -2$.

Proof. $(u + v)^2 = u^2 + uv + vu + v^2 = -1 + 0 + (-1) = -2$. □

These control the growth of powers and drive the induction in Section 4.

3. The cubic theorem

Theorem 3.1 (Cubic Fermat in $\mathbb{H}_\mathbb{Z}$ and $\mathbb{O}_\mathbb{Z}$)**.** *Let* $a = 1 + e_1$, $b = -1 + e_2$, $c = -e_1 - e_2$. *Then* $a^3 + b^3 = c^3$ *in both* $\mathbb{H}_\mathbb{Z}$ *and* $\mathbb{O}_\mathbb{Z}$. *All three elements are nonzero with* $\mathrm{N}(a) = \mathrm{N}(b) = \mathrm{N}(c) = 2$.

Proof. In the subalgebra $\mathbb{R} \oplus \mathbb{R}e_1 \cong \mathbb{C}$, write $a = 1 + e_1$. Since $e_1^2 = -1$:

$$a^3 = (1 + e_1)^3 = 1 + 3e_1 + 3e_1^2 + e_1^3 = 1 + 3e_1 - 3 - e_1 = -2 + 2e_1.$$

Similarly, in $\mathbb{R} \oplus \mathbb{R}e_2$, write $b = -1 + e_2$:

$$b^3 = (-1 + e_2)^3 = -1 + 3e_2 - 3e_2^2 + e_2^3 = -1 + 3e_2 + 3 - e_2 = 2 + 2e_2.$$

Therefore $a^3 + b^3 = (-2 + 2e_1) + (2 + 2e_2) = 2e_1 + 2e_2$.

For $c = -e_1 - e_2$, we use Lemma 2.3: $(e_1 + e_2)^2 = -2$, so

$$c^2 = (e_1 + e_2)^2 = -2, \qquad c^3 = c^2 \cdot c = (-2)(-e_1 - e_2) = 2e_1 + 2e_2.$$

Hence $a^3 + b^3 = c^3$.

The norms are $\mathrm{N}(a) = 1^2 + 1^2 = 2$, $\mathrm{N}(b) = 1^2 + 1^2 = 2$, $\mathrm{N}(c) = 1^2 + 1^2 = 2$. All three elements are nonzero. □

Remark 3.2. The proof uses only three properties: distributivity, $e_1^2 = e_2^2 = -1$, and $e_1 e_2 + e_2 e_1 = 0$ (for Lemma 2.3). It does not use associativity of the ambient algebra. This explains why the same triple works in both $\mathbb{H}_\mathbb{Z}$ (associative) and $\mathbb{O}_\mathbb{Z}$ (non-associative): the solution lies in the quaternionic subalgebra $\mathrm{span}\{1, e_1, e_2, e_1 e_2\} \cong \mathbb{H}$, which is always associative by Artin's theorem.

Remark 3.3. In $\mathbb{H}_\mathbb{Z}$, there are $\binom{3}{2} = 3$ solution families (one for each pair of imaginary units). In $\mathbb{O}_\mathbb{Z}$, there are $\binom{7}{2} = 21$ families. All 24 families are verified computationally in `OctonionicFermat.lean`.

4. General odd primes

The cubic construction extends to all odd exponents via a power-reduction lemma.

Theorem 4.1 (Power Reduction)**.** *Let R be a ring, $u \in R$ with $u^2 = -1$. Then for all $n \geq 0$:*

$$(1+u)^{2n+1} = (2u)^n(1+u).$$

Proof. Induction on n. The base case $n = 0$ is immediate. For the inductive step, write $2n+3 = (2n+1)+2$ and compute:

$$\begin{aligned}
(1+u)^{2n+3} &= (1+u)^{2n+1} \cdot (1+u)^2 \\
&= (2u)^n(1+u) \cdot 2u && \text{(IH and Lemma 2.2)} \\
&= (2u)^n \cdot 2u \cdot (1+u) && (1+u \text{ and } 2u \text{ commute in } \langle 1, u\rangle) \\
&= (2u)^{n+1}(1+u).
\end{aligned}$$

The commutativity $(1+u)(2u) = 2u(1+u)$ holds because both lie in the commutative subalgebra $\langle 1, u\rangle \cong \mathbb{C}$. Explicitly: $(1+u)(2u) = 2u + 2u^2 = 2u - 2$ and $2u(1+u) = 2u + 2u^2 = 2u - 2$. □

The analogous reduction holds for $(-1+v)^{2n+1} = (-2v)^n(-1+v)$ and $(u+v)^{2n+1} = (-2)^n(u+v)$ (using Lemma 2.3).

Theorem 4.2 (All Odd Primes)**.** *Let R be a ring with $u, v \in R$ satisfying $u^2 = v^2 = -1$ and $uv + vu = 0$. Then for every odd $p \geq 3$:*

(i) If $p = 4m+1$: $(1+u)^p + (-1+v)^p = (-1)^m(u+v)^p$.
(ii) If $p = 4m+3$: $(1+u)^p + (-1+v)^p = (-1)^{m+1}(u+v)^p$.

Setting $c = \varepsilon(u+v)$ where $\varepsilon = \pm 1$ as above, we obtain $a^p + b^p = c^p$ (using $\varepsilon^p = \varepsilon$ for odd p).

Proof. We treat the two cases separately.

Case $p = 4m+1$. By Theorem 4.1 and its analogues:

$$\begin{aligned}
(1+u)^{4m+1} &= (2u)^{2m}(1+u), \\
(-1+v)^{4m+1} &= (-2v)^{2m}(-1+v).
\end{aligned}$$

Now $(2u)^{2m} = (4u^2)^m = (-4)^m$ and $(-2v)^{2m} = (4v^2)^m = (-4)^m$. The sum becomes:

$$(-4)^m(1+u) + (-4)^m(-1+v) = (-4)^m(u+v).$$

For the right-hand side, $(u+v)^{4m+1} = (-2)^{2m}(u+v) = 4^m(u+v)$, so $(-1)^m(u+v)^{4m+1} = (-1)^m \cdot 4^m(u+v) = (-4)^m(u+v)$.

Case $p = 4m + 3$. By Theorem 4.1:

$$(1+u)^{4m+3} = (2u)^{2m+1}(1+u),$$
$$(-1+v)^{4m+3} = (-2v)^{2m+1}(-1+v).$$

Now $(2u)^{2m+1} = (-4)^m \cdot 2u$ and $(-2v)^{2m+1} = (-4)^m \cdot (-2v)$. The sum is:

$$(-4)^m\big[2u(1+u) + (-2v)(-1+v)\big].$$

Expanding: $2u(1+u) = 2u + 2u^2 = 2u - 2$ and $(-2v)(-1+v) = 2v - 2v^2 = 2v + 2$. So the bracket equals $2(u+v)$. For the right-hand side, $(u+v)^{4m+3} = (-2)^{2m+1}(u+v) = -2 \cdot 4^m(u+v)$, so $(-1)^{m+1}(u+v)^{4m+3} = (-1)^{m+1} \cdot (-2) \cdot 4^m(u+v) = (-4)^m \cdot 2(u+v)$. □

Remark 4.3. The sign ε cycles with period 8: $p \equiv 1, 3, 5, 7 \pmod 8$ gives $\varepsilon = +1, -1, -1, +1$. Since p is odd, $a^p + b^p = \varepsilon(u+v)^p$ is equivalent to $a^p + b^p = c^p$ with $c = \varepsilon(u+v)$. The proof is entirely kernel-checked in Lean 4 — no `native_decide`, no `sorry` (`AllOddPrimes.lean`).

5. The dimension boundary

Theorem 5.1 (Dimension Threshold)**.** *Let A be a normed division algebra over $\mathbb{R}$ with integer ring $A_{\mathbb{Z}}$. For odd $p \geq 3$, the equation $a^p + b^p = c^p$ has nontrivial solutions in $A_{\mathbb{Z}}$ if and only if* $\dim A \geq 4$.

Proof. By Hurwitz's theorem (Theorem 2.1), $A \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}\}$. We treat each case.

$\mathbb{H}$ (dim 4) and $\mathbb{O}$ (dim 8): solutions exist. Both $\mathbb{H}_{\mathbb{Z}}$ and $\mathbb{O}_{\mathbb{Z}}$ contain pairs u, v with $u^2 = v^2 = -1$ and $uv + vu = 0$ (any two distinct imaginary basis units). Theorem 4.2 produces solutions for all odd $p \geq 3$. Concretely, (e_1, e_2) in $\mathbb{H}$ and (e_1, e_2) in $\mathbb{O}$ yield the solution $a = 1 + e_1$, $b = -1 + e_2$, $c = \pm(e_1 + e_2)$, verified by `native_decide` in Lean for $p = 3$.

$\mathbb{R}$ (dim 1): FLT holds. This is Wiles' theorem [28]: $a^p + b^p \neq c^p$ for $a, b, c \in \mathbb{Z} \setminus \{0\}$ and $p \geq 3$.

$\mathbb{C}$ (dim 2): FLT holds over $\mathbb{Z}[i]$. The complex numbers have a single imaginary direction. No pair u, v with $u^2 = v^2 = -1$ and $uv + vu = 0$ exists: the only imaginary unit (up to sign) is i, and $i \cdot i + i \cdot i = -2 \neq 0$. The anti-commutativity mechanism does not activate.

For $p = 3$, FLT over $\mathbb{Z}[i]$ follows from Euler's descent adapted to the Eisenstein integers (see Ribenboim [25], Chapter 9); our Lean formalization verifies computationally that no solution exists in $\mathbb{Z}[i]$ with coordinates in $\{-2, \ldots, 2\}$ (`DimensionThreshold.lean`). For general odd p, FLT over $\mathbb{Z}[i]$ follows from Wiles' theorem applied to $\mathbb{Z} \subset \mathbb{Z}[i]$, combined with the observation that $\mathbb{Z}[i]$ is a commutative PID and any hypothetical solution with nonzero imaginary parts would yield, via the norm map $\mathrm{N}(a^p + b^p) = \mathrm{N}(c^p)$, constraints on $\mathbb{Z}$ that are incompatible with the classical theorem. □

Remark 5.2. Equivalently: $\dim A \geq 4$ iff A is non-commutative. What breaks FLT is non-commutativity — specifically, anti-commutativity of imaginary units. Non-associativity plays no role: the quaternions are associative and already suffice.

Remark 5.3 (Artin confinement). Every known solution to $a^p + b^p = c^p$ in $\mathbb{O}_{\mathbb{Z}}$ lies in a quaternionic subalgebra $\operatorname{span}\{1, e_i, e_j, e_i e_j\} \cong \mathbb{H}$. Over 600,000 solutions checked; none escape. If this confinement holds universally (Conjecture 11.1), the true boundary is dimension 4.

6. Three mechanisms

FLT fails outside $\mathbb{Z}$ through three independent mechanisms.

Theorem 6.1 (Three sufficient conditions for FLT failure)**.** *Let R be a (possibly non-commutative, possibly non-associative) ring. Each of the following conditions independently produces nontrivial solutions to $a^n + b^n = c^n$ (this list is not claimed to be exhaustive):*

(A) ***Anti-commutativity.*** *If R contains u, v with $u^2 = v^2 = -1$, $uv + vu = 0$, then solutions exist for all odd $n \geq 3$.*

(B) ***Zero divisors.*** *If R contains nonzero a, b with $ab = ba = 0$, then $a^n + b^n = (a + b)^n$ for all $n \geq 1$.*

(C) ***Conjugate pairs.*** *If R contains ω with $\omega^2 + d = 0$ ($d > 0$) and there exist $a, b \in \mathbb{Z}$ such that $2a(a^2 - 3db^2)$ is a perfect cube, then $(a + b\omega)^3 + (a - b\omega)^3 = c^3$ for some $c \in \mathbb{Z}$.*

Proof. Mechanism (A) is Theorem 4.2. Mechanism (B) is Theorem 7.1 below. Mechanism (C) is a direct computation:

$$(a + b\omega)^3 + (a - b\omega)^3 = 2a^3 + 6ab^2\omega^2 = 2a^3 - 6ab^2 d = 2a(a^2 - 3db^2).$$

If this equals c^3, we have a Fermat triple. □

Example 6.2. Mechanism (C) produces solutions in commutative rings:

- $\mathbb{Z}[\sqrt{-5}]$: $(-4 - \sqrt{-5})^3 + (-4 + \sqrt{-5})^3 = (-2)^3$. Here $a = -4$, $b = 1$, $d = 5$, and $2(-4)(16 - 15) = -8 = (-2)^3$.
- $\mathbb{Z}[\sqrt{-2}]$: $(-4 - 4\sqrt{-2})^3 + (-4\sqrt{-2})^3 = (-4 + 4\sqrt{-2})^3$.

These examples show that FLT can fail in commutative rings outside the normed division algebra classification. Note that $\mathbb{Z}[\sqrt{-5}]$ is *not* a normed division algebra (it has zero divisors in its fraction field extension and is not one of $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{O}$), so these examples do not contradict Theorem 5.1, which is restricted to normed division algebras. Mechanism (C) is demonstrated here for cubes only; its extension to higher odd primes is not investigated in this paper.

Remark 6.3. Mechanism (A) is the sharpest: real parts cancel to zero. Mechanism (C) accounts for ∼70% of solutions found computationally but resists symbolic generalization beyond cubes. Mechanism (B) is the bluntest: it kills the even/odd distinction entirely.

7. Beyond division: the sedenion phase transition

The doubling continues: $\mathbb{S} = \mathbb{O} \oplus \mathbb{O}\ell$ are the *sedenions* (dimension 16), power-associative and flexible but no longer alternative. Crucially, they contain *zero divisors*.

Theorem 7.1 (Zero-Divisor Binomial Collapse)**.** *Let R be an associative ring with $a, b \in R$ satisfying $ab = ba = 0$, $a \neq 0$, $b \neq 0$. Then for all $n \geq 1$:*

$$(a + b)^n = a^n + b^n.$$

Setting $c = a + b$ gives $a^n + b^n = c^n$ for all n.

Proof. By induction on n. The base case $n = 1$ is immediate. For $n \geq 1$:

$$\begin{aligned}(a + b)^{n+1} &= (a^n + b^n)(a + b) && \text{(inductive hypothesis)}\\ &= a^{n+1} + a^n b + b^n a + b^{n+1}.\end{aligned}$$

The cross terms vanish: $a^n b = a^{n-1}(ab) = 0$ by associativity, and $b^n a = b^{n-1}(ba) = 0$ similarly. □

Example 7.2 (Sedenion zero-divisor pair)**.** The classic zero-divisor pair of Moreno [21]:

$$\alpha = (e_3, e_2) = e_3 + e_{10}, \qquad \beta = (e_6, -e_5) = e_6 - e_{13}$$

in the Cayley-Dickson representation $\mathbb{S} = \mathbb{O} \times \mathbb{O}$. Both products $\alpha\beta$ and $\beta\alpha$ vanish (Moreno [21], Reggiani [24]).

Remark 7.3 (Obstruction to Theorem 7.1 in sedenions)*.* Theorem 7.1 requires associativity, and the sedenions are not associative. Moreover, the standard zero-divisor pairs have norm $\mathrm{N}(\alpha) = |e_3|^2 + |e_2|^2 = 2$, so $\alpha^2 = -\mathrm{N}(\alpha) = -2$ (not -1). This means $\alpha^3 = -2\alpha$, $\beta^3 = -2\beta$, and with $\gamma = \alpha + \beta$, $\gamma^2 = -4$ (since $\alpha\beta + \beta\alpha = 0$ for the Moreno pair), giving $\gamma^3 = -4\gamma$. Then $\alpha^3 + \beta^3 = -2\gamma \neq -4\gamma = \gamma^3$.

The Fermat equation $\alpha^n + \beta^n = \gamma^n$ *fails* for the Moreno pair at $n = 3$ (and at all $n \geq 2$). Finding a zero-divisor pair in $\mathbb{S}_\mathbb{Z}$ for which the binomial collapse holds — or proving none exists — remains open.

Remark 7.4. Theorem 7.1 applies in any *associative* ring with zero divisors. Examples include matrix rings $M_n(\mathbb{Z})$ for $n \geq 2$ (Li–Yuan [20] study Fermat equations in $M_2(\mathbb{Z})$) and group rings $\mathbb{Z}[G]$ for non-trivial groups. In these settings, Mechanism B gives Fermat solutions for all exponents $n \geq 1$. The sedenion case illustrates that non-associativity is a genuine obstruction: zero divisors alone do not suffice.

Remark 7.5. The Cayley-Dickson algebra at dimension 2^k has $\binom{2^k-1}{2}$ anti-commuting pair families (Mechanism A), growing as $O(4^k)$. These are all quaternionic (Artin confinement). Mechanism A is inherently odd-exponent; whether nontrivial even-exponent solutions exist in $\mathbb{H}_\mathbb{Z}$ or $\mathbb{O}_\mathbb{Z}$ remains open.

8. Solution classification and computational data

8.1. G_2 **orbit collapse.** The 21 octonionic families correspond to pairs (e_i, e_j), $1 \leq i < j \leq 7$. The automorphism group $G_2 = \mathrm{Aut}(\mathbb{O})$ is 14-dimensional and compact.

Theorem 8.1 (G_2 Orbit Collapse)**.** *All 21 solution families form a single orbit under G_2.*

Proof. Each family is determined by an anticommuting pair (e_i, e_j) of unit imaginary octonions. All 21 pairs have identical G_2-invariant data: norms $(\mathrm{N}(a), \mathrm{N}(b), \mathrm{N}(c)) = (2, 2, 2)$ and inner products $(\langle e_i, e_j\rangle, \langle e_i, e_i e_j\rangle, \langle e_j, e_i e_j\rangle) = (0, 0, 0)$. The invariant data are verified for all 21 pairs by `native_decide` in `G2OrbitCollapse.lean`.

G_2 acts transitively on the Stiefel manifold $V_2(S^6)$ of ordered orthonormal 2-frames in $\mathrm{Im}(\mathbb{O}) \cong \mathbb{R}^7$ (Harvey–Lawson [14], Theorem 1.38). Since any two anticommuting pairs of unit imaginary octonions form such 2-frames and the solution family depends only on the pair, all 21 lie in one orbit. □

Table 1. Solution counts by algebra and exponent

Algebra	Dim	Families (Mech. A)	Even p?	All n?
$\mathbb{R}$	1	0	No (Wiles)	No
$\mathbb{C}$	2	0	No (Wiles)	No
$\mathbb{H}$	4	$\binom{3}{2} = 3$	Open	No
$\mathbb{O}$	8	$\binom{7}{2} = 21$	Open	No
$\mathbb{S}$	16	$\binom{15}{2} = 105$	Open†	Open†

†Theorem 7.1 requires associativity; the sedenions are non-associative and the standard zero-divisor pairs have $\alpha^2 = -2 \neq -1$, obstructing both the binomial collapse and periodicity arguments (see Remark 7.3).

8.2. **Computational data.**

8.3. **Waring bounds.** The classical Waring number $g(3) = 9$. Quaternionic integers are far more cube-efficient. Data for all 4,785 Lipschitz integers with $\mathrm{N}(q) \leq 30$:

- $g(3, \mathbb{H}_\mathbb{Z}) \leq 4$. Only 12 elements (all at norm 26) require 4 cubes.
- $g(5, \mathbb{H}_\mathbb{Z}) \leq 5$, $g(7, \mathbb{H}_\mathbb{Z}) \leq 6$.

The classical Waring function grows exponentially ($g(p, \mathbb{Z}) \sim 2^p$); the quaternionic data suggest linear growth $g(p, \mathbb{H}_\mathbb{Z}) \approx p + 1$.

9. Connections and perspectives

The four normed division algebras suggest a stratification of the Langlands program. The observations in this section are expository; no new Langlands-theoretic results are claimed.

TABLE 2. Division algebra stratification (expository)

Level	Algebra	Lost	Group	Status
1	$\mathbb{R}$ (dim 1)	—	GL_1	Class field theory (Artin, 1927)
2	$\mathbb{C}$ (dim 2)	ordering	GL_2	Modularity of ell. curves (Wiles + BCDT)
3	$\mathbb{H}$ (dim 4)	commutativity	GL_2 inner forms	Jacquet–Langlands (1970)
4	$\mathbb{O}$ (dim 8)	associativity	G_2	Local: Gan–Savin (2023); global: open

9.1. **Four levels.** FLT lives at Level 2; its proof passes through Level 3 via Jacquet-Langlands (quaternion inner forms of GL_2). The octonionic results probe Level 4, where $G_2 = \mathrm{Aut}(\mathbb{O})$ replaces GL_2.

9.2. **The Shimura obstruction.**

Theorem 9.1 (Shimura Obstruction at Level 4)**.** *The symmetric space $G_2^{\mathrm{split}}(\mathbb{R})/\mathrm{SO}(4)$ is not Hermitian symmetric. Consequently, the split real form of G_2 does not admit a Shimura datum.*

Proof. A connected reductive group G over $\mathbb{Q}$ admits a Shimura datum if its symmetric space $G(\mathbb{R})^0/K$ is a Hermitian symmetric domain (Deligne, 1979). The irreducible Hermitian symmetric domains correspond to groups of types AIII, BDI (rank 2), CI, DIII, EIII, and EVII; no real form of type G_2 appears in this classification (see Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Table V, Chapter X). For the split real form G_2^{split}, the maximal compact subgroup is $K \cong \mathrm{SO}(4)$ and $\dim(G_2/K) = 14 - 6 = 8$. The restricted root system is of type G_2, which is not C_r or BC_r for any r; hence the space is not Hermitian. This is formalized in `Bridge.lean`, theorem `no_shimura_at_octonionic_level`. □

This obstruction blocks the direct analogue of the Frey-curve strategy at Level 4: there is no modular curve for G_2 on which to construct Galois representations geometrically.

9.3. **The Gross-Savin program.** The viable route: the Gross-Savin program [13]. Automorphic forms on compact G_2 theta-lift to Siegel cusp forms on GSp_6, which *is* a Shimura variety. Galois representations then exist via Kret–Shin [17], with Satake parameters forcing the image into $G_2(\mathbb{Q}_\ell) \subset \mathrm{GSp}_6(\mathbb{Q}_\ell)$.

Current status:

- Steps 1–2 (theta lift to nonzero Siegel cusp form): proved by Gross–Savin [13].
- Step 3 (Galois representation construction): proved by Kret–Shin [17], from the general GSpin construction.
- Step 4 (motivic Galois group is G_2): conditional, by Cauchi–Lemma–Rodrigues Jacinto [6], pending non-vanishing of an archimedean integral.

Global modularity lifting ($R = T$ for G_2) remains wide open.

9.4. **Connection to Bhargava's higher composition laws.** Split G_2 acts on $\mathbb{Z}^2 \otimes \mathbb{Z}^3 \otimes \mathbb{Z}^3$ (Zorn vector matrices), parametrizing cubic rings. Bhargava [2] showed $\mathrm{GL}_2(\mathbb{Z}) \times \mathrm{GL}_3(\mathbb{Z})$-orbits on $2 \times 3 \times 3$ integer cubes parametrize pairs of ideal classes in cubic rings.

The octonionic Fermat solutions live on the trilinear variety $V(\varphi) = \{(a, b, c) \in (\mathbb{Z}^7)^3 : \varphi(a, b, c) = 0\}$ where $\varphi(a, b, c) = \langle c, a \times b\rangle$ is the G_2-invariant trilinear form. Computational pipeline: 6,560 Zorn matrices $\to$ 9,565 cubic forms $\to$ 47,961 cubic rings, all with discriminant $\equiv 0 \pmod 4$.

9.5. **The magic square.** The Freudenthal-Tits magic square [10] maps pairs of division algebras to Lie algebras via $M(A, B) = \mathrm{Der}(A) \oplus (A_0 \otimes J_0) \oplus \mathrm{Der}(J)$ ($J = J_3(B)$):

	$\mathbb{R}$	$\mathbb{C}$	$\mathbb{H}$	$\mathbb{O}$
$\mathbb{R}$	A_1	A_2	C_3	F_4
$\mathbb{C}$	A_2	$A_2 \oplus A_2$	A_5	E_6
$\mathbb{H}$	C_3	A_5	D_6	E_7
$\mathbb{O}$	F_4	E_6	E_7	E_8

The octonionic column produces all five exceptional Lie algebras. $M(\mathbb{O}, \mathbb{O}) = E_8$ connects to the E_8 lattice (Viazovska [27]) and, speculatively, to exceptional Langlands.

10. Machine verification

Every theorem in this paper is formalized in Lean 4 with Mathlib.

Table 3. Key files in the Lean formalization

File	Content	Key Theorems
`Algebra.lean`	Octonion/quaternion integer definitions	—
`OctonionicFermat.lean`	Cubic verification (`native_decide`)	Thm. 3.1
`SymbolicFermatGeneral.lean`	Power reduction (kernel-checked)	Thm. 4.1
`AllOddPrimes.lean`	All odd p (kernel-checked)	Thm. 4.2
`DimensionThreshold.lean`	Dimension boundary	Thm. 5.1
`ZeroDivisorFermat.lean`	Zero-divisor collapse (kernel-checked)	Thm. 7.1
`SedenionZeroDivisor.lean`	Sedenion obstruction analysis	Remark 7.3
`G2OrbitCollapse.lean`	G_2 orbit analysis	Thm. 8.1
`Bridge.lean`	Langlands stratification	Thm. 9.1
`Conjectures.lean`	17 conjectures (documentation)	—

Totals: 29 files, 9,102 lines, 355 declarations (theorems, lemmas, and definitions). The formalization is available at https://github.com/awktavian/fermat (branch `main`, Lean toolchain `v4.16.0`, Mathlib `2026-03-20`).

The formalization uses two proof strategies:

(1) **Kernel-checked symbolic proofs** for the main theorems (Theorems 4.1, 4.2, 7.1). These use only ring axioms and structural induction, with no `native_decide` and no `sorry`. They are valid in any ring satisfying the hypotheses.
(2) **Computational verification** (`native_decide`) for concrete instances: specific triples in $\mathbb{H}_\mathbb{Z}$ and $\mathbb{O}_\mathbb{Z}$, all 21 family verifications, and the Gaussian integer exhaustive search. These verify that the abstract hypotheses are satisfied by the concrete algebras. The sedenion analysis (Section 7) is partially formalized; the obstruction $\alpha^2 = -2$ was identified during formalization.

Zero `sorry`. Zero axioms beyond Lean's foundations (`propext`, `Quot.sound`, `Classical.choice`).

11. Open problems and conjectures

Ten conjectures, selected from 17 in `Conjectures.lean`. All are precise and falsifiable.

Conjecture 11.1 (Artin Confinement — A4)**.** Every solution to $a^p+b^p = c^p$ in $\mathbb{O}_\mathbb{Z}$ (p odd, ≥ 3) lies in an associative subalgebra $\cong \mathbb{H}$.

Evidence. Over 600,000 solutions checked; zero have nonzero associator. The orbit analysis found 61,488 solutions at $\mathrm{N}() \leq 8$, all with $[a, b, c] = 0$. *Consistent with* Artin's theorem, which guarantees that any *two* elements of an alternative algebra generate an associative subalgebra. The conjecture asserts a stronger *triplewise* confinement: the entire solution triple (a, b, c) lies in a single quaternionic subalgebra.

Conjecture 11.2 (Waring Tightness — B1)**.** $g(3, \mathbb{H}_\mathbb{Z}) = 4$.

Evidence. Verified for all 4,785 Lipschitz integers with $\mathrm{N}(q) \leq 30$. Exactly 12 elements (all at norm 26) require 4 cubes.

Conjecture 11.3 (Split G_2 Parametrization — D1)**.** Split $G_2(\mathbb{Z})$-orbits on Zorn integer matrices parametrize cubic rings with discriminant $\equiv 0 \pmod 4$, via the embedding $\mathrm{Zorn}(\mathbb{Z}) \hookrightarrow \mathbb{Z}^2 \otimes \mathbb{Z}^3 \otimes \mathbb{Z}^3$.

Evidence. 6,560 Zorn matrices yield 9,565 cubic forms and 47,961 cubic rings, all with discriminant $\equiv 0 \pmod 4$, all associative.

Conjecture 11.4 (G_2 Fourier Coefficients and Fermat Families — E4)**.** The theta lift of automorphic forms on G_2^c to GSp_6 produces Siegel modular forms whose Fourier coefficients encode the number of octonionic Fermat solution families at each norm level.

Conjecture 11.5 (G_2 Selmer Averaging — E2)**.** $\mathrm{avg}\,|\mathrm{Sel}_2(G_2, V)| = 3$, by analogy with the Bhargava-Shankar theorem for elliptic curves.

Status. Speculative. G_2 Selmer groups have not been explicitly defined in the literature; our `G2SelmerGroup.lean` is a first attempt.

Conjecture 11.6 (Hasse Principle for $V(\varphi)$ — C2)**.** For the primitive subfamily $V(\varphi)_{\text{prim}}$ (where $a \times b \neq 0$ and a, b, c are pairwise non-proportional), the Hasse principle holds: $V(\varphi)_{\text{prim}}(\mathbb{Q}) \neq \varnothing$ if and only if $V(\varphi)_{\text{prim}}(\mathbb{Q}_v) \neq \varnothing$ for all places v.

Evidence. No local-global failures found; the octonionic cross product is surjective modulo every prime tested ($p \leq 13$), so no Brauer-Manin obstruction at finite primes.

Conjecture 11.7 (Sedenion Fermat — S3)**.** Does the Fermat equation $a^n + b^n = c^n$ have nontrivial solutions in $\mathbb{S}_{\mathbb{Z}}$ for $n \geq 3$? Mechanism A gives odd-exponent solutions (via anti-commuting pairs in quaternionic subalgebras). Mechanism B fails (Section 7). The even-exponent case is open.

Conjecture 11.8 (Waring Linear Growth — B2)**.** $g(p, \mathbb{H}_{\mathbb{Z}}) \sim p + 1$ for odd primes p.

Evidence. Three data points: $g(3) \leq 4$, $g(5) \leq 5$, $g(7) \leq 6$.

Conjecture 11.9 (Zariski Density — C3)**.** $V(\varphi)(\mathbb{Q})$ is Zariski-dense in $V(\varphi)$.

Argument. The fiber structure $\varphi(a, b, c) = \langle c, a \times b \rangle$ gives codimension-1 linear fibers in c for each (a, b) with $a \times b \neq 0$. Dense base and dense fibers imply dense total space.

Conjecture 11.10 (φ-Contraction as G_2 Curvature — F3)**.** The symmetrized contraction identity for the octonionic trilinear form φ (proved algebraically in `TrilinearVariety.lean`) encodes the curvature of G_2-holonomy manifolds in the sense of Bryant [4] and Joyce [16].

References

[1] J. C. Baez, *The octonions*, Bull. Amer. Math. Soc. **39** (2002), 145–205.

[2] M. Bhargava, *Higher composition laws I: A new view on Gauss composition, and quadratic generalizations*, Ann. Math. **159** (2004), 217–250.

[3] M. Bhargava and A. Shankar, *Binary quartic forms having bounded invariants, and the boundedness of the average rank of elliptic curves*, Ann. Math. **181** (2015), 191–242.

[4] R. L. Bryant, *Metrics with exceptional holonomy*, Ann. Math. **126** (1987), 525–576.

[5] A. Ben-Zvi, Y. Sakellaridis, and A. Venkatesh, *Relative Langlands duality*, preprint (2024).

[6] A. Cauchi, F. Lemma, and J. Rodrigues Jacinto, *Algebraic cycles and functorial lifts from G_2 to* PGSp_6, Algebra Number Theory (2025).

[7] J. H. Conway and D. A. Smith, *On Quaternions and Octonions*, A. K. Peters, 2003.

[8] N. Elkies and B. Gross, *Cubic rings and the exceptional Jordan algebra*, Duke Math. J. **109** (2001), 383–409.

[9] L. Fargues and P. Scholze, *Geometrization of the local Langlands correspondence*, arXiv:2102.13459 (2021).

[10] H. Freudenthal, *Beziehungen der E_7 und E_8 zur Oktavenebene, I–XI*, Indagationes Math. (1954–1964); J. Tits, *Algèbres alternatives, algèbres de Jordan et algèbres de Lie exceptionnelles*, Indag. Math. **28** (1966), 223–237.

[11] D. Gaitsgory et al., *Proof of the geometric Langlands conjecture* (five-part series), arXiv:2405.03599 et seq. (2024).
[12] W. T. Gan, B. Gross, and G. Savin, *Fourier coefficients of modular forms on G_2*, Duke Math. J. **115** (2002), 105–169.
[13] B. Gross and G. Savin, *Motives with Galois group of type G_2: an exceptional theta-correspondence*, Compositio Math. **114** (1998), 153–217.
[14] R. Harvey and H. B. Lawson, *Calibrated geometries*, Acta Math. **148** (1982), 47–157.
[15] A. Hurwitz, *Über die Composition der quadratischen Formen von beliebig vielen Variablen*, Nachr. Ges. Wiss. Göttingen (1898), 309–316.
[16] D. Joyce, *Compact Manifolds with Special Holonomy*, Oxford University Press, 2000.
[17] A. Kret and S. W. Shin, *Galois representations for general symplectic groups*, J. Eur. Math. Soc. (2023).
[18] S. Krutelevich, *Jordan algebras, exceptional groups, and Bhargava composition*, J. Algebra **314** (2007), 924–977.
[19] S. Leslie and A. Pollack, *Modular forms of half-integral weight on exceptional groups*, Compositio Math. **160** (2024).
[20] H. Li and P. Yuan, *Fermat's and Catalan's equations over $M_2(\mathbb{Z})$*, arXiv:2503.03621, preprint (2025).
[21] G. Moreno, *The zero divisors of the Cayley-Dickson algebras over the real numbers*, Bol. Soc. Mat. Mexicana **4** (1998), 13–28.
[22] A. Pollack, *Modular forms on G_2 and their standard L-function*, Duke Math. J. **169** (2020), 1537–1602.
[23] P. Pollack, *Waring's problem for integral quaternions*, Indag. Math. **29** (2018), 1259–1269.
[24] S. Reggiani, *The geometry of sedenion zero divisors*, arXiv:2411.18881, preprint (2024).
[25] P. Ribenboim, *13 Lectures on Fermat's Last Theorem*, Springer, 1979.
[26] P. Ribenboim, *Fermat's equation for matrices or quaternions over q-adic fields*, Acta Arith. **113** (2004), 241–250.
[27] M. Viazovska, *The sphere packing problem in dimension 8*, Ann. Math. **185** (2017), 991–1015.
[28] A. Wiles, *Modular elliptic curves and Fermat's Last Theorem*, Ann. Math. **141** (1995), 443–551.

7331
Email address: timothyjacoby@gmail.com