

Notes on Type Theory

[DRAFT: APRIL 10, 2025]

Steve Awodey

with contributions from Andrej Bauer

Contents

4	Homotopy Type Theory	5
4.1	Identity types	6
4.1.1	The naive interpretation	8
4.2	The groupoid model	9
4.3	Weak factorization systems	13
4.4	Natural models	13
4.4.1	Modeling the type formers	16
4.4.2	Strictification	16
4.5	Universes	16
	Bibliography	17

Chapter 4

Homotopy Type Theory

The extensional dependent type theory of the previous chapter is in some ways a very natural system that admits an intuitively clear model in the locally cartesian closed category of sets and related categories. But for computational purposes, and specifically for the important application of type theory to proof checking in a computer proof assistant such as Agda or Lean, it has some serious defects: the equality relation between terms (or types) is not *decidable*: there is no algorithm that will determine whether two closed terms of a given type $s, t : A$ are (judgementally) equal $s \equiv t : A$. Indeed, there is no normalization procedure for reducing terms to normal forms—otherwise we could use it to decide whether two terms were equal by normalizing them and then comparing their normal forms. Relatedly, one cannot effectively decide whether a given type (e.g. an equality type such as $\text{Eq}_A(s, t)$) is inhabited (which would be a decision procedure for the *provability* of $s =_A t$), even given a candidate “proof term” $p : \text{Eq}_A(s, t)$ (which would be a decision procedure for *being a proof*).

For this reason, the extensional system is often replaced in applications by a weaker one, called *intensional type theory*, which enjoys better computational behavior, such as decidability of equality and type-checking, and normalization. A good discussion of these and several related issues, such as canonicity and consistency can be found in Chapter 3 of the book [AG].

However, this is only one side of the story. The intensional theory was mainly a technical device for specialists in computational type theory (and a conceptual challenge from the semantic point of view) until around 2006, when it was discovered that this theory admitted a homotopical (and higher-categorical) interpretation, which led to the discovery of Homotopy type theory (HoTT) [Awo12]. This interpretation not only helped to clarify the intensional theory, and prove useful in investigating its computational properties, but also opened up a wide range of applications outside of the conventional areas of type theory, *vis.* computational and constructive mathematics. For, quite independently of such applications, the homotopical interpretation permits the use of intensional type theory as a powerful and expressive *internal language* for formal reasoning in homotopy theory and higher category theory, both highly abstract areas of mathematics, for which new and rigorous tools for calculation and proof are quite welcome. Moreover, the fortuitous fact that

this system also has the good computational behavior that it does has led to the use of computational proof assistants in homotopy theory and higher category theory, even ahead of some more down-to-earth branches of mathematics, where such exotic semantics were not needed.

The homotopical interpretation was already anticipated by a 2-dimensional one in the category of groupoids, a special case of a higher categorical model that already suffices to make some of the essential features of such models clear. Thus we shall briefly review this model below, after introducing the intensional theory, and before considering the general homotopical semantics using weak factorization systems. Such “weak” interpretations also bring to a head the coherence issues that we deferred in the previous chapter, and we conclude with one approach to strictifying such interpretations using natural models, aka, categories with families.

4.1 Identity types

We begin by recalling from Section ?? the rules for *equality* types in the *extensional* system: The formation, introduction, elimination, and computation rules for equality types were as follows:

$$\begin{array}{c} \frac{s : A \quad t : A}{s =_A t \text{ type}} \qquad \frac{a : A}{\mathbf{refl}(a) : (a =_A a)} \\[1em] \frac{p : s =_A t}{s \equiv t : A} \qquad \frac{p : s =_A t}{p \equiv \mathbf{refl}(s) : (s =_A s)} \end{array}$$

The *Identity types* in the intensional theory, also written $x =_A y$, or sometimes $\mathbf{Id}_A(x, y)$, have the same formation and introduction rules as the Equality types, but the *elimination rule* of “equality reflection” is replaced by the following elimination rule:

$$\frac{x : A, y : A, z : \mathbf{Id}_A(x, y) \vdash C(x, y, z) \text{ type}, \quad x : A \vdash c(x) : C(x, x, \mathbf{refl}(x))}{x : A, y : A, z : \mathbf{Id}_A(x, y) \vdash J(x, y, z, c) : C}$$

in which the variable x is bound in the occurrence of c within the eliminator J . The associated *computation rule* then becomes:

$$x : A \vdash J(x, x, \mathbf{refl}(x), c) \equiv c(x) : C(x, x, \mathbf{refl}(x))$$

In HoTT, the elimination rule is called *path induction*, for reasons that will become clear.

To see how the elimination rule works, let us derive the basic laws of identity, namely reflexivity, symmetry, and transitivity, as well as Leibniz’s Law the *indiscernibility of identicals*, also known as the substitution of equals for equals.

- Reflexivity: states that $x =_A y$ is a reflexive relation, but this is just the \mathbf{Id} -formation and intro rules:

$$x : A, y : A \vdash x =_A y \text{ type}, \quad x : A \vdash \mathbf{refl}(x) : x =_A x$$

- Symmetry: can be stated as $x : A, y : A, u : x =_A y \vdash ? : y =_A x$, which can be proved with an **Id-elim** as follows:

$$\frac{x : A \vdash \mathbf{refl}(x) : x =_A x}{x : A, y : A, u : x =_A y \vdash J(x, y, u, \mathbf{refl}) : y =_A x}$$

- Transitivity: we wish to show

$$x : A, y : A, z : A, u : x =_A y, v : y =_A z \vdash ? : x =_A z$$

regarding $z : A$ as a fixed parameter, which we can move to the front of the context, we want to apply an **Id-elim** with respect to the assumption $u : x =_A y$, so we can set x to y , and look for a premiss of the form:

$$z : A, y : A, v : y =_A z \vdash ? : y =_A z$$

We cannot simply take v , however, since the order of the types in the context is still wrong for **Id-elim**, but we can move the assumption $v : y =_A z$ to the right with a λ -abstraction to obtain

$$z : A, y : A \vdash \lambda v. v : y =_A z \rightarrow y =_A z,$$

and now we *can* apply the planned **Id-elim** with respect to $u : x =_A y$ with the “motive” being $y =_A z \rightarrow x =_A z$ to obtain

$$z : A, y : A, x : A, u : x =_A y \vdash J(x, y, u, \lambda v. v) : y =_A z \rightarrow x =_A z$$

from which follows the desired

$$x : A, y : A, z : A, u : x =_A y, v : y =_A z \vdash J(x, y, u, \lambda v. v) v : x =_A z.$$

- Substitution: to show

$$\frac{x : A \vdash C(x) \text{ type}}{x : A, y : A, u : x =_A y \vdash ? : C(x) \rightarrow C(y)}$$

it suffices to have a premiss of the form

$$x : A \vdash c(x) : C(x) \rightarrow C(x)$$

for this, we can take $c(x) = \lambda z : C(x). z : C(x) \rightarrow C(x)$ to obtain

$$x : A, y : A, u : x =_A y \vdash J(x, y, u, x. \lambda z : C(x). z) : C(x) \rightarrow C(y).$$

Note that the variable x is bound in the **J** term.

Many more properties of Id -types and their associated J-terms are shown in the introductory texts [Uni13, Rij25]. One key fact is that the higher identity types $\text{Id}_{\text{Id}_A(a,b)}(p, q)$ are no longer degenerate, but themselves may have terms that are non-identical, i.e. not propositionally equal, leading to so-called *higher types*. This “failure of UIP” (uniqueness of identity proofs) in the intensional system was first shown using the groupoid model, which sheds considerable light on the intensional system.

Exercise 4.1.1. Show that given any $a, b, c : A$ and $p : a =_A b$ and $q : b =_A c$, one can define a composite $p \cdot q : a =_A c$ (using the transitivity of $=_A$). Then show that, for any $p : a =_A b$, the symmetry term $\sigma(p) : b =_A a$ satisfies the (propositional) equation $\sigma(p) \cdot p = \text{refl}$. Is either of $\sigma(p) \cdot p = \text{refl}$ or $\text{refl} \cdot \sigma(p) = \text{refl}$ judgemental? What about associativity of $p \cdot q$

Exercise 4.1.2. Show that $p \cdot q$ from the previous exercise is (propositionally) associative.

Exercise 4.1.3. Show that any term $f : A \rightarrow B$ acts on identities $p : a =_A b$, in the sense that there is a term $\text{ap}(f)(p) : fa =_B fb$. Is $\text{ap}(f)$ “functorial” (in the evident sense)?

Exercise 4.1.4. Observe that the Substitution property means that the assignment

$$(a : A) \mapsto C(a) \text{ type}$$

is functorial (in some sense). Is it strictly functorial?

4.1.1 The naive interpretation

We can try to interpret the (intensional) identity types in the naive way, as we did for extensional dependent type theory. This would give the formation and introduction rules as a type family $\text{Id}_A \rightarrow A \times A$ with a partial section over the diagonal substitution $\delta_A : (x : A) \rightarrow (x : A, y : A)$.

$$\begin{array}{ccc} & & \text{Id}_A \\ & \nearrow \text{refl} & \downarrow p \\ A & \xrightarrow{\delta_A} & A \times A \end{array}$$

where we are writing Id_A for the extended context (A, A, Id_A) and p for the dependent family $(x : A, y : A \vdash \text{Id}_A(x, y))$. The elimination rule then takes the form:

$$\begin{array}{ccc} A & \xrightarrow{c} & C \\ \text{refl} \downarrow & \searrow J & \downarrow \\ \text{Id}_A & \xlongequal{\quad} & \text{Id}_A \end{array} \quad (4.1)$$

for any type family $C \rightarrow \text{Id}_A$, with the computation rule asserting that the top triangle commutes (the bottom triangle commutes by the assumption that J is a section of $C \rightarrow \text{Id}_A$).

But now recall that in extensional type theory, *any* map $f : B \rightarrow A$ can be regarded as a type family over A , namely by taking the *graph factorization*

$$B \cong \Sigma_{a:A} \Sigma_{b:B} \mathbf{Eq}_A(a, fb) \longrightarrow B \times A.$$

So we can take the family C in the elimination to be $\mathbf{refl} : A \rightarrow \mathbf{Id}_A$, to obtain:

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ \mathbf{refl} \downarrow & \nearrow \mathbf{j} & \downarrow \mathbf{refl} \\ \mathbf{Id}_A & \xlongequal{\quad} & \mathbf{Id}_A \end{array}$$

We therefore get an iso $A \cong \mathbf{Id}_A$, making the identity type isomorphic to the extensional equality type $\mathbf{Eq}_A = A \rightarrow A \times A$.

Exercise 4.1.5. Prove that in the extensional theory, the graph factorization does indeed make any map $f : B \rightarrow A$ isomorphic to a family of types over its codomain. *Hint:* Consider the following two-pullback diagram.

$$\begin{array}{ccc} \Sigma_{a:A} \Sigma_{b:B} \mathbf{Eq}_A(a, fb) & \longrightarrow & A \\ \downarrow & \lrcorner & \downarrow \\ A \times B & \xrightarrow{A \times f} & A \times A \\ p_2 \downarrow & \lrcorner & \downarrow p_2 \\ B & \xrightarrow{f} & A \end{array}$$

4.2 The groupoid model

Exercises 4.1.1 – 4.1.4 from the last section suggest an interpretation of the intensional version of dependent type theory, namely with types as *groupoids* and type families as *functors*. Such an interpretation was first given by [HS98] in order to show that the principle of Uniqueness of Identity Proofs (UIP) – which holds in the extensional theory – indeed fails in the intensional one. We shall briefly sketch this result here.

In order to give a model of intensional type theory we should define what it means to be *be* a model of (intensional) type theory. We will do this in section ?? below – for now we simply describe a single model in the category **Gpd** of groupoids, which will turn out to be an instance of the general notion. For the extensional theory, we defined a model simply to be an interpretation into an LCCC \mathcal{E} , with contexts Γ interpreted as objects of \mathcal{E} , substitutions $\sigma : \Delta \rightarrow \Gamma$ as arrows of \mathcal{E} , type families $\Gamma \vdash A$ as objects of \mathcal{E}/Γ , and terms $\Gamma \vdash a : A$ as sections of the associated families. Substitution into families and terms was (weakly) interpreted as pullback (in the sense that there was an unresolved coherence issue), and the Σ and Π type formers were adjoints to pullback. Finally, the equality type

$x : A, y : A \vdash \mathbf{Eq}_A(x, y)$ was interpreted as the diagonal $A \rightarrow A \times A$.

$$\begin{array}{ccc} (x : A, y : A, z : \mathbf{Eq}_A(x, y)) & \xlongequal{\quad} & A \\ \downarrow & & \downarrow \\ (x : A, y : A) & \xlongequal{\quad} & A \times A \end{array}$$

We may simplify the notation for category of contexts by using Σ -types, writing e.g. $(x : A, y : A, z : \mathbf{Eq}_A(x, y)) = \Sigma_{x:A} \Sigma_{y:A} \mathbf{Eq}_A(x, y)$ or even \mathbf{Eq}_A , and $(x : A, y : A) = A \times A$, etc.

The groupoid interpretation of the intensional theory is based on the idea that the identity type of a type (interpreted as a groupoid) \mathbf{G} can be interpreted by the *path groupoid* of \mathbf{G} , which we shall write as $\mathbf{G}^!$.

Definition 4.2.1. If the groupoid $\mathbf{G} = G_0 \rightrightarrows G_1$ has objects G_0 and arrows G_1 , the *path groupoid* $\mathbf{G}^! = (|(\mathbf{G}^!)| \rightrightarrows |\mathbf{G}^!|)$ has as objects $|\mathbf{G}^!| = G_1$, and as arrows $|(\mathbf{G}^!)|$ the set of all commutative squares in \mathbf{G} , with the obvious source and target maps.

In other words, the path groupoid is the arrow category \mathbf{G}^\downarrow . Recall that the category \mathbf{Gpd} of (small) groupoids is a cartesian closed subcategory of \mathbf{Cat} , and that there is a *walking arrow* groupoid \mathbf{I} with exactly two objects and two (mutually inverse) non-identity arrows,

$$\mathbf{I} = \left(0 \begin{array}{c} \xleftarrow{\quad} \\ \xrightarrow{\quad} \end{array} 1 \right)$$

The notation $\mathbf{G}^!$ for the path groupoid is then correctly the exponential of \mathbf{G} by \mathbf{I} in \mathbf{Gpd} (and in \mathbf{Cat}). Observe that there are functors $\mathbf{dom}, \mathbf{cod} : \mathbf{G}^! \rightrightarrows \mathbf{G}$, as well as one $\mathbf{id} : \mathbf{G} \rightarrow \mathbf{G}^!$, making $\mathbf{G}^! \rightrightarrows \mathbf{G}$ into an internal groupoid in \mathbf{Gpd} , for any object \mathbf{G} . This will be our interpretation of the identity type of the type interpreted by \mathbf{G} .

More formally, we interpret:

- Contexts Γ : groupoids, i.e. objects of \mathbf{Gpd} ,
- Substitutions $\sigma : \Delta \rightarrow \Gamma$: homomorphisms of groupoids, i.e. arrows of \mathbf{Gpd} ,
- Types $\Gamma \vdash A$: functors $A : \mathbf{G} \rightarrow \mathbf{Gpd}$, where \mathbf{G} interprets Γ ,
- Terms $\Gamma \vdash a : A$: natural transformations $a : 1 \rightarrow A$ between functors, where 1 is the terminal functor in $\mathbf{Gpd}^{\mathbf{G}}$,
- Context extension $(\Gamma, A) \rightarrow \Gamma$: the Grothendieck construction $\int_{\mathbf{G}} A \rightarrow \mathbf{G}$.

In order to model the type formers Σ , Π , etc. of intensional type theory in \mathbf{Gpd} , we must deal with the fact that \mathbf{Gpd} is not locally cartesian closed, although it is cartesian closed. Recall that in order to model extensional type theory in presheaves we used the fact that $\mathbf{Set}^{\mathbf{C}^{\text{op}}}$ is always a CCC and that for any $P \in \mathbf{Set}^{\mathbf{C}^{\text{op}}}$ we have an equivalence

$$\mathbf{Set}^{\mathbf{C}^{\text{op}}}/_P \simeq \mathbf{Set}^{f^{P^{\text{op}}}},$$

and thus every slice is also a CCC. For groupoids, something similar is the case, but instead of the full slice category \mathbf{Gpd}/\mathbf{G} we use the subcategory of *fibrations* $\mathbf{Fib}_{\mathbf{G}} \hookrightarrow \mathbf{Gpd}/\mathbf{G}$, for which we have an equivalence

$$\mathbf{Fib}_{\mathbf{G}} \simeq \mathbf{Gpd}^{\mathbf{G}} \simeq \mathbf{Gpd}(\mathbf{Set}^{\mathbf{G}^{\text{op}}}),$$

Since the proof that \mathbf{Gpd} is a CCC doesn't depend on the classical logic of \mathbf{Set} , the category of internal groupoids in a topos like $\mathbf{Set}^{\mathbf{G}^{\text{op}}}$ is also a CCC. Thus we have that $\mathbf{Fib}_{\mathbf{G}}$ is a CCC for any groupoid \mathbf{G} .

Definition 4.2.2. A (split op-) fibration of groupoids $p : \mathbf{A} \rightarrow \mathbf{G}$ is a functor satisfying the condition: for every $a \in \mathbf{A}$ and $\gamma : pa \rightarrow g$ there is given a “lift” $\ell(a, \gamma) : a \rightarrow \tilde{g}$ with $p(\ell(a, \gamma)) = p$, and moreover,

1. $\ell(a, 1_{pa}) = 1_a : a \rightarrow a$,
2. for $\gamma' : g = p(\tilde{g}) \rightarrow h$, the lift of the composite is the composite of the lifts:

$$\ell(a, \gamma' \circ \gamma) = \ell(\tilde{g}, \gamma') \circ \ell(a, \gamma) : a \rightarrow \tilde{h}.$$

Proposition 4.2.3. *The category $\mathbf{Fib}_{\mathbf{G}}$ of fibrations of groupoids and functors $f : \mathbf{A} \rightarrow \mathbf{B}$ over \mathbf{G} that preserve the lifts is equivalent to the functor category $\mathbf{Gpd}^{\mathbf{G}}$.*

It follows in particular that each category $\mathbf{Fib}_{\mathbf{A}} \simeq \mathbf{Gpd}^{\mathbf{A}} \simeq \mathbf{Gpd}(\mathbf{Set}^{\mathbf{A}})$ is a CCC. The interpretation of a context Γ is now taken to be a groupoid \mathbf{G} , and the interpretation of a type family $\Gamma \vdash A$ is a functor $A : \mathbf{G} \rightarrow \mathbf{Gpd}$. The context extension $(\Gamma, A) \rightarrow \Gamma$ is interpreted by the fibration $\int_{\mathbf{G}} A \rightarrow \mathbf{G}$ given by the Grothendieck construction mediating the equivalence $\mathbf{Gpd}^{\mathbf{G}} \simeq \mathbf{Fib}_{\mathbf{G}}$.

For the base change functors along a fibration $p : \mathbf{A} \rightarrow \mathbf{G}$, we have adjoints as follows:

$$\begin{array}{ccc} \mathbf{A} & \mathbf{Gpd}^{\mathbf{A}} & \xrightarrow[\sim]{f} \mathbf{Fib}_{\mathbf{A}} \\ p \downarrow & p^* \uparrow & \Sigma \left(\begin{array}{c} \uparrow \\ p^* \\ \downarrow \end{array} \right) \Pi \\ \mathbf{G} & \mathbf{Gpd}^{\mathbf{G}} & \xrightarrow[\sim]{f} \mathbf{Fib}_{\mathbf{G}} \end{array}$$

For this, it still needs to be shown that:

1. the pullback of a fibration is a fibration,
2. the composite of fibrations is a fibration,
3. there is a pushed-forward fibration of a fibration along a fibration (right adjoint to pullback).

The proof uses the CCC structure in the categories $\mathbf{Fib}_{\mathbf{A}} \simeq \mathbf{Gpd}^{\mathbf{A}} \simeq \mathbf{Gpd}(\mathbf{Set}^{\mathbf{A}})$.

Identity types

To interpret the Id -type of a type (interpreted as, say) $\mathbf{A} = (A_1 \rightrightarrows A_0)$ in \mathbf{Gpd} (or indeed in any relative version $\mathbf{Gpd}(\mathbf{Set}^G)$), we shall use the path groupoid

$$\text{Id}_{\mathbf{A}} = \mathbf{A}^I \rightarrow \mathbf{A} \times \mathbf{A},$$

which is easily seen to be a fibration, and therefore corresponds to a functor $\text{Id}_{\mathbf{A}} : \mathbf{A} \times \mathbf{A} \rightarrow \mathbf{Gpd}$, namely that with *discrete groupoids* as its values:

$$\text{Id}_{\mathbf{A}}(a, b) = \{p : a \rightarrow b\} \subseteq A_1.$$

Note that for two objects $a, b \in \mathbf{A}$ there may be many different arrows $f : a \rightarrow b$ in $\text{Id}_{\mathbf{A}}(a, b)$, but for two such parallel arrows $f, g : a \rightrightarrows b$ in \mathbf{A} , regarded as objects in the path groupoid $\text{Id}_{\mathbf{A}} = \mathbf{A}^I$, there need be no arrow between them in the (discrete) groupoid $\text{Id}_{\mathbf{A}}(a, b)$; and indeed, there will be one (which is then unique) just if $f = g$. Thus we will have the desired violation of UIP, once we have shown that this interpretation satisfies the rules for intensional Id -types.

To show that, consider the diagram below, which we have already encountered as (4.2). We take any fibration $p : \mathbf{C} \rightarrow \text{Id}_{\mathbf{A}}$ and any section $c : \mathbf{A} \rightarrow \mathbf{C}$ over the insertion of identity arrows into the path groupoid $\text{refl} : \mathbf{A} \rightarrow \mathbf{A}^I = \text{Id}_{\mathbf{A}}$, and we need a diagonal filler J .

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{c} & \mathbf{C} \\ \text{refl} \downarrow & \searrow J & \downarrow p \\ \text{Id}_{\mathbf{A}} & \xlongequal{\quad} & \text{Id}_{\mathbf{A}} \end{array}$$

Since the diagram commutes by assumption, for any $a \in \mathbf{A}$ we have $pca = 1_a$. Let $\alpha : a \rightarrow b$ be any object in $\text{Id}_{\mathbf{A}} = \mathbf{A}^I$ and observe that there is always an arrow $\chi_\alpha : 1_a \rightrightarrows \alpha$ in $\text{Id}_{\mathbf{A}} = \mathbf{A}^I$, namely $(1_a, \alpha)$.

$$\begin{array}{ccc} a & \xrightarrow{1_a} & a \\ 1_a \downarrow & \chi_\alpha & \downarrow \alpha \\ a & \xrightarrow{\alpha} & b \end{array}$$

Since $p : \mathbf{C} \rightarrow \text{Id}_{\mathbf{A}}$ is a fibration, there is a lift $\ell(ca, \chi_\alpha) : ca \rightarrow \tilde{\alpha}$. We then set

$$J(\alpha) = \tilde{\alpha}$$

to obtain a functor $J : \text{Id}_{\mathbf{A}} \rightarrow \mathbf{A}$ making the two triangles in the diagram commute.

Exercise 4.2.4. Prove this!

Exercise 4.2.5. Show that the composition of fibrations $\mathbf{B} \rightarrow \mathbf{A}$ and $\mathbf{A} \rightarrow \mathbf{G}$ is a fibration. (This will be used for the interpretation of the type $\Gamma \vdash \Sigma_{\mathbf{A}} \mathbf{B}$, where $\Gamma \vdash \mathbf{A}$ and $\Gamma, \mathbf{A} \vdash \mathbf{B}$.)

Definition 4.4.1. For a small category \mathbb{C} , a natural transformation $f : Y \rightarrow X$ of presheaves on \mathbb{C} is called *representable* if for every $C \in \mathbb{C}$ and $x \in X(C)$, there is given a $D \in \mathbb{C}$, a $p : D \rightarrow C$, and a $y \in Y(D)$ such that the following square is a pullback.

$$\begin{array}{ccc} yD & \xrightarrow{y} & Y \\ yp \downarrow & \lrcorner & \downarrow f \\ yC & \xrightarrow{x} & X \end{array} \quad (4.2)$$

We will show that a representable natural transformation is essentially the same thing as a *category with families* in the sense of Dybjer [Dyb96]. Indeed, let us write the objects of \mathbb{C} as Γ, Δ, \dots and the arrows as $\sigma : \Delta \rightarrow \Gamma, \dots$, thinking of \mathbb{C} as a “category of contexts”. Let $u : \dot{\mathbf{U}} \rightarrow \mathbf{U}$ be a representable map of presheaves, and write its elements as:

$$\begin{aligned} A \in \mathbf{U}(\Gamma) & \quad \text{iff} \quad \Gamma \vdash A \\ a \in \dot{\mathbf{U}}(\Gamma) & \quad \text{iff} \quad \Gamma \vdash a : A, \end{aligned}$$

where $A = u \circ a$, as indicated in:

$$\begin{array}{ccc} & & \dot{\mathbf{U}} \\ & \nearrow a & \downarrow u \\ y\Gamma & \xrightarrow{A} & \mathbf{U} \end{array}$$

Thus we regard \mathbf{U} as the *presheaf of types*, with $\mathbf{U}(\Gamma)$ the set of all types in context Γ , and $\dot{\mathbf{U}}$ as the *presheaf of terms*, with $\dot{\mathbf{U}}(\Gamma)$ the set of all terms in context Γ , while the component $u_\Gamma : \dot{\mathbf{U}}(\Gamma) \rightarrow \mathbf{U}(\Gamma)$ is the typing of the terms in context Γ .

The naturality of $u : \dot{\mathbf{U}} \rightarrow \mathbf{U}$ just means that for any substitution $\sigma : \Delta \rightarrow \Gamma$, we have an action on types and terms:

$$\begin{aligned} \Gamma \vdash A & \quad \mapsto \quad \Delta \vdash A\sigma \\ \Gamma \vdash a : A & \quad \mapsto \quad \Delta \vdash a\sigma : A\sigma. \end{aligned}$$

While, by functoriality, given any further $\tau : \Phi \rightarrow \Delta$, we have

$$(A\sigma)\tau = A(\sigma \circ \tau) \quad (a\sigma)\tau = a(\sigma \circ \tau),$$

as well as

$$A1 = A \quad a1 = a$$

for the identity substitution $1 : \Gamma \rightarrow \Gamma$.

Finally, the representability of the natural transformation $p : E \rightarrow U$ is exactly the operation of *context extension*: given any $\Gamma \vdash A$, by Yoneda we have the corresponding map $A : y\Gamma \rightarrow \mathbf{U}$, and we let $p_A : \Gamma.A \rightarrow \Gamma$ be (the map representing) the pullback of u

along A , as in (4.2). We therefore have a pullback square:

$$\begin{array}{ccc} \mathbf{y}\Gamma.A & \xrightarrow{q_A} & \dot{\mathbf{U}} \\ \mathbf{y}p_A \downarrow & \lrcorner & \downarrow \mathbf{u} \\ \mathbf{y}\Gamma & \xrightarrow[A]{} & \mathbf{U} \end{array} \quad (4.3)$$

where the map $q_A : \Gamma.A \rightarrow \dot{\mathbf{U}}$ now determines a term

$$\Gamma.A \vdash q_A : Ap_A.$$

We may hereafter omit the \mathbf{y} for the Yoneda embedding, letting the Greek letters serve to distinguish representable presheaves and their maps.

Exercise 4.4.2. Show that the fact that (4.3) is a pullback means that given any $\sigma : \Delta \rightarrow \Gamma$ and $\Delta \vdash a : A\sigma$, there is a map

$$(\sigma, a) : \Delta \rightarrow \Gamma.A,$$

and this operation satisfies the equations

$$\begin{aligned} p_A \circ (\sigma, a) &= \sigma \\ q_A(\sigma, a) &= a, \end{aligned}$$

as indicated in the following diagram.

$$\begin{array}{ccccc} \Delta & & & & \\ & \searrow^{(\sigma, a)} & & \searrow^a & \\ & & \Gamma.A & \xrightarrow{q_A} & \dot{\mathbf{U}} \\ & & \downarrow p_A & \lrcorner & \downarrow \mathbf{u} \\ & & \Gamma & \xrightarrow[A]{} & \mathbf{U} \\ & \searrow^\sigma & & & \end{array}$$

Show moreover that the uniqueness of (σ, a) means that for any $\tau : \Delta' \rightarrow \Delta$ we also have:

$$\begin{aligned} (\sigma, a) \circ \tau &= (\sigma \circ \tau, a\tau) \\ (p_A, q_A) &= 1. \end{aligned}$$

Comparing the foregoing with the definition of a category with families in [Dyb96], we have shown:

Proposition 4.4.3. *Let $\mathbf{u} : \dot{\mathbf{U}} \rightarrow \mathbf{U}$ be a representable natural transformation of presheaves on a small category \mathbb{C} with a terminal object. Then \mathbf{u} determines a category with families, with \mathbb{C} as the contexts and substitutions, $\mathbf{U}(\Gamma)$ as the types in context Γ , and $\dot{\mathbf{U}}(\Gamma)$ as the terms in context Γ .*

Remark 4.4.4. A category with families is usually defined in terms of a presheaf

$$\mathbf{Ty} : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Set}$$

of types on the category \mathcal{C} of contexts, together with a presheaf

$$\mathbf{Tm}' : (\int_{\mathcal{C}} \mathbf{Ty})^{\mathrm{op}} \rightarrow \mathbf{Set}$$

of typed-terms on the category $\int_{\mathcal{C}} \mathbf{Ty}$ of types-in-context. We are using the equivalence of categories, valid for any category of presheaves $\mathbf{Set}^{\mathcal{C}^{\mathrm{op}}}$,

$$\mathbf{Set}^{\mathcal{C}^{\mathrm{op}}}/_P \simeq \mathbf{Set}^{(\int_{\mathcal{C}} P)^{\mathrm{op}}}$$

between the slice category over a presheaf P and the presheaves on its category of elements $\int_{\mathcal{C}} P$, to turn the presheaf $\mathbf{Tm}' : (\int_{\mathcal{C}} \mathbf{Ty})^{\mathrm{op}} \rightarrow \mathbf{Set}$ into one $\mathbf{Tm} : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Set}$ together with a map $\mathbf{Tm} \rightarrow \mathbf{Ty}$ in $\mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Set}$.

We think of a representable map of presheaves on an arbitrary category \mathbb{C} as a “type theory over \mathbb{C} ”, with \mathbb{C} as the category of contexts and substitutions. We will show in Section 4.4.2 that such a map of presheaves is essentially determined by a class of maps in \mathbb{C} that is closed under all pullbacks, corresponding to the “incoherent” interpretation of types in context as maps $A \rightarrow \Gamma$.

Definition 4.4.5. A *natural model of type theory* on a small category \mathbb{C} is a representable map of presheaves $\mathbf{u} : \dot{\mathbf{U}} \rightarrow \mathbf{U}$.

Exercise 4.4.6 (The natural model of syntax). Let \mathbb{T} be a dependent type theory and $\mathcal{C}_{\mathbb{T}}$ its category of contexts and substitutions. Define the presheaves $\mathbf{Ty} : \mathcal{C}_{\mathbb{T}}^{\mathrm{op}} \rightarrow \mathbf{Set}$ of types-in-context and $\mathbf{Tm} : \mathcal{C}_{\mathbb{T}}^{\mathrm{op}} \rightarrow \mathbf{Set}$ of terms-in-context, along with a natural transformation

$$\mathbf{tp} : \mathbf{Tm} \rightarrow \mathbf{Ty}$$

that takes a term to its type. Show that $\mathbf{tp} : \mathbf{Tm} \rightarrow \mathbf{Ty}$ is a natural model of type theory.

4.4.1 Modeling the type formers

4.4.2 Strictification

4.5 Universes

Bibliography

- [AG] C. Angiuli and D. Gratzer. Principles of dependent type theory. Online at <https://carloangiuli.com/courses/b619-sp24/notes.pdf>. Version 2024-11-26.
- [Awo12] Steve Awodey. Type theory and homotopy. In Peter Dybjer, Sten Lindström, Erik Palmgren, and Göran Sundholm, editors, *Epistemology Versus Ontology: Essays on the Philosophy and Foundations of Mathematics in Honour of Per Martin-Löf*, pages 183–201. Springer, 2012. arXiv:1010.1810.
- [Awo16] Steve Awodey. Natural models of homotopy type theory. *Mathematical Structures in Computer Science*, 28:1–46, 11 2016.
- [Dyb96] P. Dybjer. Internal type theory. *LNCS*, 1158:120–134, 1996.
- [Hof95] Martin Hofmann. Syntax and semantics of dependent types. In *Semantics and logics of computation*, volume 14 of *Publ. Newton Inst.*, pages 79–130. Cambridge University Press, Cambridge, 1995.
- [HS98] Martin Hofmann and Thomas Streicher. The groupoid interpretation of type theory. In *Twenty-five years of constructive type theory (Venice, 1995)*, volume 36 of *Oxford Logic Guides*, pages 83–111. Oxford Univ. Press, New York, 1998.
- [Rij25] Egbert Rijke. *Introduction to Homotopy Type Theory*. Cambridge University Press, 2025.
- [Uni13] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. <https://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.