

P7 Q3

- a) A 95% confidence interval for the "true" popularity of candidate A is given by (.43, .51)

We got this as

- $1 - \alpha/20$: significance level (well, 1 minus that)
- 4 percentage points: width of a confidence interval around the parameter of interest, at the stated level of significance.

- b) An approximate 95% CI for θ , the parameter of a $\text{Bern}(\theta)$ distⁿ, is

$$\left(\hat{\theta} - \sqrt{\frac{\hat{\theta}(1-\hat{\theta})}{n}} z_{1-\alpha/2}, \hat{\theta} + \sqrt{\frac{\hat{\theta}(1-\hat{\theta})}{n}} z_{1-\alpha/2} \right)$$

where $\alpha = 0.05$ and $z_{1-\alpha/2}$ is the value such that if $Z \sim N(0,1)$, $P(Z < z_{1-\alpha/2}) = 1 - \alpha/2$. For $\alpha = 0.05$, $z_{1-\alpha/2} = 1.96$. The previous question (part (a)) lets us write

$$\sqrt{\frac{\hat{\theta}(1-\hat{\theta})}{n}} z_{1-\alpha/2} = 0.04$$

where $\hat{\theta} = 0.47$, $z_{1-\alpha/2} = 1.96$.
Solving for n ,

$$n = \frac{\hat{\theta}(1-\hat{\theta})}{(0.04/z_{1-\alpha/2})^2} = \frac{0.47 \times 0.53}{(0.04/1.96)^2} \approx 598$$

c) For a sample with $\hat{\theta} = 0.47$, we need the width of the CI to be < 0.03 if we want it to not contain 0.50. Using the previous formula for n ,

$$n = \frac{0.47 \times 0.53}{(0.03 / 1.96)^2} = 1063.27$$

So we would need > 1064 respondents.

P7 Q5. (t-distⁿ)

$$Z \sim N(0,1), \quad f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$$

$$U \sim \chi_k^2, \quad f_U(u) = \frac{1}{2^{k/2} \Gamma(k/2)} u^{k/2-1} e^{-u/2}$$

Let $V = (U/k)^{1/2}$. Then $U = KV^2$, and

$$dU/dV = 2KV. \quad \text{where } U = KV^2$$

Changing variables: $f_V(v) = \left| \frac{dU}{dV} \right| f_U(u)$

$$= \frac{2KV}{2^{k/2} \Gamma(k/2)} [KV^2]^{k/2-1} e^{-KV^2/2}$$

$$= \left(\frac{2}{2^{k/2}} \right) \left(\frac{1}{\Gamma(k/2)} \right) (K \cdot K^{k/2-1}) (V \cdot V^{2(k/2-1)}) e^{-KV^2/2}$$

$$= \frac{K^{k/2}}{\Gamma(k/2) 2^{k/2-1}} V^{k-1} e^{-KV^2/2}$$

Now $T = \frac{Z}{\sqrt{U/K}} = \frac{Z}{V}$, the quotient of two independent random variables whose densities we know.

$$\begin{aligned} f_T(t) &= \int_0^\infty v f_V(v) f_Z(vt) dv \\ &= \int_0^\infty v \frac{K^{K/2}}{\Gamma(K/2) 2^{K/2-1}} v^{K-1} e^{-Kv^2/2} \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}(vt)^2} dv \\ &= \frac{K^{K/2}}{\Gamma(K/2) 2^{K/2-1}} \cdot \frac{1}{\sqrt{2\pi}} \int_0^\infty v^K e^{-\frac{1}{2}(K+t^2)v^2} dv. \end{aligned}$$

This almost looks like the gamma density as provided in the hint. Let $X = V^2$, $V = X^{1/2}$, $dv = \frac{1}{2} X^{-1/2} dx$

$$\begin{aligned} \Rightarrow &= \frac{K^{K/2}}{\Gamma(K/2) 2^{K/2-1}} \cdot \frac{1}{\sqrt{2\pi}} \int_0^\infty (X^{1/2})^K \cdot \frac{1}{2} X^{-1/2} e^{-\frac{1}{2}(K+t^2)X} dx \\ &= \frac{K^{K/2}}{\Gamma(K/2) 2^{K/2}} \cdot \frac{1}{\sqrt{2\pi}} \int_0^\infty X^{\frac{K-1}{2}} e^{-\frac{K}{2}(1+t^2/K)X} dx \\ &= \frac{K^{K/2}}{\Gamma(K/2) 2^{K/2}} \cdot \frac{1}{\sqrt{2\pi}} \Gamma\left(\frac{K-1}{2} + 1\right) \left(\frac{2}{K}\right)^{\frac{K-1}{2}} \left(1 + \frac{t^2}{K}\right)^{-(K+1)/2} \end{aligned}$$

~~Not a Gamma density~~

$$\begin{aligned} &= \frac{K^{K/2 - K/2 - 1/2}}{\sqrt{\pi} \times \Gamma(K/2)} \cdot \Gamma\left(\frac{K+1}{2}\right) \cdot 2^{\frac{K+1}{2} - \frac{K}{2} - \frac{1}{2}} \left(1 + \frac{t^2}{K}\right)^{-(K+1)/2} \\ &= \frac{\Gamma\left(\frac{K+1}{2}\right)}{\sqrt{K\pi} \Gamma(K/2)} \left(1 + \frac{t^2}{K}\right)^{-(K+1)/2} \end{aligned}$$

Though I do expect you to do it, note that I consider this to be a very tough derivation. Good job!

A8Q1

$$a) \text{ if } f(x) = \theta^x (1-\theta)^{n-x}$$

$$L(\theta) = \theta^{\sum x} (1-\theta)^{n-\sum x}$$

$$l(\theta) = \sum x \log \theta + (n - \sum x) \log(1-\theta)$$

$$S(\theta) = \partial l / \partial \theta = \sum x / \theta - \frac{n - \sum x}{1-\theta}$$

$$= 0$$

$$\Rightarrow \hat{\theta} = \frac{1}{n} \sum x_i = \bar{x}$$

$$J(\theta) = -\partial S / \partial \theta = \frac{\sum x}{\theta^2} + \frac{n - \sum x}{(1-\theta)^2}$$
$$= n \left(\frac{\bar{x}}{\theta^2} + \frac{1-\bar{x}}{(1-\theta)^2} \right)$$

$$I(\theta) = E J(\theta) = n \left(\frac{1}{\theta} + \frac{1}{1-\theta} \right) \quad (E(X) = \theta)$$
$$= \frac{n}{\theta(1-\theta)}$$

$$\Rightarrow \text{Var}(\hat{\theta}) \approx \frac{\hat{\theta}(1-\hat{\theta})}{n}$$

Approx 95% CI is

$$\left(\hat{\theta} - \sqrt{\text{Var}(\hat{\theta})} Z_{1-\alpha/2}, \hat{\theta} + \sqrt{\text{Var}(\hat{\theta})} Z_{1-\alpha/2} \right)$$

$$= \left(0.82 - \sqrt{\frac{0.82 \times 0.18}{150}} 1.96, 0.82 + \sqrt{\frac{0.82 \times 0.18}{150}} 1.96 \right)$$

$$= (0.814, 0.826)$$

$$e) E(X) = \frac{\theta_0}{2}, \text{Var}(X) = \frac{\theta_0^2}{12}, \text{SD}(X) = \frac{\theta_0}{\sqrt{12}}$$

From lecture 2, MOM estimator of θ is $\hat{\theta} = 2\bar{X}$

$$E(\hat{\theta}) = 2E(\bar{X}) = \theta_0$$

$$\text{Var}(\hat{\theta}) = 4\text{Var}(\bar{X}) = \frac{4}{12} \times \frac{\theta_0^2}{n} = \frac{\theta_0^2}{3n}$$

$$\text{SD}(\hat{\theta}) = \theta_0/\sqrt{3n}$$

$$\text{CLT for } \bar{X}: \frac{\bar{X} - E(\bar{X})}{\text{SD}(\bar{X})} \underset{\text{approx}}{\sim} N(0,1)$$

$$\Rightarrow 2\bar{X} = \hat{\theta} \underset{\text{approx}}{\sim} N(0, 2^2)$$

$$\text{A 95\% CI for } \bar{X} \text{ is } \left(\bar{X} - \text{SD}(\bar{X})z_{1-\alpha/2}, \bar{X} + \text{SD}(\bar{X})z_{1-\alpha/2} \right)$$

$$\text{So a 95\% for } \hat{\theta} = 2\bar{X} = \left(2\bar{X} - 2\text{SD}(\bar{X})z_{1-\alpha/2}, 2\bar{X} + 2\text{SD}(\bar{X})z_{1-\alpha/2} \right)$$

Q11) A8Q2

$$a) X \sim \text{unif}(0, \theta_0). \quad P(X < x) = \frac{x}{\theta_0}$$

$$\Rightarrow P(X/\theta_0 < x) = P(X < \theta_0 x) = \frac{\theta_0 x}{\theta_0} = x$$

which is recognized as the CDF of a $\text{unif}(0,1)$ random variable. Hence $X/\theta_0 \sim \text{Unif}(0,1)$

$$\begin{aligned} b) P\left(\frac{X_{(n)}}{\theta_0} < x\right) &= P\left(\frac{X_1}{\theta_0} < x \text{ and } \dots \text{ and } \frac{X_n}{\theta_0} < x\right) \\ &= x^n, \text{ as each } P\left(\frac{X_i}{\theta_0} < x\right) = x \text{ and they are independent.} \end{aligned}$$

which is recognized as the CDF of a $\text{Beta}(n,1)$ random variable. Hence $X_{(n)}/\theta_0 \sim \text{Beta}(n,1)$

c) The distribution function of $X_{(n)}/\theta_0$ is

$$P(X_{(n)}/\theta_0 < x) = F(x) = x^n, \quad 0 < x < 1$$

$$\text{So } P\left(\frac{1}{q} X_{(n)} < \theta_0 < q X_{(n)}\right) = 1 - \alpha$$

$$= P\left(\frac{1}{X_{(n)}} > \frac{1}{\theta_0} > \frac{1}{q X_{(n)}}\right)$$

$$= P\left(\frac{1}{q} < \frac{X_{(n)}}{\theta_0} < 1\right)$$

$$= F(1) - F(1/q) = 1 - \alpha$$

$$1^n - (1/q)^n = 1 - \alpha$$

$$(1/q)^n = \alpha$$

$$1/q = \alpha^{1/n}$$

$$\boxed{q = \alpha^{-1/n}}$$

d) $\bar{X} = 6.885$, $\hat{\theta}_{\text{mom}} = 2\bar{X} =$

So the CI from before = $(5.9789, 21.5611)$

$X_{(n)} = \text{Max}(X_i) = 13.35$. So we know

$\theta_0 \geq 13.35$, hence the 1st interval includes a bunch of nonsense (inadmissible) values.

2nd interval: $\alpha = 0.05$, $n = 10$, so $q = (0.05)^{-1/10} = 1.349$.

$\Rightarrow (13.35, 18.01)$

I much prefer this one; it is narrower and contains only possible values for θ_0 .

A8 Q5

$$a) \hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

$$y_i \sim N(\beta_0 + \beta_1 x_i, \sigma^2)$$

note $\bar{y} = \beta_0 + \beta_1 \bar{x} + \bar{\varepsilon} \sim N(\beta_0 + \beta_1 \bar{x}, \sigma^2/n)$, a normal random variable. Hence $y_i - \bar{y}$ is also normally distributed, with

$$E(y_i - \bar{y}) = \beta_1 (x_i - \bar{x})$$

$$\text{Var}(y_i - \bar{y}) = \sigma^2 + \sigma^2/n - 2\text{cov}(y_i, \bar{y})$$

$$\text{now, } \text{cov}(y_i, \bar{y}) = \frac{1}{n} \sum_{j=1}^n \text{cov}(y_i, y_j) = \sigma^2/n,$$

$$\text{so } \text{Var}(y_i - \bar{y}) = \sigma^2$$

$\hat{\beta}_1$ is another linear combination of normal random variables, so is again normal. We can get its mean and variance:

$$E \hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x}) E(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

$$= \frac{\sum_{i=1}^n (x_i - \bar{x}) \beta_1 (x_i - \bar{x})}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

$$= \beta_1$$

$$\text{Var}(\hat{\beta}_1) = \left(\frac{1}{\sum_{i=1}^n (x_i - \bar{x})^2} \right)^2 \sum_{i=1}^n (x_i - \bar{x})^2 \text{Var}(y_i - \bar{y})$$

$$= \left(\frac{1}{\sum_{i=1}^n (x_i - \bar{x})^2} \right)^2 \cdot \sum_{i=1}^n (x_i - \bar{x})^2 \cdot \sigma^2$$

$$= \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

$$\text{so } \hat{\beta}_1 \sim N\left(\beta_1, \frac{\sigma^2}{\sum_{i=1}^n (x_i - \bar{x})^2}\right) \quad \heartsuit$$

A9 Q3

$$X_{ij} \sim N(\mu_i, \sigma^2) \quad \sigma^2 \text{ known.}$$

$$i=1 \dots K, \quad j=1 \dots n$$

It's K groups, each of size n . Do they all have equal means?

Note the question is NOT "are their means all equal to some particular value?". We don't presuppose what μ_0 is - just that

$$H_0: \mu_1 = \dots = \mu_K = \mu_0$$

vs

$$H_1: \exists i \neq j \text{ with } \mu_i \neq \mu_j$$

a) Each $\mu_i \in \mathbb{R}$, so $\Omega = \mathbb{R} \times \dots \times \mathbb{R} = \mathbb{R}^K$

$$b) \Omega_0 = \{\mu_0\} \times \dots \times \{\mu_0\} = \{\mu_0\}^K$$

c) Composite. Under both, there are quantities to be estimated.

d) Under H_0 , we just have $X_{ij} \sim N(\mu_0, \sigma^2)$, which directly yields the result.

Sorry, I forgot e! ↘

f) Under H_1 , we have $X_{ij} \sim N(\mu_i, \sigma^2)$, so the likelihood over j for fixed i is

$$L_i(\mu_i) = \prod_{j=1}^n f_i(X_{ij}) = C \times \exp\left(-\frac{1}{2\sigma^2} \sum_{j=1}^n (X_{ij} - \mu_i)^2\right)$$

All the X_{ij} are mutually independent though, so the full likelihood is obtained as

$$L(\mu_1, \dots, \mu_K) = \prod_{i=1}^K L_i(\mu_i)$$

$$= C \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \mu_i)^2\right)$$

okay I'll do
e here \rightarrow

e) $X_{ij} \sim N(\mu_0, \sigma^2)$, so previous results from the course allow you to state that

$$\hat{\mu}_0 = \frac{1}{nK} \sum_{i=1}^K \sum_{j=1}^n X_{ij}$$

Don't let the double sum confuse you; we have nK independent observations from a $N(\mu_0, \sigma^2)$ distⁿ, so the MLE is the sample mean.

Also, on a test you would have to actually find the MLE by differentiating the log-likelihood, etc.

g) Note that μ_i only appears in the likelihood through $L_i(\mu_i)$, so all other terms can be treated as constant wrt μ_i . We had

$$L_i(\mu_i) = C \exp\left(-\frac{1}{2\sigma^2} \sum_{j=1}^n (X_{ij} - \mu_i)^2\right)$$

so unsurprisingly, $\hat{\mu}_i = \frac{1}{n} \sum_{j=1}^n X_{ij}$.

$$h) \Lambda = \frac{L_0(\mu)}{L(\mu)} = \frac{C \times \exp\left(\frac{1}{2\sigma^2} \sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \bar{X}_{..})\right)}{C \times \exp\left(\frac{1}{2\sigma^2} \sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \bar{X}_{i.})^2\right)}$$

$$\log \Lambda = \frac{1}{2\sigma^2} \sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \bar{X}_{..})^2 + \frac{1}{2\sigma^2} \sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \bar{X}_{i.})^2$$

$$-2 \log \Lambda = \frac{1}{\sigma^2} \left[\sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \bar{X}_{..})^2 - \sum_{i=1}^K \sum_{j=1}^n (X_{ij} - \bar{X}_{i.})^2 \right]$$

$$i) \dim \Omega_0 = 1, \dim \Omega = K$$

$$\Rightarrow -2 \log \Lambda \sim \chi_{K-1}^2$$

$$j) \text{ Let } W \sim \chi_{K-1}^2. \text{ Then}$$

$$R_\alpha = \{x: -2 \log \Lambda > W_{1-\alpha}\}$$

$$\text{and } P_0 = P(W > -2 \log \Lambda)$$

where $W_{1-\alpha}$ is the $1-\alpha$ quantile of the χ_{K-1}^2 distⁿ, ie

$$P(W < W_{1-\alpha}) = 1 - \alpha.$$

A10Q1

a) See lecture 10 slides.


b), c). In both these questions, $d\sqrt{n} \rightarrow \infty$.

$$\begin{aligned}\eta(d, n, \alpha) &= 1 - P(d\sqrt{n} - z_{1-\alpha/2} < Z < d\sqrt{n} + z_{1-\alpha/2}), \quad Z \sim N(0, 1) \\ &= 1 - (\Phi(d\sqrt{n} + z_{1-\alpha/2}) - \Phi(d\sqrt{n} - z_{1-\alpha/2})) \\ &\xrightarrow{d\sqrt{n} \rightarrow \infty} 1 - (1 - 1) \\ &= 1.\end{aligned}$$

as $\Phi(x)$ is strictly increasing and bounded above by 1

d) The normal distⁿ is symmetric, and its CDF satisfies

$$\Phi(-x) = 1 - \Phi(x)$$



$$\begin{aligned}\Rightarrow \eta(-d, n, \alpha) &= 1 - (\Phi(-d\sqrt{n} + z_{1-\alpha/2}) - \Phi(-d\sqrt{n} - z_{1-\alpha/2})) \\ &= 1 - (1 - \Phi(d\sqrt{n} - z_{1-\alpha/2}) - (1 - \Phi(d\sqrt{n} + z_{1-\alpha/2}))) \\ &= 1 - (\Phi(d\sqrt{n} + z_{1-\alpha/2}) - \Phi(d\sqrt{n} - z_{1-\alpha/2})) \\ &= \eta(d, n, \alpha)\end{aligned}$$

$$\begin{aligned}\text{e) } \eta(0, n, \alpha) &= 1 - (\Phi(z_{1-\alpha/2}) - \Phi(-z_{1-\alpha/2})) \\ &= 1 - (1 - \alpha) \\ &= \alpha.\end{aligned}$$