Contents

- Test computation of 1D specimen with 2D system.
- Compute 2D and 1D image of fluorescence microscope.
- Compute DIC image of phase object.
- Plot the results.

Test computation of 1D specimen with 2D system.

```
% Compare images computed by 1D sum over source and 2D sum over source.

% Specimen: double-slit.
x=[-3:0.01:3];
slitwidth=0.3;
specimen1d= single(abs(x-0.5*slitwidth)<1E-10 | abs(x+0.5*slitwidth)<1E-10);
specimen2d= single(repmat(specimen1d,length(x),1));</pre>
```

Compute 2D and 1D image of fluorescence microscope.

```
params.NAo=0.95;
params.Nac=0.95;
params.wavelength=0.55;
params.nEmbb=1;
params.nImm=1;

fluor=microlith(x,0);
fluor.computesys('Fluorescence',params);
fluor.computeimage(specimen2d,'CPU');


fluor1d=microlith(x,0);
fluor1d.computesys('Fluorescence',params);
fluor1d.computeimage(specimen1d,'CPU');
```


Compute DIC image of phase object.


```
params.NAo=0.95;
params.NAc=0.9;
params.wavelength=0.55;
params.nEmbb=1;
params.nImm=1;
params.shearangle=0;
params.shear=0.2;
params.bias=pi/4;
tic2D=tic;
DIC2d=microlith(x,0);
DIC2d.computesys('DIC',params);
DIC2d.Ic(DIC2d.mm>0)=0;
DIC2d.computeimage(exp(1i*specimen2d),'CPU');
time2D=toc(tic2D);
tic1D=tic;
DIC1d=microlith(x,0);
DIC1d.computesys('DIC',params);
DIC1d.Ic(DIC1d.mm>0)=0;
DIC1d.computeimage(exp(1i*specimen1d), 'CPU');
```


time1D=toc(tic1D);


Plot the results.














Published with MATLAB® R2013b