

Rapport d'analyses statistiques

Axelle Dupont, sbim, Hopital Saint Louis, Paris

September 20, 2017

Contents

1	Objectives	3
2	Methods	3
3	Results	4
3.1	Descriptive results	4
3.1.1	Patients characteristics	4
3.1.2	Treatments before alloSCT	5
3.1.3	Transplant conditions	6
3.1.4	Post-AlloSCT Response	9
3.2	Survival analysis in all patients	12
3.2.1	Overall survival, EFS and GRFS	12
3.2.2	TRM and cause of death	17
3.3	Survival analysis after a complete remission post alloSCT	19
3.4	Univariate Analysis and multivariate analysis	25

List of Tables

1	Patients characteristics	5
2	Treatments before alloSCT	5
3	Transplant conditions	8
4	Post-AlloSCT Response	10
5	characteristics according to conditionning	11
6	Univariate analysis of 5 years OS survival	26
7	Multivariate analysis of 5 years OS (stratified on the delay between diagnosis and alloSCT)	26
8	Univariate analysis of 5 years GRFS	28
9	Multivariate analysis of 5 years GRFS	28
10	Univariate analysis of 5 years EFS	30
11	Multivariate analysis of 5 years EFS	30
12	Univariate analysis of 5 years cause specific mortality : HSCT related . .	32
13	Multivariate analysis of 5 years cause specific mortality : HSCT related .	32

List of Figures

1	Overall survival	12
2	CIF of relapse or progression and death without relapse or progression . .	13
3	CIF of relapse or progression and death without relapse or progression according to conditionning	14
4	EFS and GRFS	15
5	EFS according to conditionning	16
6	TRM	17
8	CIF of GVHD and Death without GVHD (acute or chronic)	18
9	OS in patients with a complete remission post alloSCT	19
10	CIF of relapse and death without relapse (in patients with a complete remission post alloSCT)	20
11	RFS and EFS in patients with a complete remission post alloSCT	21
12	CIF of Related HSCT Death and Non-related HSCT Death (in patients with a complete remission post alloSCT)	22
13	CIF of GVHD and Death without GVHD (acute or chronic)(in patients with a complete remission post alloSCT)	23
7	CIF of Related HSCT Death and Non-related HSCT Death	24

1 Objectives

The primary objective of the study was to assess the survival, the risk of relapse and GVHD of patients who underwent allogeneic stem-cell transplantation (alloSCT) for aggressive T-cell lymphomas. The second objective was to determine the variables associated with these outcomes.

2 Methods

A retrospective analysis was conducted. A descriptive analysis of the variables recorded was performed. Different endpoints were defined : death, Event Free Survival (EFC), GRFS. GRFS was defined as death, progression/relapse, grade 3-4 acute GVHD or extensive chronic GVHD.

Survival curves were estimated using Kaplan-Meier product-limit estimator. Competing risk survival analysis methods were applied to estimate the cumulative incidence (CIF) of developing events over time from alloSCT. These methods allow for the fact that a patient may experience an event which is different from that of interest. These events are known as competing risk events, and may preclude the onset of the event of interest, or may modify the probability of the onset of that event. In particular, a transplanted patient may die before a relapse occurs.

Factors associated with overall survival were analyzed using Cox proportional hazards models. The proportional hazards assumption was checked by examination of Schoenfeld residuals. Occurrence of a grade 3-4 acute GVHD or chronic GVHD was treated as a time dependent covariable. For the different endpoints, univariable analyses were first carried out, then a multivariable analysis was used where all factors with P-value < 0.05 in the univariable analyses were considered. If needed, factors were then sequentially removed from the adjusted model based on the AIC criteria.

3 Results

3.1 Descriptive results

285 patients were initially selected. We excluded 1 patient that underwent two alloSCT. The final analysis was performed on 284 patients and 284 grafts.

3.1.1 Patients characteristics

Parameters	Values	N	Statistics*
		284	
Patient sex	Female	93	32.75 %
	Male	191	67.25 %
Age at diagnosis		284	45.01(15;68)
Stage at diagnosis	I	13	6.47 %
	II	17	8.46 %
	III	45	22.39 %
	IV	126	62.69 %
	NA	83	
Subtypes	AITL	82	28.87 %
	ALCL ALK-	20	7.04 %
	ALCL ALK?	2	0.7 %
	ALCL ALK+	21	7.39 %
	ATLL	16	5.63 %
	EATL	3	1.06 %
	HS	12	4.23 %
	LGL	1	0.35 %
	NK leukemia	1	0.35 %
	NK/T nasal	16	5.63 %
	NOS	110	38.73 %
	NOS	110	38.73 %
	AITL	82	28.87 %
Subtypes	ALCL	43	15.14 %
	ATLL	16	5.63 %
	NK/T nasal	16	5.63 %
	Others	17	5.99 %
	angers	8	2.82 %
	Becquerel[941]	4	1.41 %
	C.H.R.U Brest[659]	2	0.7 %
	caen	4	1.41 %
	CHU clermont ferrand	7	2.46 %
Centres	Geneve	6	2.11 %
	Gustave Roussy[666]	3	1.06 %
	H A Michallon[270]	5	1.76 %

H Bretonneau[272]	3	1.06 %
H Charles Nicolle[932]	1	0.35 %
H Claude Huriez[277]	8	2.82 %
H de l'ARCHET I[523]nice	3	1.06 %
H E Herriot[671]	5	1.76 %
H Haut-Leveque[267]	31	10.92 %
H Hautepierre[672]	11	3.87 %
H Jean Minjoz[233]	5	1.76 %
H La Miletrie[264]	5	1.76 %
H Mondor Hematol[252]	4	1.41 %
H Necker[160]	9	3.17 %
H Percy[665]	4	1.41 %
H Purpan[624]	8	2.82 %
H Sud/Pontchaillou[661]	7	2.46 %
H Sud[955]	1	0.35 %
Hotel Dieu[253]	32	11.27 %
liege	8	2.82 %
limoges	3	1.06 %
montpellier	10	3.52 %
nancy	1	0.35 %
Paoli Calmettes[230]	39	13.73 %
Pellegrin-Enfants[978]	1	0.35 %
Pitie-Salpetriere[262]	8	2.82 %
St Antoine[775]	10	3.52 %
St Etienne[250]	4	1.41 %
St Louis[207]	24	8.45 %

Table 1: Patients characteristics

3.1.2 Treatments before alloSCT

Parameters	Values	N	Statistics*
		284	
Previous auto	No	191	67.25 %
	Yes	93	32.75 %
Programme auto allo	No	257	90.49 %
	Yes	27	9.51 %
First graft relapse	No	219	77.11 %
	Yes	65	22.89 %

Table 2: Treatments before alloSCT

3.1.3 Transplant conditions

Parameters	Parameters	N	n(%) med[Q1;Q3](min,m
		284	
Age at graft		284	49.5 [38;57] (16;69)
Donor age		263	28 [18;39] (1;54)
Donor sex	Female	114	40.71 %
	Male	166	59.29 %
	NA	4	
Delay diagnosis and allo SCT		284	378.5 [213.2;710.8] (89;9
>12 months delay	NO	149	52.46 %
	Yes	135	47.54 %
Disease status at transplant	CR	175	61.84 %
	PR	76	26.86 %
	PD	32	11.31 %
	NA	1	
Disease status at transplant	CR (?)	7	2.47 %
	CR1	94	33.22 %
	CR2	61	21.55 %
	CR3	13	4.59 %
	PD	32	11.31 %
	PR (?)	13	4.59 %
	PR1	39	13.78 %
	PR2	18	6.36 %
	PR3	5	1.77 %
	PR4	1	0.35 %
	NA	1	
Karnofsky score		263	90 [80;100]
Karnofsky score	100	92	34.98 %
	40	1	0.38 %
	50	4	1.52 %
	60	1	0.38 %
	70	9	3.42 %
	80	70	26.62 %
	90	86	32.7 %
	NA	21	
Karnofsky score	100	92	34.98 %
	Unable to carry on normal activity	15	5.7 %
	90-80	156	59.32 %
	NA	21	
No of lines before alloSCT		254	2 [1;3] (1;9)
No of lines before alloSCT	1	73	28.74 %
	2	92	36.22 %

	3	65	25.59 %
	>=4	24	9.45 %
	NA	30	
No of lines before alloSCT	>2	89	35.04 %
	1 or 2	165	64.96 %
	NA	30	
HLA match	HLA mismatched	53	18.66 %
	HLA matched	231	81.34 %
HLA match	Alternative donnors	53	18.66 %
	Identical sibling	128	45.07 %
	Matched unrelated	103	36.27 %
HLA match	Identical sibling	128	45.07 %
	Matched unrelated	103	36.27 %
	Mismatched relative	7	2.46 %
	Mismatched unrelated	13	4.58 %
	Unrelated CB	33	11.62 %
Sex of donnor/patient	Others	205	73.48 %
	F/M	74	26.52 %
	NA	5	
CMV serostatus of donnor/patient	neg/neg	91	32.5 %
	Others	189	67.5 %
	NA	4	
Source of stem cells	BM	49	17.25 %
	CB	33	11.62 %
	PB	202	71.13 %
TBI	No	161	56.69 %
	Yes	123	43.31 %
conditioning Intensity	MAC	106	38.13 %
	NMA	27	9.71 %
	RIC	145	52.16 %
	NA	6	
Conditioning	BEAM	1	0.36 %
	BEAM + Campath	1	0.36 %
	BU CY	4	1.42 %
	BU CY + FLU+ ATG	1	0.36 %
	BU CY ATG	1	0.36 %
	EDX ATG	0	0 %
	ENX TBI 2gray	1	0.36 %
	FLU ATG	3	1.07 %
	FLU BU 1+ ATG	3	1.07 %
	FLU BU 2	1	0.36 %
	FLU BU 2+ ATG	73	25.98 %
	FLU BU 3+ ATG	21	7.47 %

	FLU BU 4+ ATG	10	3.56 %
	FLU BU EDX	8	2.85 %
	FLU BU EDX +ATG	6	2.14 %
	FLU EDX	1	0.36 %
	FLU EDX ATG	3	1.07 %
	FLU EDX MEL	1	0.36 %
	FLU ENX TBI 2gray	24	8.54 %
	FLU ENX TBI 4gray	2	0.71 %
	FLU ENX TBI 6gray	1	0.36 %
	FLU ENX TBI 6gray + campath	1	0.36 %
	FLU MEL	12	4.27 %
	FLU MEL + campath	4	1.42 %
	FLU MEL + Campath	1	0.36 %
	FLU MEL ATG	1	0.36 %
	FLU MEL TBI 2gray	1	0.36 %
	FLU TBI 2gray	21	7.47 %
	FLU TBI 2gray ATG	1	0.36 %
	FLU Tbi 8 gray	1	0.36 %
	MEL 140 TBI 10 gray	1	0.36 %
	MEL TBI VP16	1	0.36 %
	TB2F	2	0.71 %
	TBI 12 gray	1	0.36 %
	TBI 2gray	1	0.36 %
	TBI EDX	49	17.44 %
	TBI EDX +ATG	11	3.91 %
	TBI EDX FLU	5	1.78 %
	Thiotepa etoposide TBI12 gray	1	0.36 %
	NA	3	
Cells manipulation	No	275	97.86 %
	Yes	6	2.14 %
	NA	3	
Depletion	No	275	98.57 %
	Partial T depletion	4	1.43 %
	NA	5	
No of donnors	1	261	91.9 %
	2	23	8.1 %

Table 3: Transplant conditions

3.1.4 Post-AlloSCT Response

Parameters	Values	N	Statistics*
		284	
Agvhd	No	141	49.65 %
	Yes	143	50.35 %
Agvhd grade	No aGvHD present (Grade 0)	141	49.65 %
	Grade I	49	17.25 %
	Grade II	46	16.2 %
	Grade III	24	8.45 %
	Grade IV	17	5.99 %
	Present, grade unknown	7	2.46 %
Cgvhd	Early death	41	14.44 %
	no	146	51.41 %
	yes	97	34.15 %
Cgvhd grade	Early death (100D)	41	14.44 %
	Extensive	38	13.38 %
	Limited	55	19.37 %
	No cGvh	146	51.41 %
	grade unknown	4	1.41 %
Engrafted	Early death (30D)	5	1.76 %
	Engrafted	271	95.42 %
	Lost graft	2	0.7 %
	No engraftment	6	2.11 %
Cause of death	HSCT-GVHD	21	19.63 %
	HSCT-GVHD + infection	3	2.8 %
	HSCT-infection	27	25.23 %
	HSCT-toxicity	4	3.74 %
	HSCT related	3	2.8 %
	HSCT related ILD	1	0.93 %
	HSCT related MAT	1	0.93 %
	HSCT related MOF	2	1.87 %
	HSCT related MVO	1	0.93 %
	HSCT related pneumopathie interstitielle	2	1.87 %
	HSCT related PTLN	1	0.93 %
	HSCT related SDRA	1	0.93 %
	Other	1	0.93 %
	Relapse or progression of original disease	37	34.58 %
	Secondary malignancy	1	0.93 %
	Unknown	1	0.93 %
	NA	177	
Best reponse after SCT	CR	245	86.88 %
	Not evaluable	4	1.42 %

Relapse/progression	Not evaluated	3	1.06 %
	PD	14	4.96 %
	PR	16	5.67 %
	NA	2	
	Continuous progression	28	9.93 %
	No	217	76.95 %
	Non applicable	3	1.06 %
	Yes	34	12.06 %
	NA	2	

Table 4: Post-AlloSCT Response

Parameters	Values	Conditionnement		NA		
		N	Statistics*	N	Statistics*	p-value
		133	MAC	151	RIC	
Age greffe		133	44 [35;54]	151	53 [43;59]	<0.0001
Nbr de lignes		121	2 [1;3]	133	2 [2;3]	0.045
Nbr de lignes	>2	37	30.58 %	52	39.1 %	0.20
	1 or 2	84	69.42 %	81	60.9 %	
	NA	12		18		
Nbr de lignes	1	41	33.88 %	32	24.06 %	0.14
	2	43	35.54 %	49	36.84 %	
	3	30	24.79 %	35	26.32 %	
	>=4	7	5.79 %	17	12.78 %	
	NA	12		18		
Autogreffe avant	0	101	75.94 %	90	59.6 %	0.005
	1	32	24.06 %	61	40.4 %	
Statut de la maladie	CR	78	58.65 %	97	64.67 %	0.048
	PR	44	33.08 %	32	21.33 %	
	PD	11	8.27 %	21	14 %	
	NA	0		1		
Karnofsky	100	47	37.01 %	45	33.09 %	0.45
	Unable to carry on normal activity	9	7.09 %	6	4.41 %	
	90-80	71	55.91 %	85	62.5 %	
	NA	6		15		
Stade dia	I	5	5.15 %	8	7.69 %	0.035
	II	4	4.12 %	13	12.5 %	
	III	18	18.56 %	27	25.96 %	
	IV	70	72.16 %	56	53.85 %	
	NA	36		47		

Table 5: characteristics according to conditioning

3.2 Survival analysis in all patients

3.2.1 Overall survival, EFS and GRFS

Median follow-up was 20.18 (range 0.03 to 112.83). OS at 1 year was 0.68 (95 % 0.62 - 0.73), was 0.64 (95 % 0.58 - 0.7) at 2 years. OS at 4 years was 0.57 (95 % 0.5 - 0.63).




Figure 1: Overall survival

CIF for relapse/progression at 1 years was 0.18 (95 % 0.13 - 0.23), at 2 years 0.19 (95 % 0.15 - 0.24). CIF for death without relapse or progression at 1 year was 0.19 (95 % 0.14 - 0.24), at 2 years 0.22 (95 % 0.17 - 0.27).




Figure 2: CIF of relapse or progression and death without relapse or progression




Figure 3: CIF of relapse or progression and death without relapse or progression according to conditionning

EFS at 1 year was 0.63 (95 % 0.57 - 0.69), was 0.59 (95 % 0.53 - 0.65) at 2 years. EFS at 4 years was 0.52 (95 % 0.45 - 0.59).

Délai médian rechute : 94 jours.

GRFS at 1 year was 0.49 (95 % 0.44 - 0.56), was 0.47 (95 % 0.41 - 0.54) at 2 years.

GRFS at 4 years was 0.43 (95 % 0.37 - 0.5).




Figure 4: EFS and GRFS




Figure 5: EFS according to conditionning

3.2.2 TRM and cause of death

TRM at 1 year was 0.22 (95 % 0.27 - 0.17), was 0.24 (95 % 0.3 - 0.19) at 2 years. TRM at 4 years was 0.3 (95 % 0.37 - 0.24).




Figure 6: TRM

CIF for related HSCT death at 1 years was 0.2, at 2 years 0.23. CIF for non-related HSCT Death at 1 year was 0.12, at 2 years 0.13.




Figure 8: CIF of GVHD and Death without GVHD (acute or chronic)

3.3 Survival analysis after a complete remission post alloSCT

245 patients with a complete remission were included.

OS at 1 year was 0.74 (95 % 0.68 - 0.8), was 0.7 (95 % 0.64 - 0.76) at 2 years. OS at 4 years was 0.62 (95 % 0.56 - 0.7).




Figure 9: OS in patients with a complete remission post alloSCT

CIF for relapse at 1 year was 0.12 (95 % 0.07 - 0.16), at 2 years 0.13 (95 % 0.09 - 0.18).
 CIF for death without relapse at 1 year was 0.19 (95 % 0.14 - 0.24), at 2 years 0.22 (95 % 0.17 - 0.28).




Figure 10: CIF of relapse and death without relapse (in patients with a complete remission post alloSCT)




Figure 11: RFS and EFS in patients with a complete remission post alloSCT




Figure 12: CIF of Related HSCT Death and Non-related HSCT Death (in patients with a complete remission post alloSCT)




Figure 13: CIF of GVHD and Death without GVHD (acute or chronic)(in patients with a complete remission post alloSCT)




Figure 7: CIF of Related HSCT Death and Non-related HSCT Death

3.4 Univariate Analysis and multivariate analysis

variable	Variable	HR	IC	pval	p
Age at graft		1.01	[0.99 - 1.02]	0.36	0.35
subtypes	NOS	1.00			0.46
	AITL	1.21	[0.75 - 1.97]	0.43	
	ALCL	1.16	[0.64 - 2.09]	0.63	
	ATLL	1.93	[0.93 - 4.01]	0.079	
	NK/T nasal	1.82	[0.85 - 3.93]	0.13	
	Others	1.55	[0.69 - 3.48]	0.29	
Delay between diag and allo SCT > 12 mo	NO	1.00			0.54
	Yes	0.89	[0.6 - 1.3]	0.54	
Stage at diagnosis	I	1.00			0.72
	II	0.49	[0.14 - 1.74]	0.27	
	III	0.79	[0.31 - 1.98]	0.61	
	IV	0.78	[0.33 - 1.81]	0.56	
Disease status at transplant	CR	1.00			0.027
	PR	0.97	[0.61 - 1.53]	0.89	
	PD	2.08	[1.24 - 3.49]	0.006	
Karnofsky score	100	1.00			0.002
	Unable to carry on normal activity	3.05	[1.35 - 6.89]	0.007	
	90-80	2.09	[1.28 - 3.41]	0.003	
First graft relapse	No	1.00			0.049
	No previous graft	2.49	[1.08 - 5.73]	0.032	
	Yes	2.13	[0.87 - 5.22]	0.098	
No of lines before alloSCT	>2	1.00			0.084
	1 or 2	0.70	[0.47 - 1.04]	0.080	
HLA match	HLA mismatched	1.00			0.10
	HLA matched	0.68	[0.43 - 1.07]	0.092	

Sex of donnor-patient	Others	1.00			0.027
	F/M	1.61	[1.07 - 2.42]	0.022	
CMV serostatus of donnor patient	neg/neg	1.00			0.78
	Others	0.94	[0.63 - 1.42]	0.78	
Source of stem cells	BM	1.00			0.016
	CB	1.71	[0.91 - 3.21]	0.094	
	PB	0.77	[0.47 - 1.27]	0.31	
Conditionning intensity	MAC	1.00			0.83
	RIC	1.04	[0.71 - 1.53]	0.83	
Depletion	No	1.00			0.87
	Partial T depletion	1.12	[0.28 - 4.55]	0.87	
Agvhd grade 3-4					<0.0001
		2.82	[1.78 - 4.47]	<0.0001	
Cgvhd					0.17
		1.44	[0.86 - 2.41]	0.16	

Table 6: Univariate analysis of 5 years OS survival

V1	Variable	HR (95%CI)	P
Agvhd	Grade 3-4 Agvhd	2.52 (1.52–4.19)	0.0003
Sex of donnor-patient	F/M	1.33 (0.85–2.08)	0.22
Disease status at transplant	PR	0.83 (0.50–1.36)	0.46
	PD	1.58 (0.88–2.84)	0.13
Karnofsky score	Unable to carry on normal activity	3.16 (1.32–7.61)	0.010
	90-80	2.22 (1.32–3.71)	0.002
Stem cell source	CB	2.01 (1.00–4.01)	0.049
	PB	0.97 (0.56–1.65)	0.90

Table 7: Multivariate analysis of 5 years OS (stratified on the delay between diagnosis and alloSCT)

variable	Variable	HR	IC	pval	p
Age at graft		1.00	[0.99 - 1.01]	0.87	0.86
subtypes	NOS	1.00			0.23
	AITL	1.10	[0.73 - 1.68]	0.64	
	ALCL	0.91	[0.54 - 1.53]	0.73	
	ATLL	2.08	[1.13 - 3.84]	0.019	
	NK/T nasal	1.60	[0.81 - 3.16]	0.18	
	Others	1.22	[0.55 - 2.7]	0.62	
Delay between diag and allo SCT	NO	1.00			0.63
	Yes	0.92	[0.66 - 1.29]	0.63	
Stage at diagnosis	I	1.00			0.94
	II	0.89	[0.31 - 2.57]	0.83	
	III	1.07	[0.44 - 2.61]	0.89	
	IV	1.11	[0.48 - 2.56]	0.81	
Disease status at transplant	CR	1.00			0.057
	PR	1.24	[0.85 - 1.83]	0.27	
	PD	1.90	[1.14 - 3.16]	0.014	
Karnofsky score	100	1.00			0.24
	Unable to carry on normal activity	1.55	[0.61 - 3.93]	0.35	
	90-80	1.36	[0.93 - 1.98]	0.11	
First graft relapse	No	1.00			0.59
	No previous graft	1.30	[0.71 - 2.37]	0.40	
	Yes	1.13	[0.58 - 2.2]	0.72	
No of lines before alloSCT	>2	1.00			0.082
	1 or 2	0.73	[0.51 - 1.04]	0.078	
HLA match	HLA mismatched	1.00			0.077
	HLA matched	0.69	[0.47 - 1.03]	0.067	
Sex of donnor-patient	Others	1.00			0.10
	F/M	1.37	[0.95 - 1.99]	0.093	

CMV serostatus of donor patient	neg/neg	1.00			0.83
	Others	1.04	[0.72 - 1.5]	0.83	
Source of stem cells	BM	1.00			0.036
	CB	1.90	[1.06 - 3.42]	0.031	
	PB	0.99	[0.63 - 1.57]	0.98	
Conditioning intensity	MAC	1.00			0.39
	RIC	1.16	[0.83 - 1.62]	0.39	
Depletion	No	1.00			0.18
	Partial T depletion	2.13	[0.79 - 5.78]	0.14	

Table 8: Univariate analysis of 5 years GRFS

V1	Variable	HR (95%CI)	<i>P</i>
Subtypes	AITL	1.22 (0.79–1.89)	0.37
	ALCL	0.91 (0.53–1.54)	0.72
	ATLL	1.89 (1.01–3.52)	0.046
	NK/T nasal	1.76 (0.89–3.50)	0.10
	Others	1.20 (0.54–2.67)	0.66
Disease status at transplant	PR	1.30 (0.86–1.96)	0.21
	PD	2.02 (1.20–3.41)	0.008
Source of stem cells	CB	2.07 (1.10–3.87)	0.023
	PB	1.04 (0.65–1.66)	0.87

Table 9: Multivariate analysis of 5 years GRFS

variable	Variable	HR	IC	pval	p
Age at graft					0.45
subtypes	NOS	1.01	[0.99 - 1.02]	0.45	0.32
	AITL	1.00			
	ALCL	1.02	[0.65 - 1.62]	0.93	
	ATLL	1.01	[0.57 - 1.77]	0.98	
	NK/T nasal	2.02	[1.04 - 3.94]	0.038	
	Others	1.76	[0.86 - 3.63]	0.12	
Delay between diag and allo SCT > 12 mo	NO	1.01	[0.43 - 2.38]	0.98	0.40
	Yes	1.00			
Stage at diagnosis	I	0.85	[0.59 - 1.23]	0.40	0.84
	II	1.00			
	III	0.63	[0.2 - 1.96]	0.43	
	IV	0.76	[0.31 - 1.9]	0.56	
Disease status at transplant	CR	0.69	[0.3 - 1.61]	0.39	0.088
	PR	1.00			
	PD	1.23	[0.82 - 1.85]	0.32	
Karnofsky score	100	1.92	[1.1 - 3.37]	0.022	0.004
	Unable to carry on normal activity	1.00			
	90-80	1.31	[0.46 - 3.76]	0.62	
First graft relapse	No	2.04	[1.3 - 3.18]	0.002	0.17
	No previous graft	1.00			
	Yes	1.75	[0.88 - 3.48]	0.11	
No of lines before alloSCT	>2	1.37	[0.64 - 2.94]	0.41	0.12
	1 or 2	1.00			
HLA match	HLA mismatched	0.73	[0.5 - 1.08]	0.12	0.058
	HLA matched	1.00			
Sex of donnor-patient	Others	0.65	[0.43 - 1]	0.048	0.44
	F/M	1.00			
		1.18	[0.78 - 1.77]	0.44	

CMV serostatus of donor patient	neg/neg	1.00			0.97
	Others	0.99	[0.67 - 1.47]	0.97	
Source of stem cells	BM	1.00			0.036
	CB	1.96	[1.04 - 3.7]	0.039	
	PB	0.96	[0.58 - 1.59]	0.89	
Conditioning intensity	MAC	1.00			0.18
	RIC	1.28	[0.89 - 1.85]	0.19	
Depletion	No	1.00			0.056
	Partial T depletion	3.16	[1.16 - 8.59]	0.025	
Agvhd grade 3-4					0.004
		2.09	[1.31 - 3.34]	0.002	
Cgvhd					0.10
		1.56	[0.93 - 2.64]	0.094	

Table 10: Univariate analysis of 5 years EFS

V1	Variable	HR (95%CI)	<i>P</i>
N of lines	1 or 2	0.70 (0.46–1.05)	0.086
Karnofsky score	Unable to carry on normal activity	1.52 (0.53–4.39)	0.44
	90-80	2.24 (1.41–3.56)	0.0006
Cell source	CB	1.99 (0.97–4.05)	0.059
	PB	1.00 (0.58–1.73)	0.99

Table 11: Multivariate analysis of 5 years EFS

variable	Variable	HR	IC	pval	p
Age at graft					0.017
subtypes	NOS	1.02	[1 - 1.05]	0.022	0.28
	AITL	1.00			
	ALCL	1.87	[1.03 - 3.38]	0.039	
	ATLL	1.20	[0.54 - 2.64]	0.66	
	NK/T nasal	1.31	[0.39 - 4.43]	0.67	
	Others	1.68	[0.57 - 4.94]	0.35	
Delay between diag and allo SCT > 12 mo	NO	2.42	[0.97 - 6.06]	0.059	0.76
	Yes	1.00			
Stage at diagnosis	I	0.93	[0.57 - 1.5]	0.76	0.47
	II	1.00			
	III	0.48	[0.08 - 2.9]	0.43	
	IV	1.31	[0.38 - 4.54]	0.67	
Disease status at transplant	CR	0.93	[0.28 - 3.04]	0.90	0.33
	PR	1.00			
	PD	0.86	[0.48 - 1.53]	0.60	
Karnofsky score	100	1.59	[0.79 - 3.18]	0.19	0.040
	Unable to carry on normal activity	1.00			
	90-80	2.30	[0.75 - 6.98]	0.14	
First graft relapse	No	2.04	[1.12 - 3.73]	0.020	0.014
	No previous graft	1.00			
	Yes	4.33	[1.05 - 17.9]	0.043	
No of lines before alloSCT	>2	5.57	[1.3 - 23.76]	0.020	0.055
	1 or 2	1.00			
HLA match	HLA mismatched	0.61	[0.37 - 1]	0.052	0.46
	HLA matched	1.00			
Sex of donnor-patient	Others	0.80	[0.44 - 1.43]	0.45	0.026
	F/M	1.00			
		1.80	[1.09 - 2.98]	0.022	

CMV serostatus of donnor patient	neg/neg	1.00		0.99
	Others	1.00	[0.6 - 1.69]	0.99
Source of stem cells	BM	1.00		0.21
	CB	1.29	[0.57 - 2.91]	0.54
	PB	0.71	[0.39 - 1.31]	0.27
Conditionning intensity	MAC	1.00		0.29
	NMA	1.79	[0.83 - 3.85]	0.14
	RIC	1.38	[0.79 - 2.43]	0.26
Depletion	No	1.00		0.92
	Partial T depletion	0.90	[0.12 - 6.49]	0.92

Table 12: Univariate analysis of 5 years cause specific mortality : HSCT related

V1	Variable	HR (95%CI)	<i>P</i>
Previous graft relapse	No previous graft	6.50 (1.50–28.1)	0.012
	Yes	6.58 (1.49–29.0)	0.013
Conditionning intensity	NMA	2.72 (1.17–6.33)	0.021
	RIC	1.45 (0.80–2.63)	0.22
Sex of donnor-patient	F/M	2.13 (1.21–3.75)	0.009
Karnofsky score	Unable to carry on normal activity	2.61 (0.84–8.13)	0.099
	90-80	2.09 (1.11–3.91)	0.022

Table 13: Multivariate analysis of 5 years cause specific mortality : HSCT related