

SARS-CoV-2 vaccination in patients with GI and hepatobiliary carcinoma: a call for booster vaccination

Bolliipo *et al*¹ presented recommendations for SARS-CoV-2 infections in patients with chronic liver diseases (CLDs). Since their publication, at least three SARS-CoV-2 vaccinations are recommended for all persons regardless of comorbidities.² Cancer and CLD are associated with impaired immune responses to SARS-CoV-2 vaccines.^{3–6} However, patients with GI cancer, especially with hepatobiliary carcinoma (HBC), are under-represented in published studies.

In this prospective, longitudinal study, 120 patients with GI cancer, including 32.5% HBC, participated (table 1). Patients under anticancer therapy were analysed compared with patients with GI cancer in follow-up care (≥ 1 year off anticancer therapy). We present profound data on humoral response rates (SARS-CoV-2 antispikes and surrogate neutralisation antibodies (sNAB) using SARS-CoV-2 IgG II Quant chemiluminescent microparticle immunoassay (Abbott Laboratories) and cPass SARS-CoV-2 Neutralization Antibody Detection Kit (GenScript), respectively). Of note, the ELISA analysing levels of sNAB is limited when it comes to current variants of concern (VOCs; BA.1, BA.2, BA.4 and BA.5). Therefore, rates of infections are more important. Cellular response rates were not considered. Linear mixed regression models were used to compare levels of total (\log_{10} transformed) and neutralising antibodies.

Four weeks after second vaccination, levels of SARS-CoV-2 antispikes IgG were significantly lower in patients with active GI cancer (2.48 \log_{10} binding antibody unit (BAU)/mL; 95% CI 2.28 to 2.68; $p < 0.01$) and HBC (2.52 \log_{10} BAU/mL; 95% CI 2.25 to 2.78; $p < 0.01$) compared with patients in follow-up care (3.03 \log_{10} BAU/mL; 95% CI 2.77 to 2.28). Titres decreased overtime, and differences diminished (figure 1). However, patients with HBC still showed significantly lower IgG levels compared with patients in follow-up care 12 weeks after second vaccination (2.06 \log_{10} BAU/mL; 95% CI 1.79 to 2.33 vs 2.47 \log_{10} BAU/mL; 95% CI 2.24 to 2.70; $p = 0.02$). Regarding surrogate neutralisation antibody (sNAB) at the same time point, levels were more impaired in patients

Table 1 Patient baseline characteristics

Patients	GI cancer under treatment (n=52)	HBC* under treatment (n=30)	Off treatment >1 year (n=38)	P value
Age (years)				0.76
Median	66	69	69	
IQR	(61–73)	(62–78)	(61–73)	
Sex				0.33
Women	35.0% (18)	40.0% (12)	50.0% (19)	
Men	65.0% (34)	60.0% (18)	50.0% (19)	
Tumour types				
Cholangiocellular cancer		30.0% (9)	2.5% (1)	0.32
Gallbladder cancer		3.0% (1)	2.5% (1)	
Hepatocellular cancer		67.0% (20)	18.5% (7)	
Colorectal cancer	33.0% (17)		34.5% (13)	0.04
Neuroendocrine tumour	25.0% (13)		2.5% (1)	
Pancreas cancer	21.0% (11)		8.0% (3)	
Gastric cancer	8.0% (4)		16.0% (6)	
Oesophageal cancer	9.0% (5)		13.0% (5)	
GI stromal tumour	4.0% (2)		2.5% (1)	
Intention of treatment				<0.01
Neoadjuvant	9.0% (5)	23.0% (7)		
Adjuvant	23.0% (12)	0.0% (0)		
Palliative	68.0% (35)	77.0% (23)		
Type of treatment				<0.01
Chemotherapy†	42.0% (22)	27.0% (8)		
Targeted therapy	21.0% (11)	10.0% (3)		
Immune checkpoint inhibition	8.0% (4)	0.0% (0)		
Local therapy‡	4.0% (2)	33.0% (10)		
Combined therapy¶	25.0% (13)	30.0% (9)		
History of SARS-CoV-2 infection after booster vaccination	7.7% (3)	3.0% (1)	2.5% (1)	0.85
Vaccines				0.98
BNT162b2 (Pfizer and BioNTech)	82.0% (43)	90.0% (27)	82.0% (31)	
AZD1222 (AstraZeneca)	8.0% (4)	7.0% (2)	8.0% (3)	
mRNA-1273 (Moderna)	4.0% (2)	0.0% (0)	5.0% (2)	
AZD1222+BNT162b2/mRNA-1273	4.0% (2)	3.0% (1)	5.0% (2)	
Ad26.COV2.s (Johnson & Johnson, Janssen)	2.0% (1)	0.0% (0)	0.0% (0)	

Baseline characteristics were compared between the three groups using one-way analysis of variance for age and Fisher's exact tests for the categorical variables.

*Cholangiocellular cancer, gallbladder cancer and hepatocellular cancer.

†Mostly with combinations of 5-fluorouracil, platinum derivatives, gemcitabine, taxane or irinotecan

‡Including multikinase inhibitors such as sorafenib, lenvatinib, cabozantinib, epidermal growth factor receptor antibodies or vessel endothelial growth factor antibodies.

§Including transarterial embolisation, radioembolisation or endobiliary/transcutaneous radiofrequency ablation and photodynamic therapy.

¶Chemotherapy+targeted therapy or chemotherapy+local therapy.

HBC, hepatocellular carcinoma.

with HBC (57.80%; 95% CI 47.49 to 68.11; $p < 0.01$) than in patients with GI-cancer (67.81%; 95% CI 58.33 to 77.29; $p < 0.01$) compared with patients in follow-up care (84.88%; 95% CI 76.34 to 93.43) (figure 1). Patients with HBC under chemotherapy showed most extended impairment of SARS-CoV-2 antispikes IgG levels (1.94 \log_{10} BAU/mL; 95% CI 1.42 to 2.46) 12 weeks after second vaccination in an univariate analysis. While in patients with GI

cancer under chemotherapy levels of SARS-CoV-2 antispikes IgG were also lower (2.16 \log_{10} BAU/mL; 95% CI 1.80 to 2.52), patients under therapy with immune checkpoint inhibitors had highest titres (2.27 \log_{10} BAU/mL; 95% CI 1.43 to 3.11) compared with all other types of therapy.

Following first booster vaccination, that is, the third vaccination, titres of SARS-CoV-2 antispikes IgG were balanced between patients with GI




Figure 1 SARS-CoV-2 antibodies. (A) SARS-CoV-2 antispikes IgG levels. Log₁₀ SARS-CoV-2 antispikes IgG titre of patients with GI cancer*, patients with HCC/CCC and patients in follow-up care (off treatment >1 year) at week 4, 12 and 24 following second vaccination and at week four after third vaccination. Length of box represents the IQR (50% of the data), horizontal line shows the median log₁₀ SARS-CoV-2 antispikes IgG titre. (B) SARS-CoV-2 surrogate neutralisation antibody (sNAB) levels. SARS-CoV-2 sNAB titres of patients with GI cancer*, patients with HCC/CCC and patients in follow-up care (off treatment >1 year) at week 12 following second vaccination and at week 4 after third vaccination. Length of box represents the IQR (50% of the data), horizontal line shows the median SARS-CoV-2 surrogate neutralisation antibody titre. BAU, binding antibody units; CCC, cholangiocarcinoma; HCC, hepatocellular carcinoma. *Colorectal cancer, neuroendocrine tumours, pancreas cancer, gastro-oesophageal-junction cancer, gastric cancer, oesophageal cancer and GI stromal tumours.

cancer (3.28 log₁₀ BAU/mL; 95% CI 3.01 to 3.56; $p=0.29$), HBC (3.36 log₁₀ BAU/mL; 95% CI 2.94 to 3.78; $p=0.60$) and patients with GI cancer in follow-up care (3.50 log₁₀ BAU/mL; 95% CI 3.21 to 3.79). This was also true for sNAB (98.87%; 95% CI 85.95 to 100 or 99.64%; 95% CI 79.9 to 100; $p=0.95$ and 99.40%; 95% CI 87.68 to 100; $p=0.98$, respectively) (figure 1). Mild infections with SARS-CoV-2 were detected in five patients after booster vaccination.

In line with recent data on patients with solid tumours,³ immunogenicity after second vaccination for SARS-CoV-2 in patients with GI cancer under anticancer therapy, particularly under chemotherapy, was significantly reduced. The capacity to develop adequate response rates, above all concerning sNAB, was more pronounced than in previous reports including all kind of solid tumours³ and was most impaired in patients with HBC. Patients with CLD showed reduced immunogenicity for different vaccines including those for SARS-CoV-2.^{4–6} Underlying cirrhosis in most patients with HBC also contributes to impaired immune responses.⁷ Differences between antibody levels were overcome by third vaccination with only rare infections thereafter highlighting the importance of booster vaccinations especially for patients at high-risk for impaired SARS-CoV-2 immunogenicity. As waning immunity overtime can be assumed⁸ and clinical benefits of a fourth

vaccine dose have been proved,⁹ additional booster vaccinations after further antibody assessment in the vulnerable group of patients with GI cancer/ HBC should be recommended. In future studies, long-term SARS-CoV-2 immunogenicity, effects of additional booster vaccinations and new SARS-CoV-2 VOC (like BA.5 at the very moment) showing antibody escape from vaccines¹⁰ have to be considered.

Malte Benedikt Monin^{1,2}, Leona Baier,¹ Moritz Berger,³ Jens Gabriel Gorny,¹ Taotao Zhou,¹ Robert Mahn,¹ Farsaneh Sadeghfar,¹ Christian Möhring,¹ Kathrin van Bremen,^{1,2} Christoph Boesecke,^{1,2} Jürgen Rockstroh,^{1,2} Christian Strassburg^{1,2}, Anna-Maria Eis-Hübinger,⁴ Maria Agnes Gonzalez-Carmona¹

¹Department of Internal Medicine I, University Hospital Bonn, Bonn, Germany

²German Centre for Infection Research (DZIF), Partner-site Cologne-Bonn, Bonn, Germany

³Institute for Medical Biometry, Informatics and Epidemiology, University Hospital Bonn, Bonn, Germany

⁴Institute of Virology, University Hospital Bonn, Bonn, Germany

Correspondence to Dr Malte Benedikt Monin, University Hospital Bonn, 53127 Bonn, Germany; malte_benedikt.monin@ukbonn.de

Acknowledgements The authors extend their grateful thanks to all study participants. Moreover, we would like to thank Tobias Klant, member of the Institute of Virology at the University Hospital Bonn, Germany, for performing the ELISA and immunoassays.

Contributors MBM: planning study, data interpreting, drafting and editing of manuscript, final approval, submitting the manuscript, responsible for the overall content as guarantor; LB: data collecting and final approval; MB: data interpreting, drafting of images,

revision of manuscript and final approval; JGG: data collecting and interpreting, drafting of tables/ images, editing of manuscript and final approval; TZ: data collecting and final approval; RM: data collecting and final approval; FS: planning study, data collecting and final approval; CM, KvB and CS: final approval; CB and JR: revision and final approval; A-ME-H: data collecting, editing and final approval and responsible for the overall content as guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests CB received honoraria for lectures and/ or consultancies from AbbVie, Gilead, Janssen, MSD, ViiV and funding from Dt. Leberstiftung, DZIF, Hector Stiftung, NEAT ID. MAG-C has contributed to advisory boards for Roche, Eisai, BMS, MSD and AZ. MBM received travel expenses and honoraria from Gilead and Virology Education. GJR received travel expenses and honoraria from Gilead. JR has received honoraria for lectures and/ or consultancies from Abivax, Galapagos, Gilead, Merck, Janssen, Theratechnologies and ViiV. However, these activities have no potential conflicts of interest with the manuscript.

Patient consent for publication Not applicable.

Ethics approval The study was approved by the ethics committee of the Medical Faculty of the University of Bonn (No. 341/17 and 023/22). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

© Author(s) (or their employer(s)) 2022. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Monin MB, Baier L, Berger M, et al. *Gut* Epub ahead of print: [please include Day Month Year]. doi:10.1136/gutjnl-2022-328169

Received 29 June 2022

Accepted 20 July 2022

Gut 2022;0:1–3. doi:10.1136/gutjnl-2022-328169

ORCID iDs

Malte Benedikt Monin <http://orcid.org/0000-0002-8794-3612>

Christian Strassburg <http://orcid.org/0000-0001-7870-5359>

REFERENCES

- 1 Bollipo S, Kapuria D, Rabiee A, *et al*. One world, one pandemic, many guidelines: management of liver diseases during COVID-19. *Gut* 2020;69:1369–72.
- 2 Wratil PR, Stern M, Priller A, *et al*. Three exposures to the spike protein of SARS-CoV-2 by either infection or vaccination elicit superior neutralizing immunity to all variants of concern. *Nat Med* 2022;28:496–503.
- 3 Fendler A, de Vries EGE, GeurtsvanKessel CH, *et al*. COVID-19 vaccines in patients with cancer: immunogenicity, efficacy and safety. *Nat Rev Clin Oncol* 2022;19:385–401.
- 4 Cornberg M, Buti M, Eberhardt CS, *et al*. EASL position paper on the use of COVID-19 vaccines in patients with chronic liver diseases, hepatobiliary cancer and liver transplant recipients. *J Hepatol* 2021;74:944–51.
- 5 Thuluvath PJ, Roberts P, Chauhan M. Analysis of antibody responses after COVID-19 vaccination in liver transplant recipients and those with chronic liver diseases. *J Hepatol* 2021;75:1434–9.
- 6 Ruether DF, Schaub GM, Duengelhoefer PM, *et al*. SARS-CoV2-specific humoral and T-cell immune response after second vaccination in liver cirrhosis and transplant patients. *Clin Gastroenterol Hepatol* 2022;20:162–72.
- 7 Albillos A, Lario M, Álvarez-Mon M. Cirrhosis-associated immune dysfunction: distinctive features and clinical relevance. *J Hepatol* 2014;61:1385–96.
- 8 Lee LYW, Starkey T, Ionescu MC, *et al*. Vaccine effectiveness against COVID-19 breakthrough infections in patients with cancer (UKCCEP): a population-based test-negative case-control study. *Lancet Oncol* 2022;23:748–57.
- 9 Brosh-Nissimov T, Hussein K, Wiener-Well Y, *et al*. Hospitalized patients with severe COVID-19 during the Omicron wave in Israel - benefits of a fourth vaccine dose. *Clin Infect Dis* 2022;20:ciac501.
- 10 Tuekprakhon A, Nutalai R, Djokaite-Guraliuc A, *et al*. Antibody escape of SARS-CoV-2 omicron BA.4 and BA.5 from vaccine and BA.1 serum. *Cell* 2022;185:2422–33.