

## Large saddle pulmonary embolism in a woman infected by COVID-19 pneumonia

Ramezan Jafari<sup>1,2</sup>, Luca Cegolon<sup>3,4</sup>, Atefeh Jafari<sup>5</sup>, Mandana Kashaki<sup>6</sup>, Babak Otoukesh<sup>7,8</sup>, Bahareh Heshmat Ghahderijani<sup>9</sup>, Morteza Izadi<sup>2</sup>, Seyed Hassan Saadat<sup>10,11</sup>, Behzad Einollahi<sup>11</sup>, and Mohammad Javanbakht<sup>11\*</sup>

<sup>1</sup>Department of Radiology, Baqiyatallah University of Medical Sciences, Tehran, Iran; <sup>2</sup>Health Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran; <sup>3</sup>Local Health Unit N. 2 'Marca Trevigiana' Public Health Department, Treviso, Italy; <sup>4</sup>Institute for Maternal & Child Health, IRCCS 'Burlo Garofolo', Trieste, Italy; <sup>5</sup>Department of Clinical Pharmacy, School of Pharmacy, Guilan University of Medical Sciences, Rasht, Iran; <sup>6</sup>Shahid Akbarabadi Clinical Research Development Unit (ShACRDU), Iran University of Medical Sciences (IUMS), Tehran, Iran; <sup>7</sup>Bone and Joint Reconstruction Research Center, Iran University of Medical Sciences, Tehran, Iran; <sup>8</sup>Department of Orthopedic and Traumatology, Universitätsklinikum Bonn, Bonn, Germany; <sup>9</sup>MD, Radiologist; <sup>10</sup>Behavioral Science Research Center, Lifestyle Institute, Baqiyatallah University of Medical Sciences, Tehran, Iran; and <sup>11</sup>Nephrology and Urology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

\* Corresponding author: Email: mhmjvbt81@yahoo.com

A 50-year-old woman was admitted to the Accident & Emergency department of Baqiyatallah hospital in Tehran (Iran), with symptoms of fever, dyspnoea, dry cough, and fatigue for the past 4 days.

At physical examination, she had enhanced body temperature (38.1 °C), a heart rate of 100 b.p.m., a respiratory rate of 22 breaths/min, and O<sub>2</sub> saturation of 92%. Laboratory tests revealed an increased white blood cell count (WBC =  $13.5 \times 10^3/\mu\text{L}$ ), increased platelets (PLT =  $475 \times 10^3/\mu\text{L}$ ), polynuclear (80%); partial thromboplastin time (PTT = 5 s), PT<sub>Lab</sub>/INR<sub>Lab</sub> (=1.31 s); prothrombin test (PT = 18.2 s); and D-dimer (10 ng/mL). Further elevated values were found for C-reactive protein (CRP = 30.3 mg/L), erythrocyte sedimentation rate (ESR = 57 mm/h); serum glutamic-pyruvic transaminase (SGPT = 108  $\mu\text{L}$ ); lactate dehydrogenase (LDH = 983  $\mu\text{L}$ ); and amylase serum (278  $\mu\text{L}$ ). Reverse transcription-PCR for COVID-19 was positive and the patient was given oxygen and treated with hydroxychloroquine, dimenhydrinate, and naproxen.

On day 2 after admission, a chest CT scan without contrast evidenced bilateral diffuse multiple patchy and subpleural consolidations with left pleural effusion—radiological findings consistent with COVID-19 pneumonia (Panel A). On day 3 following admission, the patient suddenly developed severe dyspnoea, with O<sub>2</sub> saturation decreasing to 74%. A chest computed tomography angiography (CTA) was performed, which revealed a large saddle pulmonary embolism in the main pulmonary artery (MPA), right pulmonary artery (RPA), and left pulmonary artery (LPA) (Panel B). From day 3 of hospitalization, the patient received anticoagulation therapy with heparin and antithrombotic treatment.

The patient remained in hospital for a further 7 days before being discharged in stable condition, with O<sub>2</sub> saturation increased to 98%, body temperature of 37.2 °C, negative PCR, and normal laboratory tests.



