

COVID-19 vaccination and allergen immunotherapy (AIT)

©2021 Dustri-Verlag Dr. K. Feistle
ISSN 2512-8957

DOI 10.5414/ALX02245E
e-pub: August 27, 2021

A position paper of the German Society for Applied Allergology (AeDA) and the German Society for Allergology and Clinical Immunology (DGAKI)

Ludger Klimek¹, Oliver Pfaar², Eckard Hamelmann³, Jörg Kleine-Tebbe⁴, Christian Taube⁵, Martin Wagenmann⁶, Thomas Werfel⁷, Randolph Brehler⁸, Natalija Novak⁹, Norbert Mülleneisen¹⁰, Sven Becker¹¹, and Margitta Worm¹²

¹Center for Rhinology and Allergology, Wiesbaden, ²Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Marburg, ³University Hospital for Pediatrics and Adolescent Medicine, Children's Center Bethel, University of Bielefeld, ⁴Allergy Center Westend, Berlin, ⁵Department of Pneumology, University Hospital Essen – Ruhrlandklinik, Essen, ⁶Department of Otorhinolaryngology, University Hospital Düsseldorf, Düsseldorf, ⁷Department of Dermatology, Allergology and Venerology, Hanover Medical School, ⁸Department of Dermatology, University Hospital Münster, Division of Allergology, Occupational Dermatology and Environmental Medicine, Münster, ⁹Department of Dermatology and Allergy, Polyclinic for Dermatology and Allergology, Bonn, ¹⁰Asthma and Allergy Center, Leverkusen, ¹¹Clinic for Otorhinolaryngology, University Hospital, Tübingen, and ¹²Allergology and Immunology, Clinic for Dermatotoxicology, Venerology and Allergology, Charité Universitätsmedizin Berlin, Germany

Key words

allergen immunotherapy
– allergic rhinitis –
asthma – COVID-19 –
SARS-CoV-2

German version
published in
Allergologie,
Vol. 44, No. 5/2021,
pp. 339-348

Received
April 1, 2021;
accepted in revised form
June 11, 2021

Correspondence to
Prof. Dr. med.
Ludger Klimek
Zentrum für Rhinologie
und Allergologie,
An den Quellen 10,
65183 Wiesbaden,
Germany
Ludger.Klimek@
Allergiezentrum.org

Abstract. Background: Vaccinations against severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2) are intended to induce an immune response to protect against infection/disease. Allergen immunotherapy (AIT) is thought to induce a (different) immune response, e.g., to induce tolerance to allergens. In this position paper we clarify how to use AIT in temporal relation to COVID-19 vaccination. Four SARS-CoV-2 vaccines are currently approved in the EU, and their possible immunological interactions with AIT are described together with practical recommendations for use. Materials and methods: Based on the internationally published literature, this position paper provides specific recommendations for the use of AIT in temporal relation to a SARS-CoV-2 vaccination. Results: AIT is used in 1) allergic rhinitis, 2) allergic bronchial asthma, 3) insect venom allergy, 4) food allergy (peanut). Conclusion: For the continuation of an ongoing AIT, we recommend an interval of 1 week before and after vaccination for subcutaneous immunotherapy (SCIT). For sublingual immunotherapy (SLIT) and oral immunotherapy (OIT), we recommend taking

them up to the day before vaccination and a break of 2 – 7 days after vaccination. Initiation of a new SCIT, SLIT, or OIT should be delayed until 1 week after the day of the second vaccination. For SCIT, we generally recommend an interval of ~ 1 week to COVID-19 vaccination.

Introduction

Allergen immunotherapy

Allergen immunotherapy (AIT) is the only causally effective therapy for which long-term clinical benefit has been demonstrated in allergic respiratory diseases, for example, allergic bronchial asthma or allergic rhinoconjunctivitis and other allergic diseases [1]. Since its first description more than 100 years ago (1911) [2], AIT has been an established and internationally recognized procedure for the treatment of allergies.

AIT induces immune tolerance to a specific, individually relevant allergen [3]. Sys-

tematic meta-analyses have confirmed that AIT significantly reduces symptoms of allergic disease and the amount of necessary antisymptomatic medication in patients with allergic asthma [4] and allergic rhinoconjunctivitis [5].

This is true for both subcutaneous immunotherapy (SCIT) [6, 7] and sublingual immunotherapy (SLIT) [8].

The risk of patients with allergic rhinitis developing asthma is reduced by AIT [9, 10]. AIT is also effective in patients with IgE-mediated food allergy [11, 12] and insect venom allergy [13]. In addition, the cost-saving effect of this disease-modifying therapeutic option [14, 15, 16] has been demonstrated.

After the World Health Organization (WHO) declared the severe acute respiratory syndrome coronavirus type 2 (SARS-CoV-2)-transmitted corona virus 19 (COVID-19) infectious disease pandemic in March 2020 [17], numerous position papers and recommendations for action from international and national allergological societies on the management of allergological diseases and their therapies in the pandemic have been published [18, 19, 20, 21, 22, 23, 24, 25, 26, 58, 59, 60, 61, 62, 63].

For the two mRNA-based vaccines currently approved in Europe (Comirnaty from BioNTech [27]; mRNA-1273 from Moderna [28]) and two vector-based vaccines (CHAdOx1-S from AstraZeneca [29]; Ad26.COV2-S from Johnson & Johnson [30]), recommendations for allergological risk assessment of COVID-19 vaccinations have been published in cooperation of AeDA (Ärztverband Deutscher Allergologen), DGAKI (Deutscher Gesellschaft für Allergologie und Klinische Immunologie), and GPA (Gesellschaft für Pädiatrische Allergologie und Umweltmedizin) [31, 32, 33].

Based on the technical and directions for use of the four approved COVID-19 vaccines, AIT is not a contraindication [27, 28, 29, 30].

On this basis, scientific societies have published recommendations for COVID-19 vaccinations for patients undergoing biologic therapy [25, 26, 64].

The aim of this position paper is to present the use of AIT in its different forms of application (SCIT, SLIT, oral (OIT)) in the context of COVID-19 vaccination and to

provide detailed recommendations for action (Figure 1).

Immune responses in AIT

AIT attempts to induce a tolerogenic response to an individual antigen dose by continuous administration of that antigen. The main mechanisms include early desensitization of effector cells and progressive onset of a regulatory B and T cell response [34, 35, 36]. Although the main changes of specific immunotherapy are antigen-specific, recent data support a beneficial effect in restoring the general balance of the immune system altered toward Th2 immune response [37, 38]. A recent review analyzes the impact of the COVID-19 pandemic on the management of AIT in routine practice [39].

Because AIT is administered over a period of several years, there is considerable experience in coadministration with antiviral and other antimicrobial vaccines. Here, negative influences have not become known. On the contrary, immunologically, a restored balance of the innate immune system could improve its protective function [40].

General information on COVID-19 vaccines

COVID-19 has caused circa 110 million illnesses worldwide and claimed 2.5 million lives as of the writing of this review [41].

Effective vaccination against the novel virus represents an essential strategy to achieve maintenance of health care and public life while reducing social constraints [42].

Four vaccination goals exist in the context of COVID-19 vaccination. Most important is the prevention of severe COVID-19 courses and deaths. In addition, the protection of individuals at particularly high occupational risk of infection, prevention of disease transmission, and maintenance of public life are also important [43].

Due to extensive collaborations between academia and industry, new vaccine platforms were developed at an unprecedented pace less than a year after the discovery of the SARS-CoV-2 viral sequence, extensively




Figure 1. Recommendations for subcutaneous immunotherapy (SCIT) and sublingual immunotherapy SLIT. During the phases marked in green, vaccination can be carried out; during the phases marked in blue, there should be no vaccination.

and comprehensively tested in clinical trial programs, and approved (some provisionally or conditionally) after thorough review by regulatory authorities.

COVID-19 vaccines approved in Germany

Worldwide, 66 COVID-19 vaccines are currently in clinical development and another 176 are in a preclinical development phase [44].

Vaccines against viral infectious diseases are designed to induce humoral and cellular immune responses against the vaccinated antigen. For this purpose, classical technologies are used, but also vaccine variants that have never been approved for human pharmacology [44].

The most common route of vaccine application is intramuscular injection, which induces strong priming of dendritic cells. However, vaccines using other routes of application are also currently being developed and their pharmacological properties and ef-

fects in terms of immune induction remain to be seen [45].

Traditionally, either complete viruses are used, or pathoimmunologically significant parts of the virus (for example, those necessary for entry into somatic cells).

The mRNA-based SARS-CoV-2 vaccines from BioNTech/Pfizer and Moderna break new ground in terms of vaccine antigen delivery. BNT162b2 and mRNA-1273 are mRNA-based vaccines that do not introduce the antigen against which an immune response is to be induced (surface protein of the SARS-CoV-2 virus) but the blueprint (the mRNA) to produce the target protein in human cells. Primarily, the mRNA is taken up in muscle cells, but dendritic cells are critically involved in the presentation of the antigen. Regional lymph nodes are then where the immune response predominantly occurs. The target cells produce the viral antigen based on the information of the mRNA by transcription into the amino acid sequence of the protein structure, which becomes visible to the immune system of the vaccinee as a surface protein of such “transduced” cells. The immune system recognizes

the surface protein of the SARS-CoV-2 as foreign and starts a humoral and cellular immune response. Degradation products of the produced proteins (peptides) are presented to the T cells, which are thus activated. This enables them to recognize and kill virus-infected cells. Furthermore, activated T cells support the antibody-producing B cells.

Due to the rapid degradation of mRNA, it must be packaged in liposomes as a protective envelope so that uptake into the body's own cells (the transfection) works. Their production has been improved in recent years, but liposomes and the mRNA they contain break down very easily, so these vaccines must be stored at very low temperatures (-20°C (mRNA-1273) to -70°C (BNT162b2) between production and use in humans. The lipid nanoparticles and their components also appear to be responsible for the severe allergic reactions to mRNA vaccines [32, 33].

Vector vaccines are another new class of vaccines. The AstraZeneca vaccine (ChAdOx1-S-(AZD1222)), the vaccine Ad26COV2.S from Janssen Pharmaceutical (Johnson & Johnson), and Sputnik V (Gam-COVID-Vac Adeno-based (rAd26-S+rAdS-S)) from the Moscow Gamaleja Institute are based on harmless human or monkey viruses that are unable to self-replicate and contain the SARS-CoV-2 surface protein [46].

The adenovirus-based vector vaccines can be stored in the refrigerator at 4°C for several months without losing their efficacy.

A disadvantage of virus vector vaccines in general is that they cannot be used to vaccinate several times in succession, because the vaccinee develops neutralizing antibodies against the vector itself. As a result, booster vaccinations can be only reduced or no longer effective. This is not a problem with Janssen Pharmaceutical's Ad26COV2.S vaccine, which only needs to be administered once. When booster vaccination is necessary, an alternative strategy in cases of initial immunization with vector virus type A is booster using a heterologous vector type B vaccine. This has already become a reality in the randomized placebo-controlled trial with the Sputnik V vaccine developed in Russia recently [46].

Immunology of type 1 allergies

The type 1 allergic reaction is based on a T2 immune response in which immunoglobulin E (IgE) and the cytokines interleukin (IL)-4, IL-5, and IL-13 in particular are significantly involved. From the use of biologics, we know that these key elements of T2 inflammation do not play a role in the antiviral immune response with respect to either IgE or the aforementioned cytokines. The immune response to anti-viral vaccination was not attenuated [47, 48], whereas anti-IgE treatment increased the production of type 1 interferon from dendritic cells, thereby enhancing the antiviral response [26, 49, 50].

Practical recommendations for allergen immunotherapy

SCIT/SLIT/OIT in temporal connection with COVID-19 vaccinations

In principle, manufacturer-specific guidelines Summary of Product Characteristics must be taken into account in the temporal connection of AIT and vaccinations. Thus, there should generally be an interval of ~ 1 week between SCIT and COVID-19 vaccination. Based on experience with other vaccinations, the following procedure has proven effective [25, 40, 51].

Induction phase

If it is possible to carry out the boosting phase of AIT completely before the planned vaccination date, this can be done as usual, and the recommendations given under "Maintenance therapy" then apply. If vaccination is imminent, initiation of SCIT, SLIT, or OIT should be delayed until 1 week after the second vaccination date [25].

Maintenance therapy

For continuation of ongoing AIT, we recommend a period of ~ 1 week between SCIT and vaccination, analogous to the above procedure, as well as at least 1 week interval

after vaccination, observing the minimum interval between 2 SCIT applications recommended by the manufacturer.

For SLIT or OIT, there are different recommendations from different manufacturers on the interval between vaccination and previous and subsequent SLIT or OIT administration. Therefore, no general recommendation can be given, but the information in the Summary of Product Characteristics should be considered and an individual decision should be made thereafter. In order to be able to limit possible side effects of the SLIT or OIT or the vaccination, we recommend to pause the SLIT or OIT on the day of the vaccination and to continue it with a certain time lag (2 – 7 days). In this case, SLIT or OIT can be applied up to the day before the vaccination [25].

Allergic reactions to vaccines are very rare, occurring at 1 per 1,000,000 – 30 per 100,000 vaccinations [52, 53, 54, 55, 56, 57].

To date, there is no evidence that they are increased under AIT [25, 40].

Discussion and summary

Currently, there is no scientifically substantiated evidence for clinically relevant interactions between AIT and the COVID-19 vaccines currently available in Germany.

Both mRNA vaccines against SARS-CoV-2 are based on the same lipid-based nanoparticle carrier technology; the other two vaccines are vector vaccines.

Additional vaccines are expected to be licensed in the coming months, and it is almost inevitable that adverse drug reactions will occur in the coming months that were not observed in the studies conducted for marketing authorization. Such real-life data will play a significant role in assessing interactions with other drugs – including AIT.

Vaccine safety requires a proactive approach to maintain public confidence and reduce reluctance to vaccination among segments of the population. Vigilance, careful response, documentation, and characterization of these events are necessary to allow definition of mechanisms and appropriate approaches for prediction, prevention, and treatment. This is especially true for poten-

tial interactions of allergy therapies such as AIT.

Funding

This work has been funded by German Allergy Societies AeDA and DGAKI.

Conflict of interest

L. Klimek reports research funding, grants, and/or honoraria in the last 3 years from Allergopharma, Bioprojet, Biomay, Circassia, Viatris, HAL Allergie, ALK Abelló, Aimmune, Immunotek S.L., LETI Pharma, Stallergenes, Quintiles, Sanofi, ASIT Biotech, Lofarma, Thermofisher, Roxall, Allergy Therapeutics, AstraZeneca, GSK, Immunotek, AeDA, Pohl Boskamp GmbH, Paul Martini Foundation, outside the submitted work; and is a member of the following organizations: AeDA, DGHNO, German Academy of Allergology and Clinical Immunology, German Allergy League; ENT-BV, GPA, EAACI.

O. Pfaar reports grants and/or honoraria from ALK-Abelló, Allergopharma, Stallergenes Greer, HAL Allergy Holding B. V./HAL Allergie GmbH, Bencard Allergie GmbH/Allergy Therapeutics, Immunotek S.L., Lofarma, Biomay, Circassia, ASIT Biotech Tools S. A., Laboratorios LETI/LETI Pharma, MEDA Pharma/MYLAN, Anergis S. A., Mobile Chamber Experts (a GA2LEN Partner), Indoor Biotechnologies, GlaxoSmith-Kline, Astellas Pharma Global, EUFOREA, ROXALL Medizin, Novartis, Sanofi Aventis, Sanofi Genzyme, Med Update Europe GmbH, streamedup! GmbH, Pohl-Boskamp GmbH, John Wiley and Sons AS, Paul Martini Foundation (PMS), Regeneron Pharmaceuticals Inc., Ingress-Health HWM during the last 36 months and all outside the present work.

R. Brehler reports honoraria from ALK Abelló, Allergopharma, Allmiral, AstraZeneca, Bencard, Gesellschaft zur Förderung der Dermatologischen Forschung und Fortbildung e.V., GSK, HAL Allergie, LETI Pharma, MedUpdate, Merck, Novartis, Sanofi, Stallergenes, outside the submitted work; and membership in the following organizations: AeDA, DGAKI, EAACI, ABD.

M. Wagenmann reports honoraria from ALK-Abello, Allergopharma, AstraZeneca, Bencard, Genzyme, GlaxoSmithKline, HAL Allergy, LETI, Meda Pharma, Novartis, Sanofi Aventis, Stallergenes, all outside the present work.

M. Worm reports other/other conflicts of interest by Regeneron Pharmaceuticals, DBV Technologies S.A, Stallergenes GmbH, HAL Allergy GmbH, Bencard Allergie GmbH, Allergopharma GmbH & Co. KG, ALK-Abelló Arzneimittel GmbH, Mylan Germany GmbH, Leo Pharma GmbH, Sanofi-Aventis Deutschland GmbH, Aimmune Therapeutics UK Limited, Actelion Pharmaceuticals Deutschland GmbH, Novartis AG, Biotest AG, AbbVie Deutschland GmbH & Co. KG, Lilly Deutschland GmbH, all outside the scope of the present work.

S. Becker reports honoraria from HAL Allergie GmbH, Bencard Allergie GmbH, Allergopharma GmbH & Co. KG, ALK-Abelló Arzneimittel GmbH, Mylan Germany GmbH, Sanofi-Aventis Deutschland GmbH, Novartis AG, AstraZeneca, Ambu, Karl Storz, outside the submitted work.

N. Novak reports honoraria from Alk Abello, Stallergens Geer, Hal Allergy, Leti Pharma, Sanofi Genzyme, Abbvie, Leo Pharma, Novartis, streamed up, and Blueprint outside the submitted work.

The other authors declare no conflicts of interest.

References

- [1] Bousquet J, Lockey R, Malling HJ. Allergen immunotherapy: therapeutic vaccines for allergic diseases. A WHO position paper. *J Allergy Clin Immunol.* 1998; 102: 558-562. [CrossRef PubMed](#)
- [2] Durham SR, Leung DY. One hundred years of allergen immunotherapy: time to ring the changes. *J Allergy Clin Immunol.* 2011; 127: 3-7. [CrossRef PubMed](#)
- [3] Jutel M, Van de Veen W, Agache I, Azkur KA, Akdis M, Akdis CA. Mechanisms of allergen-specific immunotherapy and novel ways for vaccine development. *Allergol Int.* 2013; 62: 425-433. [CrossRef PubMed](#)
- [4] Dhami S, Kakourou A, Asamoah F, Agache I, Lau S, Jutel M, Muraro A, Roberts G, Akdis CA, Bonini M, Cavkaytar O, Flood B, Gajdanowicz P, Izuhara K, Kalayci Ö, Mosges R, Palomares O, Pfaar O, Smolinska S, Sokolowska M, et al. Allergen immunotherapy for allergic asthma: A systematic review and meta-analysis. *Allergy.* 2017; 72: 1825-1848. [CrossRef PubMed](#)
- [5] Dhami S, Nurmatov U, Arasi S, Khan T, Asaria M, Zaman H, Agarwal A, Netuveli G, Roberts G, Pfaar O, Muraro A, Ansotegui IJ, Calderon M, Cingi C, Durham S, van Wijk RG, Halken S, Hamelmann E, Hellings P, Jacobsen L, et al. Allergen immunotherapy for allergic rhinoconjunctivitis: A systematic review and meta-analysis. *Allergy.* 2017; 72: 1597-1631. [CrossRef PubMed](#)
- [6] Klimek L, Brehler R, Hamelmann E, Kopp M, Ring J, Treudler R, Jakob T, Worm M, Pfaar O. Development of subcutaneous allergen immunotherapy (part 2): preventive aspects and innovations. *Allergo J Int.* 2019; 28: 107-119. [CrossRef](#)
- [7] Klimek L, Brehler R, Hamelmann E, Kopp M, Ring J, Treudler R, Jakob T, Worm M, Pfaar O. Evolution of subcutaneous allergen immunotherapy (part 1): from first developments to mechanism-driven therapy concepts. *Allergo J Int.* 2019; 28: 78-95. [CrossRef](#)
- [8] Durham SR, Emminger W, Kapp A, de Monchy JG, Rak S, Scadding GK, Wurtzen PA, Andersen JS, Tholstrup B, Riis B, Dahl R. SQ-standardized sublingual grass immunotherapy: confirmation of disease modification 2 years after 3 years of treatment in a randomized trial. *J Allergy Clin Immunol.* 2012; 129: 717-725.e5. [CrossRef PubMed](#)
- [9] Nurmatov U, Dhami S, Arasi S, Roberts G, Pfaar O, Muraro A, Ansotegui IJ, Calderon M, Cingi C, Durham S, van Wijk RG, Halken S, Hamelmann E, Hellings P, Jacobsen L, Knol E, Larenas-Linnemann D, Lin SY, Maggina V, Oude-Elberink H, et al. Allergen immunotherapy for allergic rhinoconjunctivitis: a systematic overview of systematic reviews. *Clin Transl Allergy.* 2017; 7: 24. [CrossRef PubMed](#)
- [10] Queisser A, Hagedorn S, Wang H, Schaefer T, Konantz M, Alavi S, Deng M, Vogel W, von Müssenhäusen A, Kristiansen G, Duensing S, Kirfel J, Lengerke C, Perner S. Ecotropic viral integration site 1, a novel oncogene in prostate cancer. *Oncogene.* 2017; 36: 1573-1584. [CrossRef PubMed](#)
- [11] Blumchen K, Trendelenburg V, Ahrens F, Gruebl A, Hamelmann E, Hansen G, Heinzmann A, Nemat K, Holzhauser T, Roeder M, Rosenfeld L, Hartmann O, Niggemann B, Beyer K. Efficacy, safety, and quality of life in a multicenter, randomized, placebo-controlled trial of low-dose peanut oral immunotherapy in children with peanut allergy. *J Allergy Clin Immunol Pract.* 2019; 7: 479-491.e10. [CrossRef PubMed](#)
- [12] Chu DK, Wood RA, French S, Fiocchi A, Jordana M, Waserman S, Brożek JL, Schünemann HJ. Oral immunotherapy for peanut allergy (PACE): a systematic review and meta-analysis of efficacy and safety. *Lancet.* 2019; 393: 2222-2232. [CrossRef PubMed](#)
- [13] Dhami S, Zaman H, Varga EM, Sturm GJ, Muraro A, Akdis CA, Antolin-Amérigo D, Bilò MB, Bokanovic D, Calderon MA, Cichocka-Jarosz E, Oude Elberink JN, Gawlik R, Jakob T, Kosnik M, Lange J, Mingomataj E, Mitsias DI, Mosbech H, Ollert M, et al. Allergen immunotherapy for insect venom allergy: a systematic review and meta-analysis. *Allergy.* 2017; 72: 342-365. [CrossRef PubMed](#)
- [14] Asaria M, Dhami S, van Ree R, Gerth van Wijk R, Muraro A, Roberts G, Sheikh A. Health economic analysis of allergen immunotherapy for the management of allergic rhinitis, asthma, food allergy

- and venom allergy: A systematic overview. *Allergy*. 2018; 73: 269-283. [CrossRef PubMed](#)
- [15] Jutel M, Agache I, Bonini S, Burks AW, Calderon M, Canonica W, Cox L, Demoly P, Frew AJ, O'Hehir R, Kleine-Tebbe J, Muraro A, Lack G, Larenas D, Levin M, Martin BL, Nelson H, Pawankar R, Pfaar O, van Ree R, et al. International consensus on allergen immunotherapy II: Mechanisms, standardization, and pharmacoeconomics. *J Allergy Clin Immunol*. 2016; 137: 358-368. [CrossRef PubMed](#)
- [16] Meadows A, Kaambwa B, Novielli N, Huissoon A, Fry-Smith A, Meads C, Barton P, Dretzke J. A systematic review and economic evaluation of subcutaneous and sublingual allergen immunotherapy in adults and children with seasonal allergic rhinitis. *Health Technol Assess*. 2013; 17: vi, xi-xiv, 1-322. [CrossRef PubMed](#)
- [17] WHO. Coronavirus disease (COVID-19) weekly epidemiological update and weekly operational update. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>. (Date of access: 23.03.2021).
- [18] Pfaar O, Torres MJ, Akdis CA. COVID-19: A series of important recent clinical and laboratory reports in immunology and pathogenesis of SARS-CoV-2 infection and care of allergy patients. *Allergy*. 2021; 76: 622-625. [CrossRef PubMed](#)
- [19] Cardinale F, Ciprandi G, Barberi S, Bernardini R, Caffarelli C, Calvani M, Cavagni G, Galli E, Minasi D, Del Giudice MM, Moschese V, Novembre E, Paravati F, Peroni DG, Tosca MA, Traina G, Tripodi S, Marseglia GL; and the SLAIP Task Force. Consensus statement of the Italian society of pediatric allergy and immunology for the pragmatic management of children and adolescents with allergic or immunological diseases during the COVID-19 pandemic. *Ital J Pediatr*. 2020; 46: 84. [CrossRef PubMed](#)
- [20] Shaker MS, Oppenheimer J, Grayson M, Stukus D, Hartog N, Hsieh EWY, Rider N, Dutmer CM, Vander Leek TK, Kim H, Chan ES, Mack D, Ellis AK, Lang D, Lieberman J, Fleischer D, Golden DBK, Wallace D, Portnoy J, Mosnaim G, et al. COVID-19: pandemic contingency planning for the allergy and immunology clinic. *J Allergy Clin Immunol Pract*. 2020; 8: 1477-1488.e5. [CrossRef PubMed](#)
- [21] Riggioni C, Comberiati P, Giovannini M, Agache I, Akdis M, Alves-Correia M, Antó JM, Arcolaci A, Azkur AK, Azkur D, Beken B, Boccabella C, Bousquet J, Breiteneder H, Carvalho D, De Las Vecillas L, Diamant Z, Eguluz-Gracia I, Eiwegger T, Eyerich S, et al. A compendium answering 150 questions on COVID-19 and SARS-CoV-2. *Allergy*. 2020; 75: 2503-2541. [CrossRef PubMed](#)
- [22] Pfaar O, Klimek L, Jutel M, Akdis CA, Bousquet J, Breiteneder H, Chinthrajah S, Diamant Z, Eiwegger T, Fokkens WJ, Fritsch HW, Nadeau KC, O'Hehir RE, O'Mahony L, Rief W, Sampath V, Schedlowski M, Torres MJ, Traidl-Hoffmann C, Wang Y, et al. COVID-19 pandemic: Practical considerations on the organization of an allergy clinic-An EAACI/ARIA Position Paper. *Allergy*. 2021; 76: 648-676. [CrossRef PubMed](#)
- [23] Klimek L, Pfaar O, Worm M, Eiwegger T, Hagemann J, Ollert M, Untersmayr E, Hoffmann-Sommergruber K, Vultaggio A, Agache I, Bavek S, Bossios A, Casper I, Chan S, Chatzipetrou A, Vogelberg C, Firinu D, Kauppi P, Kolios A, Kothari A, et al. Anwendung von Biologika bei allergischen und Typ-2-entzündlichen Erkrankungen in der aktuellen Covid-19-Pandemie, b, c: Positionspapier des Ärzteverbands Deutscher Allergologen (AeDA)A, der Deutschen Gesellschaft für Allergologie und klinische Immunologie (DGAKI)B, der Gesellschaft für Pädiatrische Allergologie und Umweltmedizin (GPA)C, der Österreichischen Gesellschaft für Allergologie und Immunologie (ÖGAI)D, der Luxemburgischen Gesellschaft für Allergologie und Immunologie (LGAI)E, der Österreichischen Gesellschaft für Pneumologie (ÖGP)F in Kooperation mit der deutschen, österreichischen, und schweizerischen ARIA-GruppeG und der Europäischen Akademie für Allergologie und Klinische Immunologie (EAACI)H. *Allergo J*. 2020; 29: 14-27. [PubMed](#)
- [24] Klimek L, Pfaar O, Worm M, Bergmann KC, Bieber T, Buhl R, Buters J, Darsow U, Keil T, Kleine-Tebbe J, Lau S, Maurer M, Merk H, Mösges R, Saloga J, Staubach P, Stute P, Rabe K, Rabe U, Vogelmeier C, et al. Allergen immunotherapy in the current COVID-19 pandemic: A position paper of AeDA, ARIA, EAACI, DGAKI and GPA: Position paper of the German ARIA GroupA in cooperation with the Austrian ARIA GroupB, the Swiss ARIA GroupC, German Society for Applied Allergology (AEDA)D, German Society for Allergology and Clinical Immunology (DGAKI)E, Society for Pediatric Allergology (GPA)F in cooperation with AG Clinical Immunology, Allergology and Environmental Medicine of the DGHNO-KHCG and the European Academy of Allergy and Clinical Immunology (EAACI) H. *Allergol Select*. 2020; 4: 44-52. [CrossRef PubMed](#)
- [25] Klimek L, Bergmann KC, Brehler R, Pfützner W, Worm M, Hartmann K, et al. Allergologische Diagnostik und Therapien bei COVID-19 Impfungen: Praktische Handlungsempfehlungen. Empfehlungen von AeDA, DGAKI und GPA. *Allergo J. Int*. 2011; 19: 1-17, DOI 10.1007/s40629-021-00165-7.
- [26] Pfaar O, Klimek L, Hamelmann E, Kleine-Tebbe J, Taube C, Wagenmann M, Werfel T, Brehler R, Novak N, Mülleneisen N, Becker S, Worm M. COVID-19 vaccination of patients with allergies and type-2 inflammation with concurrent antibody therapy (biologics) – A Position Paper of the German Society of Allergology and Clinical Immunology (DGAKI) and the German Society for Applied Allergology (AeDA). *Allergol Select*. 2021; 5: 140-147. [CrossRef PubMed](#)
- [27] EMA. Anhang I – Zusammenfassung der Merkmale des Arzneimittels – Comirnaty Konzentrat zur Herstellung einer Injektionsdispersion COVID-19-mRNA-Impfstoff (Nukleosid-modifiziert). https://www.ema.europa.eu/en/documents/product-information/comirnaty-epar-product-information_de.pdf. (Date of access: January 11, 2021).
- [28] EMA. Summary of product characteristics: Comirnaty concentrate for dispersion for injection COVID-19 mRNA Vaccine (nucleoside modified). https://www.ema.europa.eu/en/documents/product-information/comirnaty-epar-product-information_en.pdf. (Date of access: March 23, 2021).

- [29] EMA. Summary of product characteristics: COVID-19 Vaccine AstraZeneca suspension for injection. https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-astrazeneca-epar-product-information_en.pdf. (Date of access: March 23, 2021).
- [30] EMA. Summary of product characteristics: COVID-19 Vaccine Janssen suspension for injection COVID-19 vaccine (Ad26.COVS-2-S [recombinant]). https://www.ema.europa.eu/en/documents/product-information/covid-19-vaccine-janssen-epar-product-information_en.pdf. Date of access: March 23, 2021.
- [31] Worm M, Ring J, Klimek L, Jakob T, Lange L, Treudler R, Beyer K, Werfel T, Biedermann T, Bircher A, Fischer M, Fuchs T, Heller AR, Hoffmann F, Huttegger I, Kopp MV, Kugler C, Lommatzsch M, Pfaar O, Rietschel E, et al. [Covid-19 vaccination and risk of anaphylaxis – Recommendations for practical management]. *MMW Fortschr Med*. 2021; 163: 48-51. [CrossRef](#) [PubMed](#)
- [32] Klimek L, Novak N, Hamelmann E, Werfel T, Wagenmann M, Taube C, Bauer A, Merk H, Rabe U, Jung K, Schlenter W, Ring J, Chaker A, Wehrmann W, Becker S, Mülleneisen N, Nemat K, Czech W, Wrede H, Brehler R, et al. Severe allergic reactions after COVID-19 vaccination with the Pfizer/BioNTech vaccine in Great Britain and USA: Position statement of the German Allergy Societies: Medical Association of German Allergologists (AeDA), German Society for Allergology and Clinical Immunology (DGAKI) and Society for Pediatric Allergology and Environmental Medicine (GPA). *Allergo J Int*. 2021; 30: 51-55. [CrossRef](#) [PubMed](#)
- [33] Klimek L, Worm M, Lange L, Beyer K, Rietschel E, Vogelberg C, Schnadt S, Stöcker B, Brockow K, Hagemann J, Bieber T, Wehrmann W, Becker S, Freudelsperger L, Mülleneisen NK, Nemat K, Czech W, Wrede H, Brehler R, Fuchs T, et al. Management von Anaphylaxie gefährdeter Patienten während der Covid-19-Pandemie. *Allergo J Int*. 2020; 29: 16-26. [CrossRef](#)
- [34] Barker-Tejeda TC, Bazire R, Obeso D, Mera-Berriatua L, Rosace D, Vazquez-Cortes S, Ramos T, Rico MDP, Chivato T, Barbas C, Villaseñor A, Escribese MM, Fernández-Rivas M, Blanco C, Barber D. Exploring novel systemic biomarker approaches in grass-pollen sublingual immunotherapy using omics. *Allergy*. 2021; 76: 1199-1212. [PubMed](#)
- [35] Varona R, Ramos T, Escribese MM, Jimeno L, Galán A, Würtzen PA, Vega F, Marín A, Martín S, Carrera AC, Blanco C, Barber D. Persistent regulatory T-cell response 2 years after 3 years of grass tablet SLIT: Links to reduced eosinophil counts, sIgE levels, and clinical benefit. *Allergy*. 2019; 74: 349-360. [CrossRef](#) [PubMed](#)
- [36] Celebi Sözen Z, Mungan D, Cevhertas L, Ogulur I, Akdis M, Akdis C. Tolerance mechanisms in allergen immunotherapy. *Curr Opin Allergy Clin Immunol*. 2020; 20: 591-601. [CrossRef](#) [PubMed](#)
- [37] Eljaszewicz A, Rucht F, Radzikowska U, Globinska A, Boonpiyathad T, Gschwend A, Morita H, Helbling A, Arasi S, Kahlert H, Berek N, Nandy A, Akdis M, Willers C, Moniuszko M, Akdis CA, Sokolowska M. Trained immunity and tolerance in innate lymphoid cells, monocytes, and dendritic cells during allergen-specific immunotherapy. *J Allergy Clin Immunol*. 2021; 147: 1865-1877. [CrossRef](#) [PubMed](#)
- [38] Golebski K, Layhadi JA, Sahiner U, Steveling-Klein EH, Lenormand MM, Li RCY, Bal SM, Heesters BA, Vilà-Nadal G, Hunewald O, Montamat G, He FQ, Ollert M, Fedina O, Lao-Araya M, Vijverberg SJH, Maitland-van der Zee AH, van Drunen CM, Fokkens WJ, Durham SR, et al. Induction of IL-10-producing type 2 innate lymphoid cells by allergen immunotherapy is associated with clinical response. *Immunity*. 2021; 54: 291-307.e7. [CrossRef](#) [PubMed](#)
- [39] Klimek L, Jutel M, Akdis C, Bousquet J, Akdis M, Bachert C, Agache I, Ansotegui I, Bedbrook A, Bosnic-Anticevich S, Canonica GW, Chivato T, Cruz AA, Czarlewski W, Del Giacco S, Du H, Fonseca JA, Gao Y, Haahtela T, Hoffmann-Sommergruber K, et al; ARIA-MASK Study Group. Handling of allergen immunotherapy in the COVID-19 pandemic: An ARIA-EAACI statement. *Allergy*. 2020; 75: 1546-1554. [CrossRef](#) [PubMed](#)
- [40] Klimek L, Pfaar O, Worm M, Bergmann K-C, Bieber T, Buhl R, Buters J, Darsow U, Keil T, Kleine-Tebbe J, Lau S, Maurer M, Merk H, Mösges R, Saloga J, Staubach P, Poethig D, Rabe K, Rabe U, Vogelmeier C, et al. Allergen-Immuntherapie in der aktuellen Covid-19-Pandemie. *Allergo J*. 2020; 29: 17-25. [CrossRef](#) [PubMed](#)
- [41] Hodgson SH, Mansatta K, Mallett G, Harris V, Emary KRW, Pollard AJ. What defines an efficacious COVID-19 vaccine? A review of the challenges assessing the clinical efficacy of vaccines against SARS-CoV-2. *Lancet Infect Dis*. 2021; 21: e26-e35. [CrossRef](#) [PubMed](#)
- [42] Pollard AJ, Bijker EM. A guide to vaccinology: from basic principles to new developments. *Nat Rev Immunol*. 2021; 21: 83-100. Erratum in: *Nat Rev Immunol*. 2021 Feb; 21: 129. doi: 10.1038/s41577-020-00497-5. [CrossRef](#) [PubMed](#)
- [43] RKI. RKI: Epidemiologisches Bulletin. https://www.rki.de/DE/Content/Infekt/EpidBull/epid_bull_node.html. Date of access: March 23, 2021.
- [44] WHO. Draft landscape and tracker of COVID-19 candidate vaccines. <https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccines>. Date of access: March 23, 2021.
- [45] Krammer F. SARS-CoV-2 vaccines in development. *Nature*. 2020; 586: 516-527. [CrossRef](#) [PubMed](#)
- [46] Logunov DY, Dolzhikova IV, Zubkova OV, Tukhvatulin AI, Shcheblyakov DV, Dzharullaeva AS, Grousova DM, Erokhova AS, Kovyrshina AV, Botikov AG, Izhaeva FM, Popova O, Ozharovskaya TA, Esmagambetov IB, Favorskaya IA, Zrelkin DI, Voronina DV, Shcherbinin DN, Semikhin AS, Simakova YV, et al. Safety and immunogenicity of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine in two formulations: two open, non-randomised phase 1/2 studies from Russia. *Lancet*. 2020; 396: 887-897. Erratum in: *Lancet*. 2021 Jan 9; 397: 98. doi: 10.1016/S0140-6736(20)32721-5. [CrossRef](#) [PubMed](#)
- [47] Blauvelt A, Simpson EL, Tying SK, Purcell LA, Shumel B, Petro CD, Akinlade B, Gadkari A, Eckert L, Graham NMH, Pirozzi G, Evans R. Dupilumab does not affect correlates of vaccine-induced immunity: A randomized, placebo-controlled trial in

- adults with moderate-to-severe atopic dermatitis. *J Am Acad Dermatol.* 2019; 80: 158-167.e1. [CrossRef PubMed](#)
- [48] Zeitlin PL, Leong M, Cole J, Mallory RM, Shih VH, Olsson RF, Goldman M; ALIZE study investigators. Benralizumab does not impair antibody response to seasonal influenza vaccination in adolescent and young adult patients with moderate to severe asthma: results from the Phase IIIb ALIZE trial. *J Asthma Allergy.* 2018; 11: 181-192. [Cross-Ref PubMed](#)
- [49] Durrani SR, Montville DJ, Pratt AS, Sahu S, DeVries MK, Rajamanickam V, Gangnon RE, Gill MA, Gern JE, Lemanske RF Jr, Jackson DJ. Innate immune responses to rhinovirus are reduced by the high-affinity IgE receptor in allergic asthmatic children. *J Allergy Clin Immunol.* 2012; 130: 489-495. [CrossRef PubMed](#)
- [50] Esquivel A, Busse WW, Calatroni A, Togias AG, Grindle KG, Bochkov YA, Gruchalla RS, Kattan M, Kercsmar CM, Khurana Hershey G, Kim H, Lebeau P, Liu AH, Szeftel SJ, Teach SJ, West JB, Wildfire J, Pongracic JA, Gern JE. Effects of omalizumab on rhinovirus infections, illnesses, and exacerbations of asthma. *Am J Respir Crit Care Med.* 2017; 196: 985-992. [CrossRef PubMed](#)
- [51] Jakob T, Klimek L. Allergologie in Zeiten von Covid-19. *Allergo J.* 2020; 29: 3.
- [52] Caubet J-C, Ponvert C. Vaccine allergy. *Immunol Allergy Clin North Am.* 2014; 34: 597-613. [CrossRef PubMed](#)
- [53] Dreskin SC, Halsey NA, Kelso JM, Wood RA, Hummell DS, Edwards KM, Caubet JC, Engler RJ, Gold MS, Ponvert C, Demoly P, Sanchez-Borges M, Muraro A, Li JT, Rottem M, Rosenwasser LJ. International Consensus (ICON): allergic reactions to vaccines. *World Allergy Organ J.* 2016; 9: 32. [CrossRef PubMed](#)
- [54] McNeil MM, DeStefano F. Vaccine-associated hypersensitivity. *J Allergy Clin Immunol.* 2018; 141: 463-472. [CrossRef PubMed](#)
- [55] Nilsson L, Brockow K, Alm J, Cardona V, Caubet J-C, Gomes E, Jenmalm MC, Lau S, Netterlid E, Schwarze J, Sheikh A, Storsaeter J, Skevaki C, Terreehorst I, Zanoni G. Vaccination and allergy: EAACI position paper, practical aspects. *Pediatr Allergy Immunol.* 2017; 28: 628-640. [PubMed](#)
- [56] Wood RA. Allergic reactions to vaccines. *Pediatr Allergy Immunol.* 2013; 24: 521-526.
- [57] Ring J, Beyer K, Biedermann T, Bircher A, Fischer M, Fuchs T, Heller A, Hoffmann F, Hutegger I, Jakob T, Klimek L, Kopp MV, Kugler C, Lange L, Pfaar O, Rietschel E, Rueff F, Schnadt S, Seifert R, Stöcker B, et al. Guideline (S2k) on acute therapy and management of anaphylaxis: 2021 update: S2k-Guideline of the German Society for Allergology and Clinical Immunology (DGAKI), the Medical Association of German Allergologists (AeDA), the Society of Pediatric Allergology and Environmental Medicine (GPA), the German Academy of Allergology and Environmental Medicine (DAAU), the German Professional Association of Pediatricians (BVKJ), the Society for Neonatology and Pediatric Intensive Care (GNPI), the German Society of Dermatology (DDG), the Austrian Society for Allergology and Immunology (ÖGAI), the Swiss Society for Allergy and Immunology (SGAI), the German Society of Anaesthesiology and Intensive Care Medicine (DGAI), the German Society of Pharmacology (DGP), the German Respiratory Society (DGP), the patient organization German Allergy and Asthma Association (DAAB), the German Working Group of Anaphylaxis Training and Education (AGATE). *Allergo J Int.* 2021; 30: 1-25. [CrossRef PubMed](#)
- [58] Bousquet J, Agache I, Blain H, Jutel M, Ventura MT, Worm M, Del Giacco S, Benetos A, Bilo MB, Czarlewski W, Abdul Latiff AH, Al-Ahmad M, Angier E, Annesi-Maesano I, Atanaskovic-Markovic M, Bachert C, Barbaud A, Bedbrook A, Bennoor KS, Berghea EC, et al. Management of anaphylaxis due to COVID-19 vaccines in the elderly. *Allergy.* 2021. Epub ahead of print. [CrossRef PubMed](#)
- [59] Pfaar O, Mahler V. Allergic reactions to COVID-19-vaccinations – unveiling the secret(s). *Allergy.* 2021; 76: 1621-1623. [CrossRef PubMed](#)
- [60] Pfaar O, Agache I, Bonini M, Brough HA, Chivato T, Del Giacco SR, Gawlik R, Gelincik A, Hoffmann-Sommergruber K, Jutel M, Klimek L, Knol EF, Lauerma A, Ollert M, O'Mahony L, Mortz CG, Palomares O, Riggioni C, Schwarze J, Skypala I, et al. COVID-19 pandemic and allergen immunotherapy – an EAACI survey. *Allergy.* 2021. Epub ahead of print. [CrossRef PubMed](#)
- [61] Klimek L, Novak N, Cabanillas B, Jutel M, Bousquet J, Akdis CA. Allergenic components of the mRNA-1273 vaccine for COVID-19: possible involvement of polyethylene glycol and IgG-mediated complement activation. *Allergy.* 2021. Epub ahead of print. [CrossRef PubMed](#)
- [62] Hagemann J, Onorato GL, Jutel M, Akdis CA, Agache I, Zuberbier T, Czarlewski W, Mullol J, Bedbrook A, Bachert C, Bennoor KS, Bergmann KC, Braidó F, Camargos P, Caraballo L, Cardona V, Casale T, Cecchi L, Chivato T, Chu DK, et al. Differentiation of COVID-19 signs and symptoms from allergic rhinitis and common cold: An ARIA-EAACI-GA2 LEN consensus. *Allergy.* 2021; 76: 2354-2366. DOI 10.1111/all.14815. [CrossRef PubMed](#)
- [63] Cabanillas B, Akdis C, Novak N. Allergic reactions to the first COVID-19 vaccine: a potential role of Polyethylene glycol? *Allergy.* 2020. Epub ahead of print. [CrossRef PubMed](#)
- [64] Vultaggio A, Agache I, Akdis CA, Akdis M, Bavbek S, Bossios A, Bousquet J, Boyman O, Chaker AM, Chan S, Chatzipetrou A, Feleszko W, Firinu D, Jutel M, Kauppi P, Klimek L, Kolios A, Kothari A, Kowalski ML, Matucci A, et al. Considerations on biologicals for patients with allergic disease in times of the COVID-19 pandemic: An EAACI statement. *Allergy.* 2020; 75: 2764-2774. [Cross-Ref PubMed](#)