


Journal Pre-proof

Surgical site infections - guidance for elective surgery during the SARS-CoV-2 pandemic - international recommendations and clinical experience

Assadian Ojan, Golling Markus, Krüger Colin M, Leaper David, Mutters Nico T, Roth Beat, Kramer Axel



PII: S0195-6701(21)00070-0

DOI: <https://doi.org/10.1016/j.jhin.2021.02.011>

Reference: YJHIN 6318

To appear in: *Journal of Hospital Infection*

Received Date: 28 October 2020

Revised Date: 10 February 2021

Accepted Date: 10 February 2021

Please cite this article as: Ojan A, Markus G, Colin M K, David L, Nico T M, Beat R, Axel K, Surgical site infections - guidance for elective surgery during the SARS-CoV-2 pandemic - international recommendations and clinical experience, *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2021.02.011>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier Ltd on behalf of The Healthcare Infection Society.

Surgical site infections - guidance for elective surgery during the SARS-CoV-2 pandemic - international recommendations and clinical experience

Assadian Ojan¹, Golling Markus², Krüger Colin M³, Leaper David⁴, Mutters Nico T⁵, Roth Beat⁶, Kramer Axel⁷

¹Regional Hospital Wiener Neustadt, Austria; and Institute for Skin Integrity and Infection Prevention, University of Huddersfield, UK

²Surgical Clinic I, Diakonie-Klinikum Schwäbisch Hall gGmbH

³Immanuel Klinikum Rüdersdorf, Dep. of Surgery, Center for Robotic, Rüdersdorf b. Berlin, Germany

⁴Emeritus Professor of Surgery, University of Newcastle upon Tyne; and Emeritus Professor of Clinical Sciences, ISlaIP, University of Huddersfield, UK

⁵Institute for Hygiene and Public Health, Bonn University Hospital, Bonn, Germany

⁶Emeritus Chief Physician, Surgical Clinic, District Hospital Belp/Switzerland, FMH for General and Trauma Surgery, Wattenwil, Switzerland.

⁷Institute of Hygiene and Environmental Medicine, University Medicine Greifswald, Germany

Summary

Background: The COVID-19 pandemic not only had an impact on public life and healthcare facilities in general, but also affected established surgical workflows for elective procedures. The strategy to protect patients and healthcare workers from infection by SARS-CoV-2 in surgical departments has needed step-by-step development. Based on the evaluation of international recommendations and guidelines, as well as personal experiences in a clinical “hot spot” and in a 450-bed surgical clinic, an adapted surgical site infection (SSI) prevention checklist was needed to develop concise instructions, which described roles and responsibilities of health care professionals that could be used for wider guidance in pandemic conditions.

Method: Publications of COVID-19-related recommendations and guidelines, produced by health authorities and organizations, such as WHO, US-CDC, ECDC, the American College of Surgery and the Robert Koch Institute, were retrieved, assessed and referenced up to

January 31st, 2020. Additionally, clinical personal experiences in Germany were evaluated and considered.

Results: Part 1 of this guidance summarizes the experience of a tertiary care, surgical centre which utilised redundant hospital buildings for immediate spatial separation in a “hot spot” COVID-19 area. Part 2 outlines the successful screening and isolation strategy in a surgical clinic in a region of Germany with outbreaks in surrounding medical centres. Part 3 provides the synopsis of personal experiences and international recommendations suggested for implementation during the COVID-19 pandemic.

Conclusion: Understanding of COVID-19, and SARS-CoV-2-related epidemiology, is constantly and rapidly changing, requiring continuous adaptation and re-evaluation of recommendations. Established national and local guidelines for continuation of surgical services and prevention of SSI require ongoing scrutiny and focused implementation. This manuscript presents a core facility checklist to support medical institutions to continue their clinical and surgical work during the COVID-19 pandemic.

Introduction

By February 9th 2021 accumulated global data indicated that approximately 102 million people had become infected with SARS-CoV-2, approximately 2.2 million had died, and 57 million recovered. A cohort study from Heinsberg, the region with the highest SARS-CoV-2 incidence in Germany, which involved antibody- and swab-PCR-tests of 919 participants in 405 households, showed an infection rate of 15%; 22% of affected individuals were completely asymptomatic, and fatality rate was 0.37% [1].

According to the authors of the COVIDSurg Collaborative [2], who commented on an international cohort study of mortality and pulmonary complications in patients with perioperative SARS-CoV-2 infection undergoing surgery, an estimated 28.4 million elective operations were cancelled because of the disruption caused by COVID-19. The authors opined that it was the correct decision to postpone operations at a time when patients were at risk of being infected with SARS-CoV-2 perioperatively. Indeed, when a patient developed COVID-19 perioperatively, postoperative pulmonary complications occurred in 51.2% of patients and were associated with a 30-day mortality of 23.8%, with pulmonary complications at 38% [2]. Data from the UK show a clear difference between patients, infected with the virus, who were either conservatively treated and those who had laparoscopic appendicectomy. At 30-day follow-up, complications were significantly higher in the operative group ($p < 0.001$). Thus COVID-19 may change the management of acute appendicitis to favour non-operative management, which has now been shown to be safe and effective in the short-term [3]. A meta-analysis confirmed a high rate of postoperative mortality among COVID-19 patients of 20% and a postoperative ICU admission rate of 15% [4]. Consequently, it was suggested that there is an urgent need for investigations by governments and health providers into measures that ensure safety of patients and healthcare workers (HCW), as elective surgery restarts. A clear guideline, which has been adapted for this unique epidemiological challenge, is essential for the protection of patients and HCW against infection. A proposal for guidance is presented in this manuscript, which is based on the authors' experience and the evaluation of the current scientific literature.

Methods

COVID-19-related recommendations and guidelines, produced by health authorities and organizations, such as WHO, US-CDC, ECDC, the American College of Surgery and the Robert Koch Institute, were searched, assessed and referenced [3-26] up to January 31st, 2021. After consideration of all these guidelines the WHO-Patient Care Checklist was selected as a “master document”, modified and adapted, to be shared with partners in relevant medical and clinical fields. Additionally, our own experiences and data from the management of elective surgical procedures in one hospital in a local COVID-19 “hot spot”, and in another hospital where nosocomial outbreaks with temporary closure occurred in 3 hospitals within a radius of 50 km, were included.

Results and Discussion

This document is divided into three parts. In part 1 and 2, the clinical experiences are presented and discussed. In part 3 all recommendations are listed (CORE CHECKLIST) and directly linked to the challenges for interventions undertaken on SARS-CoV-2 patients.

PART 1 – Clinical experiences from a COVID-19 “hot spot” centre in Germany

Local experience: The Diakonie Klinikum (DIAK) is a regional, tertiary care centre, which provides health care to a region of approximately 300,000 inhabitants. The COVID-19 pandemic stressed the resources of the hospital to a degree never seen before. The displacement of standard care soon became evident: the surgical unit came to a complete standstill, with no elective operations, to accommodate the influx of COVID-19 patients in the ICU and general wards. It was soon decided to establish separate “*black*” and “*grey*” areas for confirmed and suspected SARS-CoV-2 infected patients. Fortunately, a recently built hospital complex allowed the reopening of redundant wards in the old hospital, so that COVID-19 patients could be isolated in separate wards. Similarly, the ICU and IMCU (intermediate care unit) were split into separate entities treating COVID-19 and uninfected patients. In both areas, externally diagnosed SARS-CoV-2 patients were delivered into the hospital through separate entrances. Nurses and doctors from other departments had to be

recruited to the ICUs and COVID-19 wards; teaching and training had to be implemented and was supported by a continuously updated software programme on the current knowledge of patient care (SOPHIA®, Standard Operating Procedure Healthcare Information Assistant). This became relevant in surgery with respect to differentiating between a surgical site infection (SSI) and COVID-19- infection, and in intensive care for decision of early compared with late intubation in critical respiratory conditions, or the need for tracheostomy [27]. In the latter, the discussion is ongoing and a clear decision for or against invasive measures is pending [28]. Furthermore, the hospital management had daily update meetings on current local and regional developments, and the decisions from those meetings were communicated through the public relations office and the intranet to hospital staff.

COVID-19 infections: Despite the initial implementation of infection and prevention control measures against COVID-19, 45 (15.2%) of 297 treated patients died, and 47 required ICU care, with 10 (21.3%) deaths. Some patients with respiratory insufficiency were transferred to other hospitals for extracorporeal membrane oxygenation (ECMO). It was initially thought that early intubation was beneficial since prolonged spontaneous breathing, with or without non-invasive assistance, increased respiratory work and worsened the risk of respiratory failure. However, this needed to be balanced against unacceptability of intubating patients too early merely out of concern that the medical staff might be at greater risk of contracting COVID-19 if patients were ventilated non-invasively. A stepwise treatment strategy was later introduced, with appropriate intensive-care monitoring and regulation of all relevant preventive precautions. This depended on the availability of intensive-care beds. In a US single centre comparative cohort study the ventilatory parameters and lung mechanics of consecutive early and late intubated and ventilated patients with COVID-19 ARDS, were measured using descriptive analysis [29]. The authors concluded that larger cohort studies were needed to detect a difference in mortality between early and late intubation. Other studies have found no advantage for early or late intubation [27, 30]. Current mortality data suggest an advantage for patients in whom intubation is completely avoided, but these data may be confounded.

Conclusion: Two aspects determined the successful management of COVID-19: the immediate reopening of redundant wards in an old hospital to separate the patients into white, grey or black areas and the continuous evaluation of sometimes contradicting

evidence (e.g. early compared with late intubation, introduction of tracheostomy) as well as the use of digital tools (SOPHIA) for training and decision making.

PART 2 - Experiences from a 450-bed surgical clinic with immediately instigated management to prevent Covid-19 outbreaks

The Immanuel Hospital is a church-backed community hospital that provides care for approximately 300,000 inhabitants in the state of Brandenburg and Berlin. During the early phase of the pandemic in Berlin the main ICU resources of the University Hospitals were occupied by COVID-19 patients and a massive reduction of elective surgery resulted. A strategy was developed around maintaining dedicated ICU and non-ICU areas for COVID-19 patients that also allowed continuation of elective and emergency surgery for as long as possible.

The following requirements for the surgical clinic were needed:

- definition of the range of surgical procedures which could be continued according to the regulation of the Federal Minister of Health
- creation and management of patient waiting lists
- surgical patient care with reduced intensive care and inpatient bed capacity
- an admission-screening strategy to separate infected, suspected, and non-infected patients
- establishment of care structures for COVID-19 patients in the operating theatre during surgical procedures
- maintaining training and education during the pandemic

The last two points are further explained below.

Admission screening: Prior to elective surgery, patients were asked to record the following symptoms of COVID-19 in a “health diary” starting 5 days prior to admission: fever $>37.8^{\circ}\text{C}$, dry cough, shortness of breath, anosmia and loss of taste, sore throat, headache, weakness, fatigue, tachypnoea with breathing frequency ≥ 30 , hypotension (systolic BP <90 mmHg or diastolic BP ≤ 60 mmHg) and pulse oximetry values of $<85\%$. In the admission interview (in a separate screening area of the hospital) contact with COVID-19 patients, recent travel history, occupation, social contacts and contact with a hot spot or cluster were questioned.

At this consultation Class II surgical masks were routinely worn by HCW and patients, and a minimum 1.5 metres social distancing was maintained; HCW wore an FFP2 respirator if the patient had respiratory symptoms.

Patients with risk factors of symptoms were admitted to the grey area. A COVID-19 test was taken with a combined nose-throat swab (PCR) and, in addition, symptomatic patients underwent a low-dose CT scan of the thorax. If the PCR was positive, the patient was transferred to the black (COVID positive / isolation) area and the decision to operate assessed on the basis of clinical urgency. Acutely symptomatic patients were admitted directly to the black area of the ICU in a single room. If the PCR was negative, patients were transferred to the white area.

After the cumulative incidence of COVID-19 rose to >50 patients/100,000 per week, all patients were screened preoperatively using a PCR test.

As there was a shortage of nurses in Germany before the pandemic, the clinic has been forced to reduce the number of inpatient beds or carry out short-term personnel rescheduling. This did not increase planning security for patients who require intensive medical monitoring and care in the early post-operative phase.

Establishment of isolation care structure: The clinic was separated into a SARS-CoV-2 free ward (white area), wards for suspected infections (grey area) and wards for confirmed infections (black area). In the same way the ICU was divided. All patients and HCW in the white area screened negative for COVID-19; HCW were tested weekly and maintained health diaries with daily documentation of any symptoms. Based on proven efficacy of class II surgical face masks, in combination with social distance [31], patients and HCW were required to wear class II surgical masks, and alcohol-based hand hygiene was promoted.

The most challenging structural requirement for the black area was the creation of specific sluice rooms adjacent to every patient room or the areas required for donning and doffing personal protective equipment (PPE). Nurses and physicians, when possible, only treated COVID-19-positive patients. Additionally, every patient received education and instructions concerning isolation protocols, safety measures and proper hand disinfection procedures.

Patients were only allowed to contact and communicate with friends and family members via mobile telephone, tablets or laptops.

Care of COVID-19 patients during anaesthesia: All symptomatic COVID-19 patients were monitored continually for respiratory and cardiovascular function. If oxygen saturation fell < 94%, supplemental oxygen therapy was started. Advanced monitoring was started if the patient's general condition deteriorated, particularly if there was somnolence, delirium, dyspnoea or reduction in oxygen saturation, as recommended by Mueller et al. [32].

Surgical care of suspected or confirmed COVID-19 patients: The spatial care of patients needed to be reorganised because few clinics had separate surgical units for suspected or confirmed COVID-19 patients. This involved transfer into the operating theatre and ended with the outward transfer. The hygiene requirements of pandemic regulation in Germany required that central areas of the operating theatre (OT) and associated areas, such as induction and recovery rooms, must not be used simultaneously for treatment of COVID-19 and non COVID-19 patients. Therefore, the OT was used for induction of anaesthesia and early post-operative monitoring. The route of COVID-19 patients in the OT pathway was thereby reduced to the OT: preparation for anaesthesia, induction and discharge were all undertaken in the closed OT. The OT also served as a recovery room. Postoperatively, patients were transferred to the isolation area of the intensive care unit or directly to the pre-isolation ward (grey area) if SARS-CoV-2 was suspected or to the isolation ward if SARS-CoV-2 had been proven. To prevent airborne infection, a simple control protocol for the room air-conditioning system (RATS) has been devised and validated. Compared with OT standard ventilation, the RATS protocol was adapted with the air supply reduced from 360 Pa to 200 Pa and the exhaust air increased from 180 Pa to 300 Pa, thus creating a negative flow of 100 Pa whilst the laminar flow ceiling remained active. In the OT, as a minimum FFP2 respirators and 2 pairs of gloves were worn [33] in accordance with the Robert Koch Institute recommendation [34]. The automated OT door activation function was deactivated to reduce the door opening frequency, and the time the patient was in the OT was minimised. Likewise, the number of OT personnel was kept to a minimum, usually one surgeon and one assistant; one physician's assistant and one anaesthetist and one anaesthesia nurse. The work of the circulating nurse in the OT was delegated to the anaesthesia nurse; the

circulating nurse communicated by telephone with the room team for additional requirements. As the risk of exposure to patient-related aerosols was considered to be highest during intubation and extubation, and also during aerosol-generating surgical procedures, the “keep your distance” advice was observed by everyone in the OT, even during surgical procedures. Virucidal gargling was implemented [35].

Conclusion: Since the COVID-19 pandemic began, 23 COVID-19 patients have had surgery: 12 orthopaedic patients with femoral neck fracture or fractures of the pelvis; 8 patients had surgery for acute appendicitis; and one patient each had surgery for a perforated gall bladder, bowel obstruction and a Caesarean section. All operations were performed without transmission of SARS-CoV-2 to the surgical team. This was in contrast to three nearby hospitals where outbreaks amongst HCWs led to temporary closures.

It is believed that the greatest risk for a COVID-19 outbreak in hospital comes from staff and not patients. Admission of planned, as well as emergency, procedures can be managed and controlled as long as there are sufficient protective materials and adequate facility capacity. It is imperative to reduce the regular number of beds for patient care by 20% of regular bed capacity. Medical and nursing staff leave the hospital every day, with family and other outside contacts, and can introduce the virus into the hospital, from asymptomatic carriers of SARS-CoV-2 in particular. Fixed routines of regular self-testing are required, twice weekly for all asymptomatic staff with a PCR or an antigen rapid test, as well as independent maintenance of a diary of symptoms. Symptomatic staff must remain in quarantine until a negative result is available. Although there may be initial shortages of personal protective equipment (PPE), these need to be available in sufficient quantities. All staff in direct patient contact should wear an FFP2 respirator which provides additional protection for all in the event of an unexpected internal COVID-19 outbreak.

PART 3 - CORE CHECKLIST (AIDE MEMOIRE)

General information SARS-CoV-2 patients – hospital precautions

- Demand on healthcare organizations (e.g., hospital/clinics) varies. Therefore, recommendations need to be implemented, and standard operating procedures (SOPs) adapted. A simple software program, installed on mobile devices, is available

for rapid implementation of SOPs and spread of innovative measures and procedures (i.e., www.sophia.online).

- Following any shortfall of supply or stock, an adaptation of these SOPs may be necessary for a limited amount of time.
- The surgical team should, in collaboration with the purchasing department, and based on daily use and consumption, ensure procurement and stock management for personal protection equipment [PPE], together with cleaning and disinfection procedures. The PPE and disinfecting products should have proven or depending on resources definitive efficacy against coronaviruses.
- Traceability of internal and external contact points of patients and limited visitors needs to be ensured.
- Suspected (potentially infected / incubating) and proven (definitive COVID-19 infections) cases have to be isolated from other patients who are cared for in a designated “white area”. Patients with proven COVID-19 infection should be managed in a designated “black area”. However, suspected cases must be isolated in a single room until the result of a SARS-CoV-2 PCR is known (“grey area”). Suspected and proven cases need to be reported to local authorities.
- A list of contact individuals (contact > 15 min and distance < 1.5 m) and traceability of internal and external contact points of patients and limited visitors needs to be maintained.
 - Rotating teams (two or more teams with no physical contact separated on an either daily, weekly or fortnight basis).
 - Monitoring of COVID-19 rates is needed by Health Authority (including basic reproduction rate, and prevalence according to local recommendations).

Before arrival to a clinical setting / health diary and tele-triage

- It is advisable to implement a tele-triage system, before elective surgery, to implement a risk assessment. Alternatively, a special temporary area can be integrated into the entrance area or prior to hospital admission. Uncontrolled patient access, without triage, has to be prevented by strict control of access.

- If possible, 5 days before admission each patient keeps a health diary to record symptoms suggestive of COVID-19, such as fever $>37.8^{\circ}\text{C}$, runny nose, dry cough, shortness of breath, loss of smell and taste, sore throat, headache, weakness, fatigue, tachypnoea with breathing frequency ≥ 30 , hypotension (systolic BP < 90 mmHg or diastolic BP ≤ 60 mmHg) and pulse oximetry (values $<85\%$ are critical).
- The most reliable method to assess the current pre-operative SARS-CoV-2 status of a patient is PCR-testing, ideally 48 to 24 hours prior to any surgical procedure. However, the appropriate timeline for preoperative laboratory screening is not known at present. The Dutch Guidelines, [13] for elective surgery on COVID-19 asymptomatic patients, recommend either no pre-operative SARS-CoV-2 screening or within 48 h before elective surgery, depending on the mean of the 3-day average of new infections in the Netherlands (above or below 40 new infections per day). A recent survey among 479 arthroplasty surgeons from 44 countries and 6 continents agreed that “...it would be ideal for all patients to undergo RT-PCR testing for SARS-CoV-2 before the operation ... [however], if there is a paucity of available tests, then only high-risk patients should be tested” [14].
- Alternatively, a rapid SARS-CoV-2 antigen test, which takes approximately 15-20 minutes, can be performed at the point of care. It indicates virus excretion with a high degree of certainty ($>95\%$) in patients infected with COVID-19, provided the viral load is sufficiently high (i.e., positivity means infectivity). The positive rapid test enables a rapid decision to be made for immediate transfer to the black area (see below) particularly in symptomatic patients and those with a history of contact with SARS CoV infections, and PCR is unnecessary. However, a negative test does not exclude COVID-19 infection, so in this case the procedure described below is analogous to that for a non-feasible PCR test.
- If preoperative PCR testing for each patient is not feasible, then general procedures have to be followed, including a decision about the necessity of PCR testing and the use of PPE by staff before, during and after a surgical procedure.
- *Triage Step 1:* Screening for symptoms of COVID-19 with TOCC (travel history, occupation, social contacts, household members, origin from a hot spot or

cluster, contact with SARS-CoV-2 individuals) using a questionnaire; the result of which enables stratification into different risk groups.

- *Triage Step 2a:* If patients are not identified as being at risk, they still need to be informed, using an agreed protocol, about the required procedures: such as wearing a mouth-nose protection (class II surgical mask or FFP respirator depending on the classification of the patient on leaving a patient's room), proper hand hygiene, preoperative bathing, stopping smoking [36], and nutritional management.
- *Triage Step 2b:* If a patient, based on the questionnaire, is assessed as being at risk, PCR-screening should be arranged prior to admission into hospital.
- *Triage Step 3:* Depending on the epidemiological situation where the patient lives, two risk-groups have to be distinguished: i) in an epidemic timeslot, where the reproduction rate $[R]$ is < 1 , but infectivity still occurs (screening is only recommended when infection is suspected). ii) in an epidemic timeslot, where the reproduction rate $[R]$ is > 1 (every new patient admitted should be screened) or local hot spots are known. If the patient is given a RT-PCR test for SARS-CoV-2, the test should be performed no less than 72 hours, but ideally 24 hours, prior to elective surgery and the result documented in the clinical records.
- *Triage Step 4:* If a patient is tested as being SARS-CoV-2 positive, surgery should be postponed, if possible, with isolation of the patient for 10 - 14 days prior to operation either at the hospital or at home, depending on their medical condition. After the quarantine period a repeated test will verify whether infectivity persists. Admission has to be postponed until diagnostic results are available.
- Except for emergency surgery, and cancer surgery that cannot be postponed from a medical perspective, consider the GRADE system [20, 26] for surgical decisions. In case of the slightest suspicion of an infection or confirmed SARS-CoV-2 infection, surgery should be postponed until recovery. COVID-19 symptoms, including fever and increased temperature, should be constantly monitored until the day of surgery. Frequent SARS-CoV-2 testing should be considered.

At patient arrival ensure patient and staff safety

- The risk of infection and transmission should be minimized by utilizing general principles such as social and physical distancing (at least 1.00 m [37], although the Robert Koch Institute recommends at least 1.50 m [38]), wearing an appropriate mouth-nose-protection as well as proper hand hygiene. Eye protection is also associated with fewer infections [37].
- If tele-triage cannot be executed, as in the case of an emergency, all patients planned for surgery should be submitted for SARS-CoV-2/COVID-19 testing, including temperature and pulse oximetry measurements; an admission can only be allowed into a predefined holding area (*grey area*) until the screening result is available.
- Procedures to separate definitively, and suspected, infected SARS-CoV-2 patients (isolation) from patients who are not considered to be at risk should be established (based on information such as hand and respiratory hygiene, use of PPE and toilet use). Patients with COVID-19-like symptoms should be directed to designated waiting areas and transferred to the *grey area*. SARS-CoV-2 positive patients need to be transferred to a *black area*.
- In case-positive hospitals which do not possess the potential of spatial separation into these three risk-categories, patients need to be assigned and transferred to a regional tertiary hospital which is able to provide spatial separation [32, 33].
- Diagnostic-based screening for potentially SARS-CoV-2 positive patients depends on the availability of appropriate testing. During the initial phase of the pandemic, the backbone of diagnostic testing was laboratory-based PCR investigation of nasopharyngeal specimens. Currently, there are several laboratory- and point of care- (POC) based diagnostic methods available, which are either molecular biology or antigen detection-based. PCR is highly sensitive and specific, but results are not usually available immediately; antigen testing, although less accurate (see below), can give results within 15 minutes.
- The diagnostic efficacy of different commercially-available, antigen lateral-flow POC tests has been evaluated [39-41]. Depending on the diagnostic antibody test used,

sensitivities of 76.6% (95% CI: 62.8% - 86.4%) [40] and 72.6% (95% CI: 64.5% - 79.9%) [42] and specificities of 99.3% (95% CI: 98.6% - 99.6%) [41] and 99.3% (95% CI: 99.7% - 100.00%) [42] have been reported.

- If a rapid pre-surgical SARS-CoV-2 assessment is required without time to wait for a PCR result, POC antigen tests may also be considered. However, only validated POC tests with known sensitivity and specificity should be used, and surgical staff trained and informed in the interpretation of results.

Prevention of nosocomial spread after identification of SARS-CoV-2 in hospitalized patients or health care workers (HCWs)

- Decisions on quarantine and further activities of HCW (figure 1) and transfer of contact patients in the grey area, and infection control by PCR at intervals of 5 days, depends on type of contact. If a patient is tested positive for SARS-CoV-2 by an RT-PCR test after a surgical procedure, all HCW who had contact with the patient, without using PPE, and who are not known to have antibody against SARS-CoV-2, should be tested and quarantined for 8 days until their test results of the 5th day of quarantine become available. Decisions to quarantine staff should be made in consultation with the hospital Infection Prevention Control team and the Occupational Health department. The patient should also be isolated and any contact with the patient should include the use of full PPE.
- Nurses and physicians should, if possible, only treat COVID-19-positive patients.

Spatial requirements for the operating area and operating procedure

- **Operating areas:** If possible, place SARS-CoV-2 positive (CoV+) and negative (CoV-) patients into spatially separated operating areas. Central areas of surgical departments (e.g., induction areas/recovery rooms) should not be used simultaneously with CoV+ patients. Nevertheless, the anaesthetic room/operating theatre/recovery room has to be accessible for induction of anaesthesia and for early postoperative surveillance and monitoring. The pathway of a CoV+ patient in the operating unit is therefore constrained to individual operating theatres with direct access. The induction of anaesthesia, intubation and extubation should be undertaken in a closed room of the

operating unit which could also act as recovery room for patients. Alternatively, if a separate room is not available, intubation can take place in the OT where only the anaesthetic staff, wearing FFP2/3 respirators, are present. After intubation the surgical team can enter the OT, wearing class II surgical masks after 15 minutes, by which time the ventilation system will have diluted the aerosolized viral load in the OT. The post-operative transfer of a recovered CoV+ patient should take place either to the isolation area of the ICU (black area) or directly to the pre-isolation area or to the isolation ward.

- **Ventilation:** No evidence is available upon which to base a risk assessment of ventilation systems, but the following conclusions can be drawn from the results of the assessment of different room airflow patterns [42]:
 - In operating theatres (OT) with laminar air flow (LAF) additional protection for the surgical team and patients is ensured by directed ventilation in the operating area. If the exhaust air from the OT flows directly to the outside via the exhaust-air device, the OT can be used without restrictions. The usual protective measures would be sufficient in the OT and a switch to negative pressure in the OT [42] should reduce the risk further [22].
 - In an OT with mixed ventilation, due to the significantly lower ventilation volume with swirling of aerosols, it is questionable whether the FFP3 respirator ensures a seal tight enough to exclude putting the team at risk. In this event, protection of the surgical team can be achieved by wearing overpressure body exhaust suits [43].
 - OTs without mechanical ventilation are not acceptable, since released aerosols are not diluted.
- **Open surgery:** It has been suggested, that during the COVID-19 pandemic that a return to open, instead of laparoscopic surgery, is made as there may be less risk of aerosols with faster operating times. No study has demonstrated the ability for a virus to be transmitted during a surgical procedure whether open or laparoscopic [44]. Even if laparoscopic surgery is associated with a lower risk of SSI, faster healing

time and lower risk for incisional hernia, an advantage of open surgery may be the isobaric setting around the surgical field. However, during any use of electrocoagulation, tissue-specific aerosols may develop.

- **Insufflation:** Laparoscopic, minimally invasive techniques produce aerosols derived from the induced pneumoperitoneum. The risk of an aerosol carryover is reduced by insufflation-systems equipped with smoke-gas elimination (smoke evacuation) and defined CO₂ feeding and discharge; therefore, these systems are to be preferred. Alternatively, it is recommended that older insufflator-instruments are used which have disposable smoke gas-filters (acc. to ISO 29463 [45]) with a Luer-taper connection to remove smoke-gases by filtering.
- **Robotic-assisted surgery:** This allows further reduction of the number of surgeons and assistants in the high-risk field around the patient. Only the first assistant in the field of intervention is needed to change instruments, or for introducing a suture or swab into the operating field. The operator is working from a console positioned at any distance from the operating area.

Pre- and intraoperative assessment

- Any close contact with patients should include use of appropriate PPE.
- Preoperative antibiotic prophylaxis should be given according to accepted local guidelines. Selection and administration of preoperative antibiotic prophylaxis, to prevent SSIs, is not specifically related to SARS-CoV-2.
- Surgical face masks should be routinely replaced prior to each operation, if visibly dirty or moist and after 2 hours of surgery.
- Observe correct hand hygiene when replacing masks.
- A number of precautions, in addition to normal practice, should be taken by the operative staff, including:
 - Prior to intubation, rinsing of the oral cavity with 1.25% aqueous povidone-iodine-solution (PVP-I), in combination with gargling if possible, should be considered; if PVP-I is contraindicated because of allergy or thyroid disorders, the combination of ethanol 21-27% with etheric oils may be applied [35],

- Limiting movement and number of people and equipment in the OT should be encouraged with cautious use of electrocautery and parallel use of suction to remove smoke, labelling specimens, for example for histology, correctly using local regulations and biohazard precautions.
- Decisions on the need for antivirals should be made, such as aqueous povidone-iodine-based formulations (according to the WHO April 2020 decision for inclusion of povidone-iodine-gargling solutions to the list of experimental treatments for COVID-19 [46], because gargle and mouth wash (PVP-I 1%) and throat spray (PVP-I 0.45%) are active against SARS-CoV-2 within 30s [48].
- It is recommended, in regions with a high prevalence of COVID-19, that surgery on patients who have not been RT-PCR-tested for SARS-CoV-2, surgeons, and the entire surgical team who perform preoperative skin antiseptic and patient pre-operative draping, wear particulate filter respirators with an exhalation valve (filtering face piece FFP2, or FFP3). During intubation as well as extubation anaesthetists and anaesthetic nurses must wear an FFP3-respirator with exclusion of the surgical team.
- Once intubation or extubation are completed, the surgical team should wear class II surgical face masks and, if available, a face shield with a neck cover that wraps around the face with an extension which can be placed inside the gown, head protection and 2 pairs of gloves [48].

Behaviour in the intensive care unit (ICU)

- If using aerosol-generating procedures (including intubation, bronchoscopy, cardiopulmonary resuscitation, or suction), pre-procedural virucidal gargling and virucidal nasal antisepsis (spray) is recommended [35].
- Allow entry of essential staff only.
- Wear a particulate respirator (filtering face piece, FFP2 or FFP3) if available, together with gown, eye protection and gloves.

Postoperative Assessment

- Clean and disinfect personal/dedicated patient equipment between patients.
 - Remove any PPE (gloves, gown, mask, eye protection) using correct hand hygiene during any replacement. During the 2014 Ebola-outbreak, it became vitally clear to use a correct dress code, putting on and taking off of PPE, to prevent infection.
 - It is recommended that members of the hygiene-team should be trained to use the dress/undress code of the PPE [32].
 - Dispose of viral-contaminated waste as clinical waste. Prevention of needle stick injury should involve use of suitable containers.

After discharge from hospital

- Provide instruction and materials to patients and caregivers on the continuous need for respiratory hygiene/cough etiquette, physical distancing as well as hand hygiene and mask-wearing principles.
- Provide advice on home isolation if necessary, infection control and limiting social contact, i.e., respect physical/social distancing.
- Record patient address and telephone number for traceability.
- All patients should wear an approved surgical mask class II postoperatively within the hospital.
- Include basic information about maintaining hygiene and monitoring incisional wounds as a routine part of discharge service.

Embedding protective measures to prevent SARS-CoV-2 transmission to surgical patients and the surgical team into the strategy of preventing SSI

All accepted measures for the prevention of SSI (the SSI-bundle approach) need to be undertaken consistently as usual, whether patients with confirmed or questionable COVID-19 infection are involved. This requires the implementation of general recommendations for

the prevention of SSI provided by WHO, CDC, NHS and RKI [49-52], which are summarized in Figure 2.

Conclusions

The COVID-virus is here to stay. Different vaccines are available, but the protective effect is unclear, especially with the spread of new mutants. At present it is important to:

- avoid operating on any patient with COVID-19 (carrier, clinical suspicion or manifestly infected), if the situation allows
- prepare hospitals and staff for pandemic situations such as COVID-19 (emergency plans, training of nurses/doctors, establishment of isolated wards, etc.) - it is possible to provide safe surgical care for SARS-CoV-2-positive patients and to minimize nosocomial transmission [53-55]
- learn to live with and deal with COVID-19 without risking dramatic economic upsets.

Conflict of interest

None declared.

Authors' contributions

OA, DL, NTM and AK drafted the manuscript together with BR and MG. MG has included experiences from a clinical "hot spot-centre", CK from a surgical clinic.

References

1. Streeck H, Schulte B, Kuemmerer B, Richter E, Hoeller T, Fuhrmann C, et al. Infection fatality rate of SARS-CoV-2 infection in a German community with a super-spreading event. med. medRxiv preprint. <https://doi.org/10.1101/2020.05.04.20090076>

2. COVIDSurg Collaborative. Mortality and pulmonary complications in patients undergoing surgery with perioperative SARS-CoV-2 infection: an international cohort study. *Lancet* 2020; 396: 27–38. [https://doi.org/10.1016/S0140-6736\(20\)31182-X](https://doi.org/10.1016/S0140-6736(20)31182-X)
3. Javanmard-Emamghissi H, Boyd-Carson H, Hollyman M, Doleman B, Adiamah A, Lund JN, et al. The management of adult appendicitis during the COVID-19 pandemic: an interim analysis of a UK cohort study. *Tech Coloproctol* 2020. <https://doi.org/10.1007/s10151-020-02297-4>
4. Abate SM, Mantefardo B, Basu B. Postoperative mortality among surgical patients with COVID-19: a systematic review and meta-analysis. *Patient Saf Surg* 2020; 14: 37. <https://doi.org/10.1186/s13037-020-00262>
5. Parvizi J, Gehrke T, Krueger CA, Chisari E, Citak M, Van Onsem S, Walter WL; International Consensus Group (ICM) and Research Committee of the American Association of Hip and Knee Surgeons (AAHKS). Resuming elective orthopaedic surgery during the COVID-19 pandemic: Guidelines developed by the International Consensus Group (ICM). *J Bone Joint Surg Am* 2020; 102(14):1205-12. Erratum in: *J Bone Joint Surg Am* 2020;102(19):e113. doi: 10.2106/JBJS.20.00844
6. AIOT Alliance of International Organizations of Orthopedics & Traumatology. AIOT Joint Statement – COVID-19 Best Practices. April 28, 2020. <https://www.efort.org/aiot-joint-statement-COVID-19/>
7. Prada C, Chang Y, Poolman R, Johal H, Bhandari M; OrthoEvidence Best Practices Working Group. Best practices for surgeons - COVID-19 evidence-based scoping review. May 30, 2020. Version 3.0. https://myoe.blob.core.windows.net/docs/COVID19/OE_Best_Practices_for_Sureons_COVID-19.pdf
8. Robert Koch Institute. Options for separate management of COVID-19 suspect cases / cases and other patients in the outpatient and pre-stationary areas. https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Getrennte_Patient_versorgung.html. July 6, 2020.
9. American College of Surgeons. Post-COVID-19 readiness checklist for resuming surgery. April, 2020 -> <https://www.facs.org/COVID-19/checklist>
10. ECDC. Checklist for hospitals preparing for the reception and care of coronavirus 2019 (COVID-19) patients. Technical Report. February 26, 2020.

- <https://www.ecdc.europa.eu/en/publications-data/checklist-hospitals-preparing-reception-and-care-coronavirus-2019-COVID-19>
11. WHO. Hospital readiness checklist for COVID-19. Regional Office for Europe, Copenhagen, 2020.
<https://apps.who.int/iris/bitstream/handle/10665/333972/WHO-EURO-2020-1012-40758-54954-eng.pdf>
 12. MedTech Europe COVID-19 Implications for clinical investigations needed for medical device recertification. May 2020. https://www.medtecheurope.org/wp-content/uploads/2020/05/20200527_MedTechEurope_COVID-19_Implications_for_Clinical_Investigations.pdf
 13. Nederlandse Vereniging voor Heelkunde. Practice guideline pre-operative work-up for SARS-CoV-2 infection in asymptomatic patients scheduled for surgery under general anesthesia. June 11, 2020, version 4.
https://www.demedischspecialist.nl/sites/default/files/leidraad_preoperatieve_work_up.pdf
 14. Kort NP, Barrena EG, Bédard M, Donell S, Epinette JA, Gomberg B, et al. Recommendations for resuming elective hip and knee arthroplasty in the setting of the SARS-CoV-2 pandemic: the European Hip Society and European Knee Associates Survey of Members. *Knee Surg Sports Traumatol Arthrosc* 2020;28(9): 2723–9.
 15. Donell ST, Thaler M, Budhiparama NC, Buttaro MA, Chen AF, Diaz-Ledezma C, et al. Preparation for the next COVID-19 wave: The European Hip Society and European Knee Associates recommendations. *Knee Surg Sports Traumatol Arthrosc* 2020; 28: 2747–55. <https://doi.org/10.1007/s00167-020-06213-z>
 16. CDC. Interim infection prevention and control recommendations for healthcare personnel during the coronavirus disease 2019 (COVID-19) Pandemic. 2020.
<https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html>
 17. American College of Surgeons. COVID-19: Elective case triage guidelines for surgical care. March 24, 2020. <https://www.facs.org/COVID-19/clinical-guidance/elective-case>

18. American College of Surgeons. COVID-19: Recommendations for Management of Elective Surgical Procedures. March 13, 2020. <https://www.facs.org/COVID-19/clinical-guidance/elective-surgery>
19. American College of Surgeons, American Society of Anesthesiologists, Association of periOperative Registered Nurses, American Hospital Association. Joint Statement: Roadmap for maintaining essential surgery during COVID-19 pandemic. August 10, 2020. https://www.facs.org/-/media/files/COVID19/joint_statement_on_roadmap_to_maintaining_essential_surgery.ashx
20. Moletta L, Pierobon ES, Capovilla G, Costantini M, Salvador R, Merigliano S, et al. International guidelines and recommendations for surgery during COVID-19 pandemic: A Systematic Review. *Int J Surg* 2020;79: 180-8. doi:10.1016/j.ijssu.2020.05.061
21. Royal College of Surgeons of England: Report: Protecting surgery through a second wave. October, 2020. <https://www.rcseng.ac.uk/coronavirus/protecting-surgery-through-a-second-wave/>
22. Schultz P, Morvan JB, Fakhry N, Morinière S, Vergez S, Lacroix C, et al.; French Society of Otorhinolaryngology, Head, Neck Surgery (SFORL); French Society of Head, Neck Carcinology (SFCCF). French consensus regarding precautions during tracheostomy and post-tracheostomy care in the context of COVID-19 pandemic. *Eur Ann Otorhinolaryngol Head Neck Dis* 2020;137(3):167-9. doi: 10.1016/j.anorl.2020.04.006.
23. Park JS, El-Sayed IH, Young VN, Pletcher SD. Development of clinical care guidelines for faculty and residents in the era of COVID-19. *Head Neck*. 2020; 42(7):1403-8. doi: 10.1002/hed.26225.
24. Mouton C, Hirschmann MT, Ollivier M, Seil R, Menetrey J. COVID-19 - ESSKA guidelines and recommendations for resuming elective surgery. *J Exp Orthopaeds* 2020 7:1 Article Number: 28.
25. Heffernan DS, Evans HL, Huston JM, Claridge JA, Blake DP, May AK, et al. Surgical Infection Society guidance for operative and peri-operative care of adult patients infected by the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). *Surg Infect* 2020; 21(4): 301-8.

26. O'Leary MP, Choong KC, Thornblade LW, Fakih MG, Fong Y, Kaiser AM. Management considerations for the surgical treatment of colorectal cancer during the global Covid-19 Pandemic. *Ann Surg* 2020;272(2):e98-e105. doi: 10.1097/SLA.0000000000004029.
 27. Lee YH, Choi KJ, Choi SH, Lee SY, Kim KC, et al. Clinical significance of timing of intubation in critically ill patients with COVID-19: A multi-center retrospective study. *J Clin Med* 2020, 9 (9): 2847. <https://doi.org/10.3390/jcm9092847> 27
 28. Avilés-Jurado FX, Prieto-Alhambra D, González-Sánchez N, de Ossó J, et al. Timing, complications, and safety of tracheotomy in critically ill patients with COVID-19. *JAMA Otolaryngol Head Neck Surg* 2020;147(1):1-8.
 29. Pandya A, Kaur NA, Sacher D, O'Corragain O, Salerno D, Desai P, et al. Ventilatory mechanics in early vs late intubation in a cohort of COVID-19 patients with acute respiratory distress syndrome: a single center's experience *Chest* 2020; S0012-3692(20)34314-2. doi:10.1016/j.chest.2020.08.2084
 30. Matta A, Chaudhary S, Bryan LK, DeJoy RD, Gul F, Torres R, et al. Timing of intubation and its implications on outcomes in critically ill patients with coronavirus disease 2019 infection. *Crit Care Explorat* 2020; 2(10):e0262
 31. Offeddu V, Yung CF, Low MSF, Tam CC. Effectiveness of masks and respirators against respiratory infections in healthcare workers: a systematic review and meta-analysis. *Clin Infect Dis* 2017;65(11):1934–42.
 32. Mueller M, Boettner F, Karczewski D, Janz V, Felix S, Kramer A, et al. Dealing with the COVID-19 pandemic in orthopaedics: experiences and procedure in Germany. *Bone Joint Open* 2020;1(6):309–15
 33. Krüger CM, Kramer A, Türlér A, Riediger H. Can surgery follow the dictates of the pandemic “keep your distance”? Requirements with COVID-19 for hygiene, resources and the team. *GMS Hyg Infect Control* 2020;15: Doc19. doi: 10.3205/dgkh000354
 34. Robert Koch-Institut (RKI). Empfehlungen des RKI zu Hygienemaßnahmen im Rahmen der Behandlung und Pflege von Patienten mit einer Infektion durch SARS-CoV-2. [last updated 2020 Jun 5, last accessed 2020 Jul 24]. Available from: https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Hygiene.html
- External link

35. Kramer A, Eggers M, Hübner NO, Walger P, Steinmann E, Exner M. Virucidal gargling and virucidal nasal spray. *GMS Hyg Infect Control*. 2021; 16:Doc02. DOI: 10.3205/dgkh000373,
36. Sørensen LT. Wound healing and infection in surgery. The clinical impact of smoking and smoking cessation: a systematic review and meta-analysis. *Arch Surg* 2012;147(4):373-83. doi: 10.1001/archsurg.2012.5.
37. Chu DK, Akl EA, Duda S, Solo K, Yaacoub S, Schünemann HJ, on behalf of the COVID-19 Systematic Urgent Review Group Effort (SURGE) study authors. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *Lancet* 2020; 395 (10242): 1973-87. [https://doi.org/10.1016/S0140-6736\(20\)31142-9](https://doi.org/10.1016/S0140-6736(20)31142-9)
38. https://www.rki.de/DE/Content/InfAZ/N/Neuartiges_Coronavirus/Hygienemassnahmen_Einsatzkraefte.pdf?__blob=publicationFile
39. Corman VM, Haage VC, Bleicker T, Schmidt ML, Mühlemann B, Zuchowski M, et al. Comparison of seven commercial SARS-CoV-2 rapid Point-of-Care Antigen tests. *medRxiv*. 2020:2020.11.12.20230292.
40. Krueger LJ, Gaeddert M, Koeppel L, Bruemmer L, Gottschalk C, Miranda IB, et al. Evaluation of the accuracy, ease of use and limit of detection of novel, rapid, antigen-detecting point-of-care diagnostics for SARS-CoV-2 2020.
41. Gremmels H, Winkel BMF, Schuurman R, Rosingh A, Rigter NAM, Rodriguez O, et al. Real-life validation of the Panbio COVID-19 Antigen Rapid Test (Abbott) in community-dwelling subjects with symptoms of potential SARS-CoV-2 infection 2020.
42. Kramer A, Külpmann R, Brunner A, Müller M, Wassilew G. Risk assessment of mixed and displacement ventilation (LAF) during orthopedic and trauma surgery on COVID-19 patients with increased release of infectious aerosols. *GMS Hyg Infect Control* 2020; 15: Doc07.
43. Makovicka JL, Bingham JS, Patel KA, Young SW, Beauchamp CP, Spangehl MJ. Surgeon personal protection: an underappreciated benefit of positive-pressure exhaust suits. *Clin Orthop Relat Res*. 2018;476(6):1341–8.
44. Vigneswaran Y, Prachand VN, Posner MC, Matthews JB, Hussain M. What is the appropriate use of laparoscopy over open procedures in the current COVID-19 climate? *J Gastrointest Surg* 2020; 24(7):1686-91. doi: 10.1007/s11605-020-04592-9.

45. ISO 29463. High-efficiency filters and filter media for removing particles in air - Part 2: Aerosol production, measuring equipment and particle-counting statistics. Berlin: Beuth, 2019.
46. World Health Organization. WHO COVID-19. Experimental Treatments. Classification of treatment types. World Health Organization 2020: R&D Blue-print. <https://www.who.int/docs/default-source/coronaviruse/COVID-classification-of-treatment-types-rev.pdf>
47. Anderson DE, Sivalingam V, Kang AEZ, Ananthanarayanan A, Arumugam H, Jenkins TM, et al. Povidone-iodine demonstrates rapid in vitro virucidal activity against SARS-CoV-2, the virus causing COVID-19 disease. *Infect Dis Ther* 2020; 9: 669–75. <https://doi.org/10.1007/s40121-020-00316-3>
48. Casanova LM, Rutala WA, Weber DJ, Sobsey MD. Effect of single- versus double gloving on virus transfer to health care workers' skin and clothing during removal of personal protective equipment. *Am J Infect Control* 2012; 40: 369–74.
49. Allegranzi B, Bischoff P, Kubilay Z, de Jonge S, Zayed B. Global guidelines for the prevention of surgical site infection. 2. Aufl, WHO 2018.
50. Berríos-Torres SI, Umscheid CA, Bratzler DW, Leas B, Stone EC, Kelz RR et al. Centers for Disease Control and Prevention guideline for the prevention of surgical site infection, 2017. *JAMA Surg* 2017;152(8):784-91. DOI:10.1001/jamasurg.2017.0904
51. Cooper K, Lamagni T, Harrington P, Wloch C, Hopkins S. Surveillance of surgical site infections in NHS hospitals in England, April 2018 to March 2019. *Public Health England*, December 2019. www.gov.uk/phe.
52. KRINKO. Prävention postoperativer Wundinfektionen. Empfehlung der Kommission für Krankenhaushygiene und Infektionsprävention (KRINKO) beim Robert Koch-Institut. *Bgbl* 2018;61:448–73. <https://doi.org/10.1007/s00103-018-2706-2>
53. Alimoglu O, Erol CI, Kayali A, Acar M, Colapkulu N, Leblebici M, et al. Emergency surgery during covid-19 pandemic; what has changed in practice. *BJS* 2020; 107(12): e581–e582
54. Caliendo V, Picciotto F, Quaglino P, Ribero S. COVID infection and sentinel lymph node procedure for melanoma: Management in a dermato-oncology center in a high-risk pandemic area. *Dermatol Ther* 2020;33(4):e13536. doi:10.1111/dth.13536

55. Jazieh AR, Chan SL, Curigliano G, Dickson N, Eaton V, Garcia-Foncillas J, et al. Delivering cancer care during the covid-19 pandemic: recommendations and lessons learned from ASCO Global Webinars. *JCO Glob Oncol* 2020;6:1461-1471. doi: 10.1200/GO.20.00423.

Journal Pre-proof




Figure 1 Decision on quarantine and further activities of HCWs depending on their exposure to SARS-CoV-2 (based on the recommendations by the Robert Koch Institute Germany 2021).




Figure 2. The SSI pathway summarizes the measures to prevent SSI