

# Finite and infinite series sheet

Gallo Tennis, CoGi / to A Mathematical Room

April 6, 2025

*Problems on series and products over finite and infinite sets: books of Terence Tao's "Analysis I", Donald E. Knuth's "The Art of Computer Programming".  
As well as problems of the Putnam Mathematical Competition and the International Math Olympiad.*

## Convergence of series

1. Let  $\sum_{n=m}^{\infty} a_n$  be a formal series of real numbers. Then  $\sum_{n=m}^{\infty} a_n$  converges if and only if, for every real number  $\varepsilon > 0$ , there exists an integer  $N \geq m$  such that

$$\left| \sum_{n=p}^q a_n \right| \leq \varepsilon \quad \text{for all } p, q \geq N.$$

**Solution.**

- ( $\implies$ ) If  $\sum_{n=m}^{\infty} a_n$  converges then, by definition, the partial sum sequence

$$(S_N)_{N=m}^{\infty} = \left( \sum_{n=m}^N a_n \right)_{N=m}^{\infty}$$

converges to  $L$ . Therefore, for any  $\varepsilon > 0$  there exists some  $M \geq m$  such that for any  $p-1, q \geq M$

$$|S_q - S_{p-1}| \leq \varepsilon.$$

Without loss of generality assume  $p-1 \leq q$ , it follows that

$$\left| \sum_{n=m}^q a_n - \sum_{n=m}^{p-1} a_n \right| = \left| \sum_{n=p}^q a_n \right| \leq \varepsilon.$$

- ( $\impliedby$ ) Doing the reverse process of the direct implication the result follows.

□

2. **(Zero test).** Let  $\sum_{n=m}^{\infty} a_n$  be a convergent series of real numbers. Then we must have  $\lim_{n \rightarrow \infty} a_n = 0$ . To put this another way, if  $\lim_{n \rightarrow \infty} a_n$  is non-zero or divergent, then the series  $\sum_{n=m}^{\infty} a_n$  is divergent.

**Solution.** If  $\sum_{n=m}^{\infty} a_n$  converges then by last problem, for any  $\varepsilon > 0$ , there exists some  $M \geq m$  such that for any  $p, q \geq M$

$$\left| \sum_{n=p}^q a_n \right| \leq \varepsilon. \quad (1)$$

Suppose for the sake of contradiction that  $\lim_{n \rightarrow \infty} a_n = L > 0$  so that for any  $\varepsilon > 0$  there exists some  $M'$  such that for any  $n \geq M'$

$$|a_n - L| < \varepsilon$$

so that in particular  $a_n \geq L - \varepsilon$ .

Therefore, for  $K := \max\{M, M'\}$  and for any  $p, q \geq M$

$$\left| \sum_{n=p}^q a_n \right| \geq \left| \sum_{n=p}^q L - \varepsilon \right| = (q - p + 1) |L - \varepsilon|.$$

Hence, let  $\varepsilon = L/2$ , we see that if the above inequality were true

$$\left| \sum_{n=p}^q a_n \right| \geq \frac{(q - p + 1)L}{2}.$$

Then let  $q = p + 1$ , it follows that

$$\left| \sum_{n=p}^q a_n \right| \geq L > \varepsilon.$$

A contradiction of inequality (1).

□

3. **(Absolute convergence test).** Let  $\sum_{n=m}^{\infty} a_n$  be a formal series of real numbers. If this series is absolutely convergent, then it is also conditionally convergent. Furthermore, in this case we have the triangle inequality

$$\left| \sum_{n=m}^{\infty} a_n \right| \leq \sum_{n=m}^{\infty} |a_n|.$$

**Solution.** If  $\sum_{n=m}^{\infty} |a_n|$  converges to  $L$ , then for any  $\varepsilon > 0$  there exists some  $M \geq m$  such that for any  $p, q \geq M$

$$\left| \sum_{n=p}^q |a_n| \right| \leq \varepsilon.$$

Hence,

$$\sum_{n=p}^q |a_n| \leq \varepsilon.$$

By triangle inequality over finite series, for any  $p, q \geq M$  we have

$$\left| \sum_{n=p}^q a_n \right| \leq \sum_{n=p}^q |a_n|.$$

We conclude that  $\sum_{n=p}^q a_n$  converges. Trivially, we also note that again, by triangle inequality over finite series, each partial sum  $S_N = \sum_{n=m}^N a_n$  is smaller than or equal to  $T_N = \sum_{n=m}^N |a_n|$ , meaning that by comparison principle  $\lim_{N \rightarrow \infty} S_N \leq \lim_{N \rightarrow \infty} T_N$ .

□

4.

5. Let  $(a_n)_{n \geq 1}$  be a sequence of positive real numbers such that the series  $\sum_{n=1}^{\infty} a_n$  converges. Show that the series

$$\sum_{n=1}^{\infty} a_n^{\frac{n}{n+1}}$$

Also converges.

## Series expansions