

Normalization Theory sheet I

Gallo Tennis

March 4, 2025

Exercises in normalization theory from J. Ullman's "Database Systems - The Complete Book". These week's topics are closures of functional dependencies sets, projections of FD's sets, minimal bases, BCNF and 3NF.

Rules about Functional Dependencies

1. **3.2.1:** Consider a relation with schema $R(A, B, C, D)$ and FD's

$$AB \rightarrow C, \quad C \rightarrow D, \quad D \rightarrow A.$$

- (a) What are all the nontrivial FD's that follow from the given FD's?
You should restrict yourself to FD's with single attributes on the right side.
- (b) What are all the keys of R ?
- (c) What are all the superkeys for R that are not keys?

Solution.

- (a) (a)
2. **3.2.2:** Repeat Exercise 3.2.1 for the following schemas and sets of FD's:
 - i) $S(A, B, C, D)$ with FD's $A \rightarrow B$, $B \rightarrow C$, and $B \rightarrow D$.
 - ii) $T(A, B, C, D)$ with FD's $AB \rightarrow C$, $BC \rightarrow D$, $CD \rightarrow A$, and $AD \rightarrow B$.
 - iii) $U(A, B, C, D)$ with FD's $A \rightarrow B$, $B \rightarrow C$, $C \rightarrow D$, and $D \rightarrow A$.

Solution.

- i) The
3. **3.2.4:** Show that each of the following are *not* valid rules about FD's by giving example relations that satisfy the given FD's (following the "if") but not the FD that allegedly follows (after the "then").
 - (a) If $A \rightarrow B$ then $B \rightarrow A$.

(b) If $AB \rightarrow C$ and $A \rightarrow C$, then $B \rightarrow C$.

(c) If $AB \rightarrow C$, then $A \rightarrow C$ or $B \rightarrow C$.

4. **3.2.5** Show that if a relation has no attribute that is functionally determined by all the other attributes, then the relation has no nontrivial FD's at all.

Solution. Let R be a relation schema. By hypothesis, for any set of attributes $X \in R$, $X^+ = X$. So if a nontrivial FD such as $X \rightarrow Y$ existed, it would imply that $Y \subseteq X$, so $X^+ = XY$, a contradiction.

□

5. **3.2.6** Let X and Y be sets of attributes. Show that if $X \subseteq Y$, then $X^+ \subseteq Y^+$, where the closures are taken with respect to the same set of FD's.

Solution. Since $X \subseteq Y$ and $Y \subseteq Y^+$, then necessarily $X \subseteq Y^+$. Since $X \subseteq X^+$, we see that each element of $X^+ \cap X$ is in Y^+ . With this in mind, we have shown the inclusion in the trivial FD's case.

Now for the case of nontrivial dependencies $x \in X^+ \setminus X$ we have to show that $x \in Y^+$. Since x is nontrivial then there exists some $z \in X$ such that $x \not\subseteq z$ and

$$z \rightarrow x.$$

Since $z \in Y$ as well, we conclude that $x \in Y^+$.

□

6. **3.2.7** Prove that $(X^+)^+ = X^+$.

Solution. Since it is obvious that $X^+ \subseteq (X^+)^+$ (the closure of a FD set always contains itself), remains showing that $(X^+)^+ \subseteq X^+$.

Let $x \in (X^+)^+$. Then there exists an $y \in X^+$ such that

$$y \rightarrow x.$$

So there must also exist a $z \in X$ such that

$$z \rightarrow y,$$

hence, by transitivity $z \rightarrow x$. We conclude that $x \in X^+$.

□

7. **3.2.8** We say a set of attributes X is *closed* (with respect to a given set of FD's) if $X^+ = X$. Consider a relation with schema $R(A, B, C, D)$ and an unknown set of FD's. If we are told which sets of attributes are closed, we can discover the FD's. What are the FD's if:

- (a) All sets of the four attributes are closed.
- (b) The only closed sets are \emptyset and $\{A, B, C, D\}$.
- (c) The closed sets are \emptyset , $\{A, B\}$, and $\{A, B, C, D\}$.

Solution.

- (a) By exercise 3.2.5 R has no non-trivial dependencies. So $\{A, B, C, D\}^+ = \{A, B, C, D\}$.
- (b) We claim that there does exist at least one superkey of the relation. First we note that the unordered pairs of attributes is

$$\binom{4}{2} > 4,$$

so there must exist at least two pairs X, Y such that $X \cap Y \neq \emptyset$ (by pigeonhole principle). In other words, since the closure of any attribute is at least two (they are not closed) we see that

$$X^+ = XZ \text{ and } Y^+ = YZ$$

or

$$X^+ = XZ \text{ and } Z^+ = ZY$$

So in the first case

$$X \rightarrow Z \text{ and } Y \rightarrow Z$$

then

$$XY \rightarrow Z.$$

In fact, since neither Z is closed, $Z \rightarrow T$, hence by transitivity $XY \rightarrow T$, we conclude that $(XY)^+ = XYZT = ABCD$.

In the second case, we get that X is a superkey by transitivity.

So the FD's will look like $\{AB \rightarrow C, C \rightarrow D\}$ or $\{A \rightarrow B, B \rightarrow C, C \rightarrow D, D \rightarrow A\}$ with respect to both cases.

- (c) Consider the sub relations $R_1 = \pi_{A,B}(R)$ and $R_2 = \pi_{C,D}(R)$. For R_1 , since $A^+ \neq A$ and $B^+ \neq B$ but $(AB)^+ = AB$ it will suffice letting $A \rightarrow B$ and $B \rightarrow A$. In fact, this is the only way it can be done; as if we let $A \rightarrow X$ with $X \subseteq R_2$ then

$$AB \rightarrow A \rightarrow X, \text{ so } (AB)^+ = ABX \neq AB.$$

For R_2 we have that $C^+ \neq C$ and $D^+ \neq D$ and that $(CD)^+ \neq CD$, since these are not closed. We then see that CD is a superkey by seeing that since

$$CD \rightarrow C, CD \rightarrow D, C \rightarrow X, D \rightarrow Y \text{ (with } X \text{ and } Y \text{ not necessarily different),}$$

then

$$CD \rightarrow XY.$$

And if $XY = X$, then since $(CD)^+ = CDX$, but since CDX is not closed, by transitivity $(CD)^+ = CDXZ = ABCD$. Then, the FD's for R_2 will be of the form $\{C \rightarrow D, D \rightarrow A\}$, $\{C \rightarrow A, D \rightarrow B\}$ or $\{C \rightarrow A, D \rightarrow A, CD \rightarrow B\}$.

□

8. **Exercise 3.2.11** Show that if an FD F follows from some given FD's, then we can prove F from the given FD's using Armstrong's axioms (defined in the box "A Complete Set of Inference Rules" in Section 3.2.7). *Hint:* Examine Algorithm 3.7 and show how each step of that algorithm can be mimicked by inferring some FD's by Armstrong's axioms.

Solution. Let $R = \{A_1, \dots, A_n\}$ be a relation. Let $\{B_1, \dots, B_m\}$ be a subset of R . Suppose F had several attributes $\{C_1, \dots, C_k\}$ on the right hand side, Algorithm 3.7 says F holds even when splitting it into several one-attributed FD's, in other words

$$B_1, \dots, B_m \rightarrow F \iff \begin{cases} B_1, \dots, B_m \rightarrow C_1 \\ \vdots \\ B_1, \dots, B_m \rightarrow C_k \end{cases}.$$

There are now some cases we are ought to study; first suppose that some C_i were an element of B_1, \dots, B_m for some $m \leq n$, then

$$\{C_i\} \subseteq \{B_1, \dots, B_m\} \implies B_1, \dots, B_m \rightarrow C_i$$

More generally if we had a larger subset of F , say $\{C_1, \dots, C_q\}$ with $q \leq k$ and each element of it satisfying the above relation, then by union rule we can say that

$$\{C_1, \dots, C_q\} \subseteq \{B_1, \dots, B_m\} \implies B_1, \dots, B_m \rightarrow C_1, \dots, C_q,$$

verifying reflexivity.

Let $\{B_1, \dots, B_m\} \rightarrow C_i$ such that $C_i \rightarrow C_j$ for some i, j . Then we see that

$$C_i \in \{B_1, \dots, B_m\}^+$$

so

$$B_1, \dots, B_m, C_i \rightarrow C_j.$$

Meaning that $C_j \in \{B_1, \dots, B_m, C_i\}^+ = \{B_1, \dots, B_m\}^+$. In other words

$$B_1, \dots, B_m \rightarrow C_i \text{ and } C_i \rightarrow C_j \implies B_1, \dots, B_m \rightarrow C_j,$$

verifying transitivity.

Lastly, since $B_1, \dots, B_m \rightarrow F$, necessarily for any other subset X of R , we have

$$B_1, \dots, B_m, X \rightarrow B_1, \dots, B_m \rightarrow F \text{ (by reflexivity)}$$

and

$$B_1, \dots, B_m, X \rightarrow X. \text{ (by reflexivity)}$$

So by union rule

$$B_1, \dots, B_m, X \rightarrow FX,$$

verifying augmentation.

□

Boyce-Codd Normal Form

1. **3.3.1:** For each of the following relation schemas and sets of FD's:

- (a) $R(A, B, C, D)$ with FD's $AB \rightarrow C$, $C \rightarrow D$, and $D \rightarrow A$.
- (b) $R(A, B, C, D)$ with FD's $B \rightarrow C$ and $B \rightarrow D$.
- (c) $R(A, B, C, D)$ with FD's $AB \rightarrow C$, $BC \rightarrow D$, $CD \rightarrow A$, and $AD \rightarrow B$.
- (d) $R(A, B, C, D)$ with FD's $A \rightarrow B$, $B \rightarrow C$, $C \rightarrow D$, and $D \rightarrow A$.
- (e) $R(A, B, C, D, E)$ with FD's $AB \rightarrow C$, $DE \rightarrow C$, and $B \rightarrow D$.
- (f) $R(A, B, C, D, E)$ with FD's $AB \rightarrow C$, $C \rightarrow D$, $D \rightarrow B$, and $D \rightarrow E$.

do the following:

- i) Indicate all the BCNF violations. Do not forget to consider FD's that are not in the given set, but follow from them. However, it is not necessary to give violations that have more than one attribute on the right side.
- ii) Decompose the relations, as necessary, into collections of relations that are in BCNF.

Solution.

- (a) Doing the closure of C : $C^+ = CDA \neq R$. So we split R in

$$\begin{aligned} R &= R_1 \bowtie R_2, \\ \text{where } R_1 &= CDA \text{ and} \\ R_2 &= C \cup (R \setminus C^+) = C \cup (ABCD \setminus CDA) = CB. \end{aligned}$$

We see that R_2 is in BCNF as it only have 2 attributes, and $R_1 = CDA$ has a projected FD's set $S_1 := \{C \rightarrow D, D \rightarrow A\}$. So $D^+ = DA \neq R_1$, we decompose it again

$$R_1 = R_3 \bowtie R_4$$

$R_3 = DA$ and $R_4 = D \cup (R_1 \setminus DA) = DC$. All three of them have two attributes so they are in BCNF.

We conclude that the BCNF decomposition of R is

$$R_2(C, B), R_3(D, A), R_4(D, C).$$

- (b) Doing the closure of B we see that $B^+ = BCD \neq R$, so $B \rightarrow C$ violates BCNF. We decompose R into $R_1 = B^+ = BCD$ and $R_2 = B \cup (R \setminus B^+) = B \cup (ABCD \setminus BCD) = AB$. Their respective set of FD's $S_1 = \{B \rightarrow C, B \rightarrow D\}$ and $S_2 = \emptyset$ (where \emptyset denotes the absence of nontrivial FD's).

We conclude that R_1 is in BCNF since $B^+ = BCD = R_1$ and R_2 is also in BCNF since it holds only trivial FD's, so the decomposition is

$$R_1(B, C, D), R_2(A, B).$$

- (c) We see that $(AB)^+ = ABCD$, $(BC)^+ = BCDA$, $(CD)^+ = CDAB$, $(AD)^+ = AD BC$, so it already is in BCNF.
- (d) We see that $A^+ = ABCD$, $B^+ = BCDA$, $C^+ = CDAB$ and $D^+ = DABC$, so it already is in BCNF.
- (e) We see that $(AB)^+ = ABCD \neq ABCDE$, so we decompose R into $R_1 = ABCD$ and $R_2 = (AB) \cup (R \setminus (AB)^+) = (AB) \cup (E) = ABE$. Their respective set of FD's are $S_1 = \{AB \rightarrow C, B \rightarrow D\}$ and $S_2 = \emptyset$.

R_1 is not in BCNF since $B^+ = BD \neq ABCD$. We decompose R_1 into $R_3 = BD$ and $R_4 = B \cup (ABCD \setminus BD) = ABC$. R_3 is in BCNF,

and R_4 has a FD's set $\{AB \rightarrow C\}$, so it is in BCNF. We conclude that the BCNF decomposition is

$$R_2(A, B, E), R_3(B, D), R_4(A, B, C).$$

- (f) We see that $C \rightarrow D$ violates BCNF since $C^+ = CDBE \neq ABCDE$. We decompose R into $R_1 = BCDE$ and $R_2 = C \cup (ABCDE \setminus BCDE) = AC$. R_2 is in BCNF as it has only two attributes. R_1 has FD's set $\{C \rightarrow D, D \rightarrow B, D \rightarrow E\}$, suffices taking $D^+ = DBE$ to see that it is not yet in BCNF. We decompose R_1 into $R_3 = DBE$ and $R_4 = D \cup (BCDE \setminus DBE) = D \cup (C) = CD$. We see that the FD's set of R_3 is $\{D \rightarrow B, D \rightarrow E\}$ so it is in BCNF. We conclude that the BCNF decomposition is

$$R_2(A, C), R_3(D, B, E), R_4(C, D).$$

□

2. **3.3.2:** Consider a relation R whose schema is the set of attributes $\{A, B, C, D\}$ with FD's $A \rightarrow B$ and $A \rightarrow C$. Either is a BCNF violation, because the only key for R is $\{A, D\}$. Suppose we begin by decomposing R according to $A \rightarrow B$. Do we ultimately get the same result as if we first expand the BCNF violation to $A \rightarrow BC$? Why or why not?

Solution.

- (a) We start with a FD's set $S = \{A \rightarrow B, A \rightarrow C\}$; decomposing R with $A \rightarrow B$ we get $R_1 = A^+ = ABC$ and $R_2 = A \cup (R \setminus ABC) = AD$. With FD's set $S_1 = \{A \rightarrow B, A \rightarrow C\}$ and $S_2 = \emptyset$.
- (b) We decompose with $A \rightarrow BC$, $A^+ = ABC$; so $R_1 = ABC$ and $R_2 = AD$. With FD's set $S_1 = \{A \rightarrow BC\}$ and $S_2 = \emptyset$.

We conclude that the decompositions yield the same subrelations, but the FD's set for the second case is smaller (but equivalent under splitting).

□

3. **3.3.3:** Let R be as in Exercise 3.3.2, but let the FD's be $A \rightarrow B$ and $B \rightarrow C$. Again compare decomposing using $A \rightarrow B$ first against decomposing by $A \rightarrow BC$ first.

Solution.

- (a) Decomposing with $A \rightarrow B$ and seeing $A^+ = ABC$ gives us subrelations $R_1 = ABC$ and $R_2 = AD$. R_2 is in BCNF. The FD's set for R_1 is $S_1 = \{A \rightarrow B, B \rightarrow C\}$, so $B^+ = BC \neq ABC$. We decompose R_1 in

$$R_3 = BC \text{ and } R_4 = B \cup (R_1 \setminus BC) = AB.$$

We conclude that the BCNF decomposition for this case is

$$R_2(A, D), R_3(B, C), R_4(A, B).$$

- (b) Decomposing with $A \rightarrow BC$ we get $R_1 = ABC$ and $R_2 = AD$ with $S_1 = \{A \rightarrow BC\}$

4. **3.3.4:** Suppose we have a relation schema $R(A, B, C)$ with FD $A \rightarrow B$. Suppose also that we decide to decompose this schema into $S(A, B)$ and $T(B, C)$. Give an example of an instance of relation R whose projection onto S and T and subsequent rejoining does not yield the same relation instance. That is, $\pi_{A,B}(R) \bowtie \pi_{B,C}(R) \neq R$.

Solution.

5. Let \mathcal{F} and \mathcal{F}_{min} be two equivalent sets of FD's for a relation R , where
- (a) \mathcal{F} contains FD's with arbitrary right-hand sides (e.g., $X \rightarrow Y$ where $|Y| \geq 1$),
 - (b) \mathcal{F}_{min} is a minimal basis.

Prove or disprove the following claim:

Claim:

Decomposing R into BCNF using \mathcal{F}_{min} results in a number of subrelations that is *less than or equal to* the number of subrelations obtained using \mathcal{F} instead.

Solution.

6. Prove that the BCNF decomposition algorithm ends after finitely many times. Furthermore prove that each output subrelation is in BCNF.

Remark. BCNF decomposition's algorithm, in J. Ullman's book, is stated as following:

INPUT: A relation R_0 with a set of functional dependencies S_0 .

OUTPUT: A decomposition of R_0 into a collection of relations, all of which are in BCNF.

METHOD: The following steps can be applied recursively to any relation R and set of FD's S . Initially, apply them with $R = R_0$ and $S = S_0$.

- (a) Check whether R is in BCNF. If so, nothing more needs to be done. Return $\{R\}$ as the answer.
- (b) If there are BCNF violations, let one be $X \rightarrow Y$. Use Algorithm 3.7 to compute X^+ . Choose $R_1 = X^+$ as one relation schema and let R_2 have attributes X and those attributes of R that are not in X^+ .
- (c) Use Algorithm 3.12 to compute the sets of FD's for R_1 and R_2 ; let these be S_1 and S_2 , respectively.
- (d) Recursively decompose R_1 and R_2 using this algorithm. Return the union of the results of these decompositions.

Solution.

- (a) (**BCNF algorithm ends**) Let $R_0(A_1, \dots, A_n)$ be any relation, and $S_0 = \{X_1 \rightarrow Y_1, \dots, X_m \rightarrow Y_m\}$ be a set of functional dependencies on R_0 with Y_i any subset of attributes in R_0 .

By induction on the amount n of attributes of R_0 .

Basis If $n = 1$, then S_0 yields only the trivial dependency $A_i \rightarrow A_i$. So it is in BCNF, hence suffices applying BCNF algorithm 0 times.

If $n = 2$ then as we used in the earlier exercises, R_0 is in BCNF (proof of this is skipped, as it is in the book), hence suffices applying BCNF algorithm 0 times.

Hypothesis We claim that there exists an integer $k \geq 0$ such that after applying BCNF algorithm k times onto a relation R_0 with n attributes results in each subrelation $R_i \subset R_0$ having 1 or 2 attributes or satisfying BCNF.

Thesis We see that for a relation R_i with n attributes, after a first decomposition becomes:

$$R_i := \begin{cases} R_{i+1} = X^+ \\ R_{i+2} = X \cup (R \setminus X^+) \end{cases} \quad .$$

For R_{i+1} we see that $|X^+| < n$ since X is not a superkey. For R_{i+2} we see that $|X^+| > |X| \geq 1$ since there exists Y such that $X \rightarrow Y$ is nontrivial. Hence

$$|X \cup (R \setminus X^+)| \leq |X| + |R \setminus X^+| < |X| + |R \setminus X| = n.$$

We conclude that applying BCNF algorithm results into a shrinkage of the input relation into two smaller relations for any $i \geq 0$.

Let k' the amount of times to apply BCNF algorithm until each subrelation is in BCNF. We suppose R_0 has $n + 1$ attributes, then one can apply BCNF algorithm and see that the outputted relations R_1, R_2 have at most n elements. So by hypothesis, these both can be decomposed into BCNF subrelations with k applications. We conclude that suffices letting $k' = k + 1$.

- (b) (**BCNF algorithm yields BCNF subrelations**) Since BCNF algorithm is applicable over non-BCNF relations, if it ends (which does), then each subrelation is in BCNF.

□

Dependency Preservation and 3NF

1. **3.4.1** Let $R(A, B, C, D, E)$ be decomposed into relations with the following three sets of attributes: $\{A, B, C\}$, $\{B, C, D\}$, and $\{A, C, E\}$. For each of the following sets of FD's, use the chase test to tell whether the decomposition of R is lossless. For those that are not lossless, give an example of an instance of R that returns more than R when projected onto the decomposed relations and rejoined.
 - (a) $B \rightarrow E$ and $CE \rightarrow A$.
 - (b) $AC \rightarrow E$ and $BC \rightarrow D$.
 - (c) $A \rightarrow D$, $D \rightarrow E$, and $B \rightarrow D$.
 - (d) $A \rightarrow D$, $CD \rightarrow E$, and $E \rightarrow D$.

Solution.

2. **3.4.2** For each of the sets of FD's in Exercise 3.4.1, are dependencies preserved by the decomposition?
3. **3.5.1** For each of the relation schemas and sets of FD's of Exercise 3.3.1:
 - (a) Indicate all the 3NF violations.
 - (b) Decompose the relations, as necessary, into collections of relations that are in 3NF.
4. **Exercise 3.5.2** Consider the relation **Courses**(C, T, H, R, S, G), whose attributes may be thought of informally as course, teacher, hour, room, student, and grade. Let the set of FD's for **Courses** be $C \rightarrow T$, $HR \rightarrow C$, $HT \rightarrow R$, $HS \rightarrow R$, and $CS \rightarrow G$. Intuitively, the first says that a course has a unique teacher, and the second says that only one course can meet in a given room at a given hour. The third says that a teacher can be in only one room at a given hour, and the fourth says the same about students. The last says that students get only one grade in a course.
 - (a) What are all the keys for **Courses**?
 - (b) Verify that the given FD's are their own minimal basis.
 - (c) Use the 3NF synthesis algorithm to find a lossless-join, dependency-preserving decomposition of R into 3NF relations. Are any of the relations not in BCNF?
5. **Exercise 3.5.3** Consider a relation **Stocks**(B, O, I, S, Q, D), whose attributes may be thought of informally as broker, office (of the broker), investor, stock, quantity (of the stock owned by the investor), and dividend (of the stock). Let the set of FD's for **Stocks** be $S \rightarrow D$, $I \rightarrow B$, $IS \rightarrow Q$, and $B \rightarrow O$. Repeat Exercise 3.5.2 for the relation **Stocks**.

6. **Exercise 3.5.4** Verify, using the chase, that the decomposition of Example 3.27 has a lossless join.
7. **Exercise 3.5.5** Suppose we modified Algorithm 3.20 (BCNF decomposition) so that instead of decomposing a relation R whenever R was not in BCNF, we only decomposed R if it was not in 3NF. Provide a counterexample to show that this modified algorithm would not necessarily produce a 3NF decomposition with dependency preservation.

Solution.