

Convergence Sheet II

Gallo Tennis / to A Mathematical Room

April 25, 2025

Problems on convergence of sequences, functions and continuity. The problems are from the books of M. Spivak "Calculus", T. Tao "Analysis I", Charles C. Pugh "Real Mathematical Analysis", Vladimir A. Zorich "Mathematical Analysis I" and Titu Andreescu "Putnam and Beyond".

Suprema and Infima of sequences

1. Let E be a subset of $\mathbb{R}^*$. Then the following statements are true.
 - (a) For every $x \in E$ we have $x \leq \sup(E)$ and $x \geq \inf(E)$.
 - (b) Suppose that $M \in \mathbb{R}^*$ is an upper bound for E , i.e., $x \leq M$ for all $x \in E$. Then we have $\sup(E) \leq M$.
 - (c) Suppose that $M \in \mathbb{R}^*$ is a lower bound for E , i.e., $x \geq M$ for all $x \in E$. Then we have $\inf(E) \geq M$.

Solution.

2. **(Least upper bound property).** Let $(a_n)_{n=m}^{\infty}$ be a sequence of real numbers, and let x be the extended real number

$$x := \sup(a_n)_{n=m}^{\infty}.$$

Then we have $a_n \leq x$ for all $n \geq m$. Also, whenever $M \in \mathbb{R}^*$ is an upper bound for a_n (i.e., $a_n \leq M$ for all $n \geq m$), we have $x \leq M$.

Finally, for every extended real number y for which $y < x$, there exists at least one $n \geq m$ for which

$$y < a_n \leq x.$$

Solution. The first two propositions follow directly from the definitions of suprema and infima of sequences and sets of extended real numbers.

For the final proposition, if $x = +\infty$, and $y \in \mathbb{R}$ then given that $a_n \in \mathbb{R}$, (a_n) is divergent. Therefore, for any K there exists some $n \geq m$ such that $a_n \geq K$. In particular there exists some $n_0 \geq m$ such that $a_{n_0} \geq K_0 > y$.

If $x \in \mathbb{R}$ then $y \in \mathbb{R}$ for any $y < x$. Suppose for the sake of contradiction that there existed at least one $y_0 < x$ such that for any $n \geq m$, $a_n \leq y_0$.

Then immediately we get a contradiction, as y_0 is an upper bound smaller than x .

Note that $x, y = \infty$ but $y < x$ would mean that $\infty < \infty$ which is false.

□

3. (**Monotone bounded sequences converge**). Let $(a_n)_{n=m}^{\infty}$ be a sequence of real numbers which has some finite upper bound $M \in \mathbb{R}$, and which is also increasing (i.e., $a_{n+1} \geq a_n$ for all $n \geq m$). Then $(a_n)_{n=m}^{\infty}$ is convergent, and in fact

$$\lim_{n \rightarrow \infty} a_n = \sup(a_n)_{n=m}^{\infty} \leq M.$$

Proof. Since (a_n) is bounded, its supremum cannot be infinite. Let ε_n be the real number $\sup(a_n) - a_n$. We see that (ε_n) is a decreasing positive sequence, has supremum $\sup(a_n) - a_1$. It is verified that $\inf(\varepsilon_n) = 0$ since otherwise, if $\inf(\varepsilon_n) = c > 0$ then for any $n \geq 1$, by problem (2)

$$c = \inf(\varepsilon_n) = \inf(\sup(a_n) - a_n) \leq \sup(a_n) - a_n$$

so that

$$a_n \leq \sup(a_n) - c < \sup(a_n)$$

which contradicts the fact that $\sup(a_n)$ is the least upper bound of (a_n) .

We claim that given any positive integer $k \geq 1$ there exists some K such that for any $n, m \geq K$,

$$|a_n - a_m| \leq \varepsilon_k.$$

To prove this claim we see that for $n, m \geq k$ by symmetry if $n \geq m$, since (a_n) is monotone

$$|a_n - a_m| = a_n - a_m \leq \sup(a_n) - a_n \leq \sup(a_n) - a_k = \varepsilon_k.$$

We conclude that suffices letting $K := k$.

Then, let $\varepsilon > 0$ be any real number, we claim that there exists some $N \geq 1$ such that for any $n \geq N$

$$|a_n - \sup(a_n)| \leq \varepsilon.$$

To prove this claim we see that since $\inf(\varepsilon_n) = 0$, by problem (2), for any $\varepsilon > 0$ we can find at least one k such that $\varepsilon > \varepsilon_k \geq 0$, hence this shows that

$$|a_n - a_m| \leq \varepsilon$$

for a given ε and for every $n, m \geq K = k$, so the sequence converges by Cauchy criterion. Then, by construction

$$|\sup(a_n) - a_n| = \varepsilon_n \leq \varepsilon.$$

□

4. Explain why Proposition 6.3.10 fails when $x > 1$. In fact, show that the sequence $(x^n)_{n=1}^\infty$ diverges when $x > 1$. (Hint: prove by contradiction and use the identity $(1/x)^n x^n = 1$ and the limit laws in Theorem 6.1.19.) Compare this with the argument in Example 1.2.3; can you now explain the flaws in the reasoning in that example?

Remark. Proposition 6.3.10. Let $0 < x < 1$. Then we have

$$\lim_{n \rightarrow \infty} x^n = 0.$$

Solution. Let $x > 1$. Suppose for the sake of contradiction that $\lim_{n \rightarrow \infty} x^n$ converged to $L \in \mathbb{R}$. Consider the sequence $(\frac{1}{x})^n$, this sequence converges to 0 by squeeze theorem with $\frac{1}{n}$. Therefore, by limit laws (since both sequences supposedly converge) we should get that

$$\lim_{n \rightarrow \infty} \left(\frac{1}{x}\right)^n \cdot x^n = \lim_{n \rightarrow \infty} \left(\frac{1}{x}\right)^n \lim_{n \rightarrow \infty} x^n = 0 \cdot L = 0.$$

In spite of this, the limit should be equal to

$$\lim_{n \rightarrow \infty} \left(\frac{1}{x^n}\right) \cdot x^n = \lim_{n \rightarrow \infty} \frac{x^n}{x^n} = 1.$$

We conclude that $0 = 1$ which is false. We conclude that x^n does not converge if $x > 1$, note that in this case the test of falseness we used was not the epsilon definition but the limit laws theorem.

Regarding Example 1.2.3, that essentially says that for any $x \in \mathbb{R}$

$$L = \lim_{n \rightarrow \infty} x^n = \lim_{m+1 \rightarrow \infty} x^{m+1} = x \lim_{m+1 \rightarrow \infty} x^m,$$

but by limit laws:

$$x \lim_{m+1 \rightarrow \infty} x^m = xL,$$

concluding that

$$xL = L.$$

So that $L = 0$ or $x = 1$. the two main flaws are: first, assuming that the sequence converges to a finite limit L for any x (that we have shown it does not) and the second (related to the first), that we can apply limit laws in possibly divergent sequences. The condition to the limit laws theorem to hold is the sequences to converge in first place.

□

5. **(Limits are limit points.)** Let $(a_n)_{n=m}^\infty$ be a sequence which converges to a real number c . Then c is a limit point of $(a_n)_{n=m}^\infty$, and in fact it is the only limit point of $(a_n)_{n=m}^\infty$.

Proof.

Existence If (a_n) converges to c then for any $\varepsilon > 0$ there exists an $N \geq m$ such that for any $n \geq N$

$$|a_n - c| \leq \varepsilon.$$

In order for c to be a limit point, it must happen that for any $\varepsilon > 0$ and any $N' \geq m$ there exists an $n' \geq N'$ such that

$$|a_{n'} - c| \leq \varepsilon.$$

Let $n' = \max\{N, N'\}$ (fixed N and variable N') then in particular for any $n \geq n' \geq N$

$$|a_n - c| \leq \varepsilon \text{ } ((a_n) \text{ converges})$$

and in particular since $n' \geq N'$ and by convergence

$$|a_{n'} - c| \leq \varepsilon \text{ } (c \text{ is a limit point}).$$

Uniqueness Note that limit are limit points does not guarantee that limit points are limits. Let d be a limit point then for any given $\varepsilon_0 > 0$ and any $N' \geq m$ there exists an $n' \geq N'$ such that

$$|a_{n'} - d| \leq \varepsilon_0.$$

in particular, since a_n converges to c , for ε_0 there exist a $N \geq m$ such that for any $n \geq N$

$$|a_n - c| \leq \varepsilon_0.$$

Choose $N' := N$ (in the limit point condition) so that for any $n \geq N'$ (by convergence condition) and in particular for n' (by limit point condition)

$$|a_{n'} - d| \leq \varepsilon_0 \text{ and } |a_{n'} - c| \leq \varepsilon_0.$$

We the notice that by triangle inequality

$$|c - d| \leq |c - a_{n'}| + |a_{n'} - d| \leq \varepsilon_0.$$

Since we let $\varepsilon_0 > 0$ be arbitrary in the condition of d to be a limit point, we conclude that $c = d$.

□

6. Let $(a_n)_{n=m}^{\infty}$ be a sequence of real numbers, let L^+ be the limit superior of this sequence, and let L^- be the limit inferior of this sequence (thus both L^+ and L^- are extended real numbers).

(c) We have

$$\inf(a_n)_{n=m}^{\infty} \leq L^- \leq L^+ \leq \sup(a_n)_{n=m}^{\infty}.$$

(d) If c is any limit point of $(a_n)_{n=m}^\infty$, then we have

$$L^- \leq c \leq L^+.$$

- (e) If L^+ is finite, then it is a limit point of $(a_n)_{n=m}^\infty$. Similarly, if L^- is finite, then it is a limit point of $(a_n)_{n=m}^\infty$.
- (f) Let c be a real number. If $(a_n)_{n=m}^\infty$ converges to c , then we must have $L^+ = L^- = c$. Conversely, if $L^+ = L^- = c$, then $(a_n)_{n=m}^\infty$ converges to c .

Solution.

(c) We have the trivial inequality

$$a_m^- = \inf(a_n)_{n=m}^\infty \leq a_m \leq \sup(a_n)_{n=m}^\infty = a_m^+.$$

Claim. We claim that for any $m_0, m_1 \geq m$

$$a_{m_0}^+ \geq a_{m_1}^-.$$

Proof. To prove this claim we see that if $m_0 \leq m_1$, then

$$\sup(a_n)_{n=m_0}^\infty \geq \sup(a_n)_{n=m_1}^\infty \geq \inf(a_n)_{n=m_1}^\infty$$

so it holds. If $m_0 \geq m_1$ then

$$\inf(a_n)_{n=m_1}^\infty \leq \inf(a_n)_{n=m_0}^\infty \leq \sup(a_n)_{n=m_0}^\infty$$

so it also holds in this case. ■

Suppose for the sake of contradiction that $L^- = \sup(a_n^-)_{n=m}^\infty > \inf(a_n^+)_{n=m}^\infty = L^+$, then by problem (2) there exists some $m_0 \geq m$ such that

$$L^+ \leq a_{m_0}^+ < L^-$$

hence $a_{m_0}^+ < L^-$, then again, by problem (2) we must have that

$$a_{m_0}^+ < a_{m_1}^- \leq L^- \text{ for some } m_1 \geq m$$

contradicting the first claim. We conclude that $L^+ \geq L^-$.

- (d) One may think at first that for any $n \geq m$ we have $L^- \leq a_n \leq L^+$, this is false, and a counterexample is the sequence $(a_n)_{n=1}^\infty$ whose terms $a_1, a_2, a_3, \dots$ are

$$1, -1/2, 1/3, \dots$$

then the sequence a_n^+ is

$$1, 1/3, 1/5, \dots$$

whose infimum $\inf(a_n^+)_{n=1}^\infty$ is just 0, but we see that $a_1 > 0$. Similarly for other case $\sup(a_n^-)_{n=1}^\infty = 0$ but we see that $a_2 < 0$.

With that precision done, suppose for the sake of contradiction that $c > L^+$ (by symmetry $c < L^-$), then we must have that for an $n_0 \geq m$, $c > a_{n_0}^+ \geq L^+$.

Since c is a limit point, for any $\varepsilon > 0$ and any $N \geq m$ there exists some $n \geq N$ such that

$$|a_n - c| \leq \varepsilon.$$

In particular let $N = n_0$ and let $\varepsilon := \frac{c - a_{n_0}^+}{2}$, therefore

$$a_n \geq \frac{a_{n_0}^+ - c}{2} + c = \frac{a_{n_0}^+ + c}{2} > a_{n_0}^+.$$

Therefore, $a_n > a_{n_0}^+$, but since $n \geq n_0$ ($a_{n_0}^+$ “reaches” a_n) we also have $a_{n_0}^+ \geq a_n$, a contradiction.

- (e) By symmetry, we have to verify that for any $\varepsilon > 0$ and any $N \geq m$ we can find some $n \geq N$ such that

$$|a_n - L^+| \leq \varepsilon.$$

So that

$$L^+ - \varepsilon \leq a_n \leq \varepsilon + L^+.$$

We notice that $(a_m^+)_{n=m}^\infty$ is decreasing (we have used this fact in item (1), for example), since $a_m^+ := \sup\{a_n : n \geq m\} \geq a_{m+1}^+ := \sup\{a_n : n \geq m+1\}$ but it is also bounded below by the finite real number $\inf(a_n^+)$, hence by problem (3) this sequence converges to $\inf(a_n^+)$.

Therefore, for any $\varepsilon > 0$ there exists some $N \geq m$ such that for any $n \geq N$

$$|a_n^+ - L^+| < \varepsilon$$

hence

$$L^+ - \varepsilon < a_n^+ < \varepsilon + L^+$$

by problem (2) we also know that there must exist some $n_0 \geq m$ such that

$$L^+ - \varepsilon < a_n \leq a_n^+ < \varepsilon + L^+.$$

We conclude that for any $n \geq N$

$$|a_n - L^+| < \varepsilon$$

since limits are limit points (problem (5)), we conclude.

- (f) ($\implies$) If (a_n) converges to c then c is a limit point and the sequence is bounded, therefore L^+ and L^- are finite. So by item (d)

$$L^- \leq c \leq L^+.$$

Suppose that

$$L^- < c < L^+.$$

By definition of convergence, for any $\varepsilon > 0$ there exists some $N \geq m$ such that for any $n \geq N$

$$|a_n - c| \leq \varepsilon$$

and in particular we can set $\varepsilon := L^+ - c$, hence we can find some N_0 such that for any $n \geq N_0$

□

7. **(Comparison principle.)** Suppose that $(a_n)_{n=m}^\infty$ and $(b_n)_{n=m}^\infty$ are two sequences of real numbers such that $a_n \leq b_n$ for all $n \geq m$. Then we have the inequalities:

$$\begin{aligned} \sup(a_n)_{n=m}^\infty &\leq \sup(b_n)_{n=m}^\infty \\ \inf(a_n)_{n=m}^\infty &\leq \inf(b_n)_{n=m}^\infty \\ \limsup_{n \rightarrow \infty} a_n &\leq \limsup_{n \rightarrow \infty} b_n \\ \liminf_{n \rightarrow \infty} a_n &\leq \liminf_{n \rightarrow \infty} b_n \end{aligned}$$

Solution.

- (a) Suppose for the sake of contradiction that $\sup(a_n) > \sup(b_n)$, then $\sup(a_n) > b_n$ for any $n \geq m$. Hence $\sup(a_n)$ is an upper bound for b_n for any $n \geq m$. By hypothesis, since $b_n \geq a_n$ then

$$\sup(a_n) > \sup(b_n) \geq b_n \geq a_n$$

so $\sup(b_n)$ is a smaller upper bound than $\sup(a_n)$, a contradiction.

- (b) The case for infimum is analogous to the one before.
(c) Suppose for the sake of contradiction that

$$\limsup_{n \rightarrow \infty} a_n > \limsup_{n \rightarrow \infty} b_n$$

hence, $L_a^+ > L_b^+$, then for any $N \geq m$ it holds that

$$a_n > L_b^+$$

for some $n \geq N$. This implies that fixing $n_0 \geq N$ there must also exist some $N' \geq m$ such that for any $n \geq N'$

$$b_n < a_{n_0}.$$

Then, we let $N = N'$ so that we can fix any $a_n \geq N'$ so in particular we must have

$$b_{n_0} < a_{n_0}.$$

A contradiction, since $a_n \leq b_n$ for any $n \geq m$.

(d) The case for the infimum is analogous to the one before.

□

8. **(Squeeze test.)** Let $(a_n)_{n=m}^{\infty}$, $(b_n)_{n=m}^{\infty}$, and $(c_n)_{n=m}^{\infty}$ be sequences of real numbers such that

$$a_n \leq b_n \leq c_n$$

for all $n \geq m$. Suppose also that $(a_n)_{n=m}^{\infty}$ and $(c_n)_{n=m}^{\infty}$ both converge to the same limit L . Then $(b_n)_{n=m}^{\infty}$ is also convergent to L .

Solution. Since (a_n) and (c_n) both converge to L , then by problem (6f) we must have

$$L_a^+ = L_c^+ = L = L_c^- = L_a^-.$$

And all of them are finite. By problem (7) we also know that since $a_n \leq b_n \leq c_n$

$$\begin{aligned} \limsup_{n \rightarrow \infty} a_n &\leq \limsup_{n \rightarrow \infty} b_n \leq \limsup_{n \rightarrow \infty} c_n \\ \liminf_{n \rightarrow \infty} a_n &\leq \liminf_{n \rightarrow \infty} b_n \leq \liminf_{n \rightarrow \infty} c_n. \end{aligned}$$

We conclude that $L_b^- = L_b^+ = L$.

□

9. **(Zero test for sequences.)** Let $(a_n)_{n=m}^{\infty}$ be a sequence of real numbers. Then the limit $\lim_{n \rightarrow \infty} a_n$ exists and is equal to zero if and only if the limit $\lim_{n \rightarrow \infty} |a_n|$ exists and is equal to zero.

Solution.

- ($\implies$) If a_n converges to 0, since $|a_n| \geq a_n$ for any $n \geq 1$ we expect that if the limit exists, then $\lim_{n \rightarrow \infty} |a_n| \geq 0$. To prove that the limit exists, we note that since (a_n) converges to zero, then for any $\varepsilon > 0$ there must exist some $N \geq m$ such that for any $n \geq N$

$$|a_n| \leq \varepsilon$$

we see that this immediately tells us that

$$||a_n|| \leq \varepsilon.$$

Thus, in fact $(|a_n|)$ does converge, and it converges to zero.

- ($\impliedby$) If $\lim_{n \rightarrow \infty} |a_n|$ exists and it is equal to zero, similarly, for any $\varepsilon > 0$ there must exist some $N \geq m$ such that if $n \geq N$ then

$$||a_n|| \leq \varepsilon.$$

Analogously to the case before, since $|a_n| \geq 0$, then $||a_n|| = |a_n|$, thus

$$|a_n| \leq \varepsilon.$$

□

10. Let us say that a sequence $(a_n)_{n=M}^{\infty}$ of real numbers has $+\infty$ as a limit point iff it has no finite upper bound, and that it has $-\infty$ as a limit point iff it has no finite lower bound. With this definition, show that

$$\limsup_{n \rightarrow \infty} a_n$$

is a limit point of $(a_n)_{n=M}^{\infty}$, and furthermore that it is larger than all the other limit points of $(a_n)_{n=M}^{\infty}$; in other words, the limit superior is the largest limit point of a sequence. Similarly, show that the limit inferior is the smallest limit point of a sequence. (One can use Proposition 6.4.12 in the course of the proof.)

11. Let $(a_n)_{n=N}^{\infty}$ be a sequence of real numbers, and let $(b_m)_{m=M}^{\infty}$ be another sequence of real numbers such that each b_m is a limit point of $(a_n)_{n=N}^{\infty}$. Let c be a limit point of $(b_m)_{m=M}^{\infty}$. Prove that c is also a limit point of $(a_n)_{n=N}^{\infty}$. (In other words, limit points of limit points are themselves limit points of the original sequence.)
12. Let x be a real number. Then the limit $\lim_{n \rightarrow \infty} x^n$ exists and is equal to zero when $|x| < 1$, exists and is equal to 1 when $x = 1$, and diverges when $x = -1$ or when $|x| > 1$.
13. For any $x > 0$, we have $\lim_{n \rightarrow \infty} x^{1/n} = 1$.
14. If a and b are positive numbers and p an arbitrary nonzero real number, then the mean of order p of the numbers a and b is the quantity

$$S_p(a, b) = \left(\frac{a^p + b^p}{2} \right)^{\frac{1}{p}}.$$

In particular for $p = 1$ we obtain the arithmetic mean of a and b , for $p = 2$ their square-mean, and for $p = -1$ their harmonic mean.

- (a) Show that the mean $S_p(a, b)$ of any order lies between the numbers a and b .
- (b) Find the limits of the sequences

$$\{S_n(a, b)\}, \quad \{S_{-n}(a, b)\}.$$

15. Show that if $a > 0$, the sequence $x_{n+1} = \frac{1}{2} \left(x_n + \frac{a}{x_n} \right)$ converges to the square root of a for any $x_1 > 0$. Estimate the rate of convergence, that is, the magnitude of the absolute error $|x_n - \sqrt{a}| = |\Delta_n|$ as a function of n .
Solution. Notice that since $x_n, a > 0$, by *AM – GM* inequality

$$\frac{1}{2} \left(x_n + \frac{a}{x_n} \right) \geq \sqrt{x_n \cdot \frac{a}{x_n}} = \sqrt{a}.$$

In fact for any $n \geq m$ we have the above relation, and thus we can conclude (x_n) is bounded below by $\sqrt{a}$.

We claim that (x_n) is a decreasing sequence. To prove this claim, consider the difference sequence $x_{n+1} - x_n$. We want to show that $x_{n+1} - x_n \leq 0$. We observe that

$$x_{n+1} - x_n = \left(x_n - \frac{a}{x_n}\right) - x_n = \frac{a - x_n^2}{2x_n} = \frac{(\sqrt{a} - x_n)(\sqrt{a} + x_n)}{2x_n}.$$

Since $\sqrt{a} \leq x_n$, then $\sqrt{a} - x_n \leq 0$ so the whole ratio becomes negative or zero. This implies that $x_{n+1} \leq x_n$.

Since (x_n) is a decreasing sequence and it is bounded below by $\sqrt{a}$, it converges. The only thing that remains checking is $\sqrt{a} = \inf(x_n)$. Suppose for the sake of contradiction that there existed another infimum $L > \sqrt{a}$. Thus, for any $n \geq 1$

$$x_n \geq L$$

since x_n converges, for any $\varepsilon > 0$ we can find some $N \geq m$ such that for any $n \geq N$

$$x_n - L \leq \varepsilon.$$

Let $\varepsilon := \frac{L - \sqrt{a}}{2}$, then

$$x_n - L \leq \frac{L - \sqrt{a}}{2} \implies x_n \leq \frac{L + \sqrt{a}}{2} \leq \frac{x_n + \sqrt{a}}{2}$$

but then

$$2x_n \leq x_n + \sqrt{a} < x_n + L \leq 2x_n$$

which is a contradiction ($2x_n < 2x_n$). We conclude that $\sqrt{a} = \inf(x_n)$, and thus is the limit of the sequence.

To find the magnitude of the absolute error, since $x_n - \sqrt{a} = \Delta_n$, then $x_n = \Delta_n + \sqrt{a}$. This implies that

$$x_{n+1} = \Delta_{n+1} + \sqrt{a} = \frac{1}{2} \left((\Delta_n + \sqrt{a}) + \frac{a}{\Delta_n + \sqrt{a}} \right)$$

thus

$$\Delta_{n+1} = \frac{1}{2} \left((\Delta_n + \sqrt{a}) + \frac{a}{\Delta_n + \sqrt{a}} \right) - \sqrt{a}$$

operating the right hand side, we get

$$\Delta_{n+1} = \frac{1}{2} \left(\frac{\Delta_n^2 + 2\Delta_n\sqrt{a} + 2a}{\Delta_n + \sqrt{a}} \right) - \sqrt{a}$$

therefore

$$\Delta_{n+1} = \frac{\Delta_n^2 + 2\Delta_n\sqrt{a} + 2a - 2\sqrt{a}\Delta_n - 2a}{2(\Delta_n + \sqrt{a})} = \frac{\Delta_n^2}{2(\Delta_n + \sqrt{a})}$$

we conclude that

$$\Delta_{n+1} = \frac{\Delta_n^2}{2x_n}.$$

This implies that the error for any subsequent term is quadratic, although, since x_n converges to $\sqrt{a}$, $(x_n - \sqrt{a})^2 < 1$ for some $n \geq N'$, and from there on, the error will decrease “faster”.

□

16. We mark all the points on a circle obtained from a fixed point by rotations of the circle through angles of n radians, where $n \in \mathbb{Z}$ ranges over all integers. Describe all the limit points of the set so constructed.

Solution. Note that we cannot have directly a full circumference rotation, since $\pi \notin \mathbb{Z}$. We would like to check if eventually, for a fixed point, it is achievable that it returns to its same origin.

Consider the unitary circumference C . Let x_n be any point in C that is at an angle of n radians with respect to $(1, 0)$.

Subsequences

1. Let $(a_n)_{n=0}^\infty$, $(b_n)_{n=0}^\infty$, and $(c_n)_{n=0}^\infty$ be sequences of real numbers. Then $(b_n)_{n=0}^\infty$ is a subsequence of $(a_n)_{n=0}^\infty$. Furthermore, if $(b_n)_{n=0}^\infty$ is a subsequence of $(a_n)_{n=0}^\infty$, and $(c_n)_{n=0}^\infty$ is a subsequence of $(b_n)_{n=0}^\infty$, then $(c_n)_{n=0}^\infty$ is a subsequence of $(a_n)_{n=0}^\infty$.

Solution.

- (a) To show (b_n) is a subsequence of (a_n) suffices letting $f(n) = n$.
- (b) Since (b_n) is a subsequence of (a_n) there exists an increasing function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $b_n = a_{f(n)}$. Since (c_n) is a subsequence of (b_n) there exists an increasing function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $c_n = b_{g(n)}$. Therefore, $c_n = b_{g(n)} = a_{(g \circ f)(n)}$. The only thing that remains checking is that $g \circ f$ is an increasing function. To verify this we note that since $f(n) \geq f(n+1)$ and $g(n) \geq g(n+1)$, then

$$(g \circ f)(n+1) = g(f(n+1)) > g(f(n)) = (g \circ f)(n).$$

□

2. Let $(a_n)_{n=0}^\infty$ be a sequence of real numbers, and let L be a real number. Then the following two statements are logically equivalent (each one implies the other):

- (a) The sequence $(a_n)_{n=0}^\infty$ converges to L .
- (b) Every subsequence of $(a_n)_{n=0}^\infty$ converges to L .

Solution.

(a) $\implies$ (b) The following claim will result handfull.

Claim. For any function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $f(n+1) > f(n)$, it holds that for any $n > 0$

$$f(n) \geq n.$$

Proof. By induction on n ,

Basis If $n = 0$, then $f(0)$ is or zero or a positive integer, either case suffices.

Hypothesis Assume that for any n it holds that $f(n) \geq n$.

Thesis We want to prove that for the positive integer $n+1$ it holds that $f(n+1) \geq n+1$. By hypothesis $f(n) \geq n$, therefore since $f(n+1) > f(n)$ then $f(n+1) \geq f(n) + 1 \geq n+1$.

■

Since (a_n) converges to L then for any $\varepsilon > 0$ there exists some $N \geq 0$ such that if $n \geq N$

$$|a_n - L| \leq \varepsilon.$$

Using the claim we get that in particular for any $n \geq N$, $f(n) \geq N$ and thus

$$|a_{f(n)} - L| \leq \varepsilon.$$

(b) $\implies$ (a) If any subsequence of (a_n) converges to L , in particular the subsequence $(a_{f(n)})$ with $f(n) = n$ converges, this is, the original sequence.

□

3. Let $(a_n)_{n=0}^{\infty}$ be a sequence of real numbers, and let L be a real number. Then the following two statements are logically equivalent:

(a) L is a limit point of $(a_n)_{n=0}^{\infty}$.

(b) There exists a subsequence of $(a_n)_{n=0}^{\infty}$ which converges to L .

Solution.

4. Can you find two sequences $(a_n)_{n=0}^{\infty}$ and $(b_n)_{n=0}^{\infty}$ which are not the same sequence, but such that each is a subsequence of the other?

Solution. Consider the sequences (a_n) , (b_n) with $a_n = (-1)^n$ and $b_n = (-1)^{n+2}$. We see that $b_n = a_{f(n)}$ with $f(n) = n+2$ and $a_n = b_{g(n)}$ with $g(n) = n-2$.

□

5. Let $(a_n)_{n=0}^\infty$ be a sequence which is not bounded. Show that there exists a subsequence $(b_n)_{n=0}^\infty$ of $(a_n)_{n=0}^\infty$ such that $\lim_{n \rightarrow \infty} 1/b_n$ exists and is equal to zero.

(Hint: for each natural number j , recursively introduce the quantity

$$n_j := \min\{n \in \mathbb{N} : |a_n| \geq j; n > n_{j-1}\}$$

(omitting the condition $n > n_{j-1}$ when $j = 0$), first explaining why the set $\{n \in \mathbb{N} : |a_n| \geq j \text{ for } n > n_{j-1}\}$ is non-empty. Then set $b_j := a_{n_j}$.)

Solution. The intuition is that the hint's set construction is similar to "looking for an increasing sequence", for instance if we have the subsequence $(b_n) = n$, we can directly construct the sequence $\frac{1}{b_n}$ that converges to zero. For doing so, we start with some natural number j and a term a_j , and look the "behaviour" of the sequence from there on, and gather the following greater term.

Given any sequence (a_n) we want to prove that we can find an increasing monotone subsequence. By induction in j we claim that the set

$$A_j := \{n \in \mathbb{N} : |a_n| \geq j \text{ for some } n > f(j-1)\}$$

is nonempty for any $j \geq 1$. The weaker case in which $f(j) > f(j-1)$ will suffice.

Basis. Let $f(0) = 0$. If $j = 1$ then $A_1 = \{n \in \mathbb{N} : |a_n| \geq 1 \text{ for some } n > f(0)\}$. Since (a_n) is an unbounded sequence, there exists some $n \geq f(0)$ such that $|a_n| \geq 1$. We conclude that A_1 is not empty.

Hypothesis. Assume that for any $j \geq 1$ the set

$$A_j := \{n \in \mathbb{N} : |a_n| \geq j \text{ for some } n > f(j-1)\}$$

is nonempty.

Thesis. Consider the integer $j+1$, we want to prove that the set

$$A_{j+1} := \{n \in \mathbb{N} : |a_n| \geq j+1 \text{ for some } n > f(j)\}$$

is nonempty. By hypothesis the set A_j is nonempty, we have found "beforehand" some pivots $n' \in A_j$ such that $|a_{n'}| \geq j$ in which look from to.

We see that since (a_n) is unbounded there must exist some $n > n'$ such that $|a_n| \geq j+1$. The only thing that remains checking is that under the former condition there exists some $n > n'$ such that $n > f(j)$.

Suppose that for any $n \geq n'$, if $|a_n| \geq j+1$ then $n < f(j)$. Since by hypothesis we have $n > n' > f(j-1)$, then

$$f(j) > n > f(j-1).$$

But since (a_n) is not bounded, we can find some $n \geq f(j)$ such that

$$|a_n| \geq j + 1.$$

We conclude that for A_{j+1} is nonempty. ■

Claim. We claim that $\min(A_{j+1}) > \min(A_j)$. To prove this claim we observe that by construction of A_{j+1} , for any $n \in A_{j+1}$,

$$n > f(j) \text{ then } n > f(j-1)$$

and since

$$|a_n| \geq j + 1 \text{ then } |a_n| \geq j$$

we see that $A_{j+1} \subseteq A_j$, but that is not enough yet, we want $A_{j+1} \subset A_j$. We must prove that for any $j \geq 1$ there exists some $k > j$ such that

$$A_j \setminus A_k \neq \emptyset$$

therefore we need to show that there exists some $n_k \in A_k$ (that might not be the minimum a priori, but anyways we can delete any $n < n_k$ and the set will behave the same) such that for some $n_j \in A_j$ with $n_j < n_k$ it holds

$$|a_{n_k}| \geq k \text{ but } |a_{n_j}| < k.$$

Unwrapping A_j

$$A_j = \{n_j, n_{j+1}, \dots\}$$

with $|a_{n_j}| \geq j$. Consider $k := \max\{|a_{n_j}|, |a_{n_{j+1}}|, \dots\} + 1$, then

$$k > |a_{n_j}| \text{ for any } n_j \in A_j.$$

We then construct A_k with that k .

Hence we conclude that

$$\min(A_k) > \min(A_j).$$

Then, we can gather all the A_k 's recursively finding each k as in the claim

$$\min(A_1), \min(A_{k_1}), \min(A_{k_2}), \dots$$

This sequence is then monotone strictly increasing, thus $\left(\frac{1}{\min(A_j)}\right)_{j=1}^{\infty}$ is a strictly decreasing sequence. We see that the sequence

$$0 < \left| \frac{1}{\min(A_j)} \right| \leq 1$$

converges to zero, thus by zero test $\left(\frac{1}{\min(A_j)}\right)_{j=1}^{\infty}$ converges to zero. □

Point set topology

Limits and continuity