

Finite and infinite series sheet

Gallo Tennis, CoGi / to A Mathematical Room

April 17, 2025

*Problems on series and products over finite and infinite sets: books of Terence Tao's "Analysis I", Donald E. Knuth's "The Art of Computer Programming".
As well as series inequalities problems from A. Engel's "Problem-Solving Strategies" and Andreescu's "Problems From The Book".*

Convergence of series

1. Let $\sum_{n=m}^{\infty} a_n$ be a formal series of real numbers. Then $\sum_{n=m}^{\infty} a_n$ converges if and only if, for every real number $\varepsilon > 0$, there exists an integer $N \geq m$ such that

$$\left| \sum_{n=p}^q a_n \right| \leq \varepsilon \quad \text{for all } p, q \geq N.$$

Solution.

- ($\implies$) If $\sum_{n=m}^{\infty} a_n$ converges then, by definition, the partial sum sequence

$$(S_N)_{N=m}^{\infty} = \left(\sum_{n=m}^N a_n \right)_{N=m}^{\infty}$$

converges to L . Therefore, for any $\varepsilon > 0$ there exists some $M \geq m$ such that for any $p-1, q \geq M$

$$|S_q - S_{p-1}| \leq \varepsilon.$$

Without loss of generality assume $p-1 \leq q$, it follows that

$$\left| \sum_{n=m}^q a_n - \sum_{n=m}^{p-1} a_n \right| = \left| \sum_{n=p}^q a_n \right| \leq \varepsilon.$$

- ($\impliedby$) Doing the reverse process of the direct implication the result follows.

□

2. **(Zero test).** Let $\sum_{n=m}^{\infty} a_n$ be a convergent series of real numbers. Then we must have $\lim_{n \rightarrow \infty} a_n = 0$. To put this another way, if $\lim_{n \rightarrow \infty} a_n$ is non-zero or divergent, then the series $\sum_{n=m}^{\infty} a_n$ is divergent.

Solution. If $\sum_{n=m}^{\infty} a_n$ converges then by last problem, for any $\varepsilon > 0$, there exists some $M \geq m$ such that for any $p, q \geq M$

$$\left| \sum_{n=p}^q a_n \right| \leq \varepsilon. \quad (1)$$

Suppose for the sake of contradiction that $\lim_{n \rightarrow \infty} a_n = L > 0$ so that for any $\varepsilon > 0$ there exists some M' such that for any $n \geq M'$

$$|a_n - L| < \varepsilon$$

so that in particular $a_n \geq L - \varepsilon$.

Therefore, for any $p, q \geq K := \max\{M, M'\}$

$$\left| \sum_{n=p}^q a_n \right| \geq \left| \sum_{n=p}^q L - \varepsilon \right| = (q - p + 1) |L - \varepsilon|.$$

Hence, let $\varepsilon = L/2$, we see that if the above inequality were true

$$\left| \sum_{n=p}^q a_n \right| \geq \frac{(q - p + 1)L}{2}.$$

Then let $q = p + 1$, it follows that

$$\left| \sum_{n=p}^q a_n \right| \geq L > \varepsilon.$$

A contradiction of inequality (1).

□

3. **(Absolute convergence test).** Let $\sum_{n=m}^{\infty} a_n$ be a formal series of real numbers. If this series is absolutely convergent, then it is also conditionally convergent. Furthermore, in this case we have the triangle inequality

$$\left| \sum_{n=m}^{\infty} a_n \right| \leq \sum_{n=m}^{\infty} |a_n|.$$

Solution. If $\sum_{n=m}^{\infty} |a_n|$ converges to L , then for any $\varepsilon > 0$ there exists some $M \geq m$ such that for any $p, q \geq M$

$$\left| \sum_{n=p}^q |a_n| \right| \leq \varepsilon.$$

Hence,

$$\sum_{n=p}^q |a_n| \leq \varepsilon.$$

By triangle inequality over finite series, for any $p, q \geq M$ we have

$$\left| \sum_{n=p}^q a_n \right| \leq \sum_{n=p}^q |a_n|.$$

We conclude that $\sum_{n=p}^q a_n$ converges. Trivially, we also note that again, by triangle inequality over finite series, each partial sum $S_N = \sum_{n=m}^N a_n$ is smaller than or equal to $T_N = \sum_{n=m}^N |a_n|$, meaning that by comparison principle $\lim_{N \rightarrow \infty} S_N \leq \lim_{N \rightarrow \infty} T_N$.

□

4. **(Series laws).**

- (a) If $\sum_{n=m}^{\infty} a_n$ is a series of real numbers converging to x , and $\sum_{n=m}^{\infty} b_n$ is a series of real numbers converging to y , then $\sum_{n=m}^{\infty} (a_n + b_n)$ is also a convergent series, and converges to $x + y$.

In particular, we have

$$\sum_{n=m}^{\infty} (a_n + b_n) = \sum_{n=m}^{\infty} a_n + \sum_{n=m}^{\infty} b_n.$$

- (b) If $\sum_{n=m}^{\infty} a_n$ is a series of real numbers converging to x , and c is a real number, then $\sum_{n=m}^{\infty} (ca_n)$ is also a convergent series, and converges to cx .

In particular, we have

$$\sum_{n=m}^{\infty} (ca_n) = c \sum_{n=m}^{\infty} a_n.$$

- (c) Let $\sum_{n=m}^{\infty} a_n$ be a series of real numbers, and let $k \geq 0$ be an integer. If one of the two series $\sum_{n=m}^{\infty} a_n$ and $\sum_{n=m+k}^{\infty} a_n$ are convergent, then the other one is also, and we have the identity

$$\sum_{n=m}^{\infty} a_n = \sum_{n=m}^{m+k-1} a_n + \sum_{n=m+k}^{\infty} a_n.$$

- (d) Let $\sum_{n=m}^{\infty} a_n$ be a series of real numbers converging to x , and let k be an integer. Then $\sum_{n=m+k}^{\infty} a_n$ also converges to x .

Solution. Let $T_N = \sum_{n=m}^N a_n$ and $R_N = \sum_{n=m}^N b_n$.

- (a) The partial sum of the series $\sum_{n=m}^{\infty} a_n + b_n$ is trivially

$$\sum_{n=m}^N (a_n + b_n) = T_N + R_N.$$

we also note that by limit laws

$$\lim_{n \rightarrow \infty} T_N + R_N = \lim_{n \rightarrow \infty} T_N + \lim_{n \rightarrow \infty} R_N.$$

We conclude that

$$\sum_{n=m}^{\infty} (a_n + b_n) = \lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} T_N + R_N = x + y.$$

- (b) Similarly to the item before, due to the linear properties of both limit and partial sums we get

$$\sum_{n=m}^N ca_n = cT_N$$

so that $\lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} cT_N = c \lim_{N \rightarrow \infty} T_N = cx$.

- (c) If $\sum_{n=m}^{\infty} a_n$ converges then the first claim is trivial, follows from convergent limit properties.

5. **(Telescoping series).** Let $(a_n)_{n=0}^{\infty}$ be a sequence of real numbers which converge to 0, i.e., $\lim_{n \rightarrow \infty} a_n = 0$. Then the series

$$\sum_{n=0}^{\infty} (a_n - a_{n+1})$$

converges to a_0 .

Solution. We see that for any partial sum S_N starting from 0,

$$\sum_{n=0}^N a_n - a_{n+1} = \sum_{n=0}^N a_n - \sum_{n=0}^N a_{n+1}$$

doing a substitution $n \rightarrow n + 1$ we get

$$S_N = \sum_{n=0}^N a_n - \sum_{n=1}^{N+1} a_n = a_0 + \sum_{n=1}^N a_n - \sum_{n=1}^N a_n + a_{N+1}$$

therefore,

$$S_N = a_0 - a_{N+1}.$$

We notice since (a_n) converges and by using limit laws that

$$\lim_{N \rightarrow \infty} S_N = \lim_{N \rightarrow \infty} a_0 - \lim_{N \rightarrow \infty} a_{N+1} = a_0.$$

□

6. (Comparison test.)
7. (Geometric series.)
8. (Cauchy criterion.)
9. (Root test.)
10. (Ratio test.)

Series classification

1. Let $(a_n)_{n \geq 1}$ be a sequence of positive real numbers such that the series $\sum_{n=1}^{\infty} a_n$ converges. Show that the series

$$\sum_{n=1}^{\infty} a_n^{\frac{n}{n+1}}$$

Also converges.

2. Show that

$$(a) \quad S_0(n) = 1^0 + \cdots + n^0 = n,$$

$$S_1(n) = 1^1 + \cdots + n^1 = \frac{n(n+1)}{2} = \frac{1}{2}n^2 + \frac{1}{2}n,$$

$$S_2(n) = 1^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6} = \frac{1}{3}n^3 + \frac{1}{2}n^2 + \frac{1}{6}n,$$

$$S_3(n) = \frac{n^2(n+1)^2}{4} = \frac{1}{4}n^4 + \frac{1}{2}n^3 + \frac{1}{4}n^2,$$

and in general that

$$S_k(n) = a_{k+1}n^{k+1} + \cdots + a_1n + a_0$$

is a polynomial in n of degree $k+1$.

$$(b) \quad \lim_{n \rightarrow \infty} \frac{S_k(n)}{n^{k+1}}.$$

3. Show that

(a) the equality

$$1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!} + \frac{1}{n!n} = 3 - \frac{1}{1 \cdot 2 \cdot 2!} - \cdots - \frac{1}{(n-1) \cdot n \cdot n!}$$

holds for $n \geq 2$;

(b) $e = 3 - \sum_{n=0}^{\infty} \frac{1}{(n+1)(n+2)(n+2)!};$

(c) for computing the number e approximately the formula

$$e \approx 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!} + \frac{1}{n!n}$$

is much better than the original formula

$$e \approx 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}.$$

Power series and generating functions

Application

1. König's theorem.