

PROBLEM SET 5 (T-445-GRTH)

You need to collect **60** points to get a full score **but** you cannot get more than **X** points (in total) from a problem section with annotation **max X**.

Please make sure to:

1. Write your name/email(s) on your work (replace my name above).
2. Write your answers in `\begin{solution}` ... `\end{solution}` blocks.
3. Write clear and concise proofs: points may be deducted for vagueness.

1. INDEPENDENT SETS (**max 30**)

- 1** (10 points) Find a recursive formula for counting the number of *maximal* independent sets in P_n ($n \geq 1$). Use it to derive a formula counting the number of maximal independent sets in C_n .

Solution. We order the vertices from one end to the other. Take v_1 . From every set, create two new sets by adding the vertices v_{i+2} or v_{i+3} to the set of v_i was the last vertex. Redo this step as often as it is possible (both new vertices still exist). If only v_{i+2} exists, add it to the set. Do the same again but starting with v_2 . Count all sets. This is the number of maximal independent sets in P_n .

For C_n we can do the same, but if we start with v_1 , it is not allowed to have v_n in the same set. So the first and the second part create the same amount of independent sets. Just do the algorithm with starting at v_2 and double the number. \square

- 2** (5 points) Show that:

- a** For every simple graph G , $\alpha(G) \geq \frac{|V(G)|}{D(G)+1}$.
- b** For every planar graph G , $\alpha(G) \geq \frac{|V(G)|}{4}$.

Solution. b) From the lecture we know $\alpha(G) \geq \frac{|V(G)|}{\chi(G)}$. The 4-color-theorem says every planar graph has chromatic number ≤ 4 . In total

$$\alpha(G) \geq \frac{|V(G)|}{\chi(G)} \geq \frac{|V(G)|}{4}$$

. \square

- 3** (7 points) Show that:

- a** For every simple graph G , $\alpha(G) \leq |V(G)| - \delta(G)$.
- b** For every simple triangle-free graph G , $\alpha(G) \geq \Delta(G)$.

Solution. a) Take an arbitrary maximal independent set and a vertex v from this set. All neighbors of this vertex can not be in the maximal independent set and this are at least $\delta(G)$. So $\alpha(G) \leq |V(G)| - \delta(G)$. b) Take a vertex v with degree $\Delta(G)$. Take all adjacent vertices of v . This is an independent set, because the graph is triangle-free and has size $\Delta(G)$. So the claim holds. \square

- 4 (10 points) Show that every simple triangle-free graph has an independent set of size \sqrt{n} .

2. BIPARTITE MATCHING (max 26)

- 5 (5 points) Let $G = (X \cup Y, E)$ be a simple bipartite graph. Suppose that there is a $k \in \mathbb{N}$ such that $d(x) \geq k \geq d(y)$ for all $x \in X, y \in Y$. Show that G has a matching saturating X (covering all of X).
- 6 (6 points) Let $G = (X \cup Y, E)$ be a simple bipartite graph. Let A be the set of vertices in G of maximum degree. Show that G has a matching saturating $A \cap X$.
- 7 (6 points) Let $G = (X \cup Y, E)$ be a simple bipartite graph such that $d(x) \geq 1$ for all $x \in X$ and $d(x) \geq d(y)$ for all $\{x, y\} \in E$ ($x \in X, y \in Y$). Show that G has a matching saturating X .
- 8 (10 points) Let $G = (X \cup Y, E)$ be a simple bipartite graph containing a perfect matching. Prove that there is a vertex $x \in X$, such that for every incident edge $\{x, y\}$, there is a perfect matching that contains $\{x, y\}$.
- 9 (6 points) Let X, Y be disjoint independent sets in a simple graph G , such that $|X| = \alpha(G)$. Prove that $G[X \cup Y]$ has a matching of size $|Y|$.

Solution. For every vertex in Y , we can find a vertex in X and connect them, such that it is a matching. This is possible because otherwise, you would find a subset $S \subseteq Y$ and the set of adjacent vertices $S' \subseteq Y$ with $|S| > |S'|$. But if this holds, we can replace S' by S in X and still have an independent subset. This subset would have more vertices than $\alpha(G)$ and therefor is a contradiction. Hence we can find a matching of size $|Y|$. \square

3. MORE MATCHING (max 50)

- 10 (6 points) Let G be a simple graph with an even number of vertices such that $d(v) \geq \frac{|V(G)|}{2}$ for every vertex $v \in V(G)$. Show that G has a perfect matching.
- 11 (9 points) Let t be a tree on n vertices with l leaves. Show that t has a matching of size at least $\lceil \frac{n-l}{2} \rceil$.
- 12 (10 points) For every $k \geq 1$, show that every simple k -regular graph has a matching of size at least $\frac{n}{4 - \frac{2}{k}}$.
Hint: show that this bound holds for every maximal matching.
- 13 (10 points) Show that a tree has either one perfect matching or none. For a graph G , denote by $o(G)$ the number of components of G of odd cardinality. Prove that, for a tree T , T has a perfect matching, if and only if, $o(T - v) = 1$ for all vertices v .
- 14 (15 points) Prove that every simple bridgeless cubic graph (3-regular graph) has a perfect matching. Furthermore, show that there is a cubic graph that has a bridge and does not have a perfect matching.
Hint: Use Tutte's theorem.

- 15** (13 points) Let G be a graph and M be a maximal matching in G .
- a** Show that if there is no M -augmenting path of length three then $|M| \geq \frac{2}{3} \cdot \text{opt}$, where opt is the size of a maximum matching.
 - b** Suppose now that for a given $k > 1$, there is no M -augmenting path of length $2k+1$ or shorter. Prove a better bound on $|M|$ than the one above, and show that your bound is tight by providing an example.