

## PROBLEM SET 5 (T-445-GRTH)

You need to collect **60** points to get a full score **but** you cannot get more than **X** points (in total) from a problem section with annotation **max X**.

**Please make sure to:**

1. Write your name/email(s) on your work (replace my name above).
2. Write your answers in `\begin{solution}` ... `\end{solution}` blocks.
3. Write clear and concise proofs: points may be deducted for vagueness.

### 1. INDEPENDENT SETS (**max 30**)

- 1** (10 points) Find a recursive formula for counting the number of *maximal* independent sets in  $P_n$  ( $n \geq 1$ ). Use it to derive a formula counting the number of maximal independent sets in  $C_n$ .

**Solution.** We order the vertices from one end to the other. Take  $v_1$ . From every set, create two new sets by adding the vertices  $v_{i+2}$  or  $v_{i+3}$  to the set of  $v_i$  was the last vertex. Redo this step as often as it is possible (both new vertices still exist). If only  $v_{i+2}$  exists, add it to the set. Do the same again but starting with  $v_2$ . Count all sets. This is the number of maximal independent sets in  $P_n$ .

For  $C_n$  we can do the same, but if we start with  $v_1$ , it is not allowed to have  $v_n$  in the same set. So the first and the second part create the same amount of independent sets. Just do the algorithm with starting at  $v_2$  and double the number.  $\square$

- 2** (5 points) Show that:

- a For every simple graph  $G$ ,  $\alpha(G) \geq \frac{|V(G)|}{D(G)+1}$ .
- b For every planar graph  $G$ ,  $\alpha(G) \geq \frac{|V(G)|}{4}$ .

**Solution.**

a) For a simple graph  $G$ , if for each vertex  $v_i$  in  $G$ , you remove  $v_i$  and its adjacent neighbours, and continue removing each vertex in this fashion until there are no vertices left, then  $\{v_1, v_2, \dots, v_i\}$  is an independent set. Because each vertex can have at most  $D(G)$  neighbours, we can remove at most  $D(G) + 1$  vertices. Therefore there can be at most  $\frac{|V(G)|}{D(G)+1}$  removals, and therefore the independent set must have at least this many vertices.

Therefore for every simple graph  $G$ ,  $\alpha(G) \geq \frac{|V(G)|}{D(G)+1}$ .

b) From the lecture we know  $\alpha(G) \geq \frac{|V(G)|}{\chi(G)}$ . The 4-color-theorem says every planar graph has chromatic number  $\leq 4$ . In total

$$\alpha(G) \geq \frac{|V(G)|}{\chi(G)} \geq \frac{|V(G)|}{4}$$

.

$\square$

- 3** (7 points) Show that:

- a For every simple graph  $G$ ,  $\alpha(G) \leq |V(G)| - \delta(G)$ .
- b For every simple triangle-free graph  $G$ ,  $\alpha(G) \geq \Delta(G)$ .

**Solution.** a) Take an arbitrary maximal independent set and a vertex  $v$  from this set. All neighbors of this vertex can not be in the maximal independent set and this are at least  $\delta(G)$ . So  $\alpha(G) \leq |V(G)| - \delta(G)$ . b) Take a vertex  $v$  with degree  $\Delta(G)$ . Take all adjacent vertices of  $v$ . This is an independent set, because the graph is triangle-free and has size  $\Delta(G)$ . So the claim holds.  $\square$

- 4 (10 points) Show that every simple triangle-free graph has an independent set of size of at least  $\lfloor \sqrt{n} \rfloor$ .

**Solution.** For a simple triangle-free graph  $G$  we know that if there exists a vertex  $u$  with a degree of greater than  $\lfloor \sqrt{n} \rfloor$ , then there exists an independent set with each neighbour adjacent to  $u$ , because the neighbours cannot be adjacent to each other because if they were it would create a triangle. Otherwise all vertices would have a degree of less than  $\lfloor \sqrt{n} \rfloor$ . We know that if there exists a vertex  $v$  with a degree of less than  $\lfloor \sqrt{n} \rfloor$ , then there must exist a maximal independent set greater than  $\lfloor \sqrt{n} \rfloor$ . Therefore based on the upper and lower bounds being greater than  $\lfloor \sqrt{n} \rfloor$ , every simple triangle-free graph has an independent set of size of at least  $\lfloor \sqrt{n} \rfloor$ .  $\square$

## 2. BIPARTITE MATCHING (max 26)

- 5 (5 points) Let  $G = (X \cup Y, E)$  be a simple bipartite graph. Suppose that there is a  $k \in \mathbb{N}$  such that  $d(x) \geq k \geq d(y)$  for all  $x \in X, y \in Y$ . Show that  $G$  has a matching saturating  $X$  (covering all of  $X$ ).

**Solution.** We start with an arbitrary  $x_1 \in X$  and match it to an adjacent  $y_i \in Y$ , that is not matched to an other  $x_j$  by now. We continue like this with the next  $x$  if possible. If there is no free adjacent vertex, we reorder the matchings from before, so this vertex gets a new free adjacent vertex. This is possible, because if the other vertices would all be connected just to this adjacent vertices, it would hold  $d(y) > d(x)$  for those  $x$  and  $y$ . This is a contradiction and so, the assumption holds.  $\square$

- 6 (6 points) Let  $G = (X \cup Y, E)$  be a simple bipartite graph. Let  $A$  be the set of vertices in  $G$  of maximum degree. Show that  $G$  has a matching saturating  $A \cap X$ .

**Solution.** In a more general way but with the same explanation as in problem 5, we can state that if a bipartite has no matching saturating  $A \subset X$ , there exists a subset  $S \subset A$ , such that the number of neighbors of  $S$  are less than  $|S|$ . We use this claim in the following proofs.

Take the set of vertices  $A \cap X =: S$ . Assume that there exists a subset  $S' \subset S$  that has less neighbors than  $|S'|$ . In total (with double counting) the vertices of  $S'$  are connected to  $\Delta(G) \cdot |S'|$  vertices. That's the total degree of the adjacent vertices. If they are less than  $|S'|$ , the average degree is higher than  $\Delta(G)$ , which is a contradiction. In total the assumption holds.  $\square$

- 7** (6 points) Let  $G = (X \cup Y, E)$  be a simple bipartite graph such that  $d(x) \geq 1$  for all  $x \in X$  and  $d(x) \geq d(y)$  for all  $\{x, y\} \in E$  ( $x \in X, y \in Y$ ). Show that  $G$  has a matching saturating  $X$ .

**Solution.** We use the same claim as in problem 6. The proof is quite similar. Assume there exists a subset  $S \subset X$ , with less neighbors than  $|S|$ . Then there would be a  $y \in Y$  with  $d(y) > d(x)$  for an adjacent  $x$  with the same reason as in problem 6. This leads one more time to a contradiction and proves our claim.  $\square$

- 8** (10 points) Let  $G = (X \cup Y, E)$  be a simple bipartite graph containing a perfect matching. Prove that there is a vertex  $x \in X$ , such that for every incident edge  $\{x, y\}$ , there is a perfect matching that contains  $\{x, y\}$ .

**Solution.** Let  $G = (X \cup Y, E)$  be a simple bipartite graph containing a perfect matching. A matching is only perfect if it has  $\lfloor \frac{n}{2} \rfloor$  edges, which means  $G$  has an even amount of vertices and that  $|X| = |Y|$  so the perfect match  $M$  in  $G$  must have  $\frac{n}{2}$  edges. Then if  $M$  is perfect and  $G$  is not disconnected we know that for each match in  $M$  there must be an edge not in  $M$  that connects each match. This implies that there exists an alternating path  $P$  in  $G$ , which means for each edge in  $M$  that connects  $u, v$ , there are two corresponding edges in  $P/M$  that connect both it's vertices. Therefore the edges in  $P/M$  provide a perfect match not equal to  $M$ .  $\square$

- 9** (6 points) Let  $X, Y$  be disjoint independent sets in a simple graph  $G$ , such that  $|X| = \alpha(G)$ . Prove that  $G[X \cup Y]$  has a matching of size  $|Y|$ .

**Solution.** For every vertex in  $Y$ , we can find a vertex in  $X$  and connect them, such that it is a matching. This is possible because otherwise, you would find a subset  $S \subseteq Y$  and the set of adjacent vertices  $S' \subseteq Y$  with  $|S| > |S'|$ . But if this holds, we can replace  $S'$  by  $S$  in  $X$  and still have an independent subset. This subset would have more vertices than  $\alpha(G)$  and therefor is a contradiction. Hence we can find a matching of size  $|Y|$ .  $\square$

### 3. MORE MATCHING (max 50)

- 10** (6 points) Let  $G$  be a simple graph with an even number of vertices such that  $d(v) \geq \frac{|V(G)|}{2}$  for every vertex  $v \in V(G)$ . Show that  $G$  has a perfect matching.

**Solution.** If  $G$  is a simple graph with an even number of vertices, then arbitrarily splitting  $G$  into two sets  $G = S1 \cup S2$  such that  $|S1| = |S2|$ . Then we know from the definition that each vertex  $v \in V(G)$ ,  $d(v) \geq \frac{|V(G)|}{2}$ , so we know that each set contains  $\frac{|V(G)|}{2}$  and each vertex must have a degree of at least  $\frac{|V(G)|}{2}$ . Therefore a vertex can be connected to at most  $\frac{|V(G)|}{2} - 1$  vertices in it's own set, and therefore every vertex must be connected to a vertex in the opposite set. Therefore every pair of vertices have at least one edge connecting them, implying that a perfect match must exist between  $S1$  and  $S2$  and therefore in  $G$  as well.  $\square$

- 11 (9 points) Let  $t$  be a tree on  $n$  vertices with  $l$  leaves. Show that  $t$  has a matching of size at least  $\lceil \frac{n-l}{2} \rceil$ .

**Solution.** We can show that any tree is a bipartite graph because at any level of the tree there are no nodes connected to other nodes at that level, so if we put all odd numbered levels in one set, and all even numbered levels in another set so we have two sets of nodes where each node is only connected to a node in the other set. Therefore every tree is bipartite. From this claim  $\square$

- 12 (10 points) For every  $k \geq 1$ , show that every simple  $k$ -regular graph has a matching of size at least  $\frac{n}{4-\frac{2}{k}}$ .

Hint: show that this bound holds for every maximal matching.

- 13 (10 points) Show that a tree has either one perfect matching or none. For a graph  $G$ , denote by  $o(G)$  the number of components of  $G$  of odd cardinality. Prove that, for a tree  $T$ ,  $T$  has a perfect matching, if and only if,  $o(T - v) = 1$  for all vertices  $v$ .

- 14 (15 points) Prove that every simple bridgeless cubic graph (3-regular graph) has a perfect matching. Furthermore, show that there is a cubic graph that has a bridge and does not have a perfect matching.

Hint: Use Tutte's theorem.

- 15 (13 points) Let  $G$  be a graph and  $M$  be a maximal matching in  $G$ .

- a Show that if there is no  $M$ -augmenting path of length three then  $|M| \geq \frac{2}{3} \cdot \text{opt}$ , where  $\text{opt}$  is the size of a maximum matching.
- b Suppose now that for a given  $k > 1$ , there is no  $M$ -augmenting path of length  $2k+1$  or shorter. Prove a better bound on  $|M|$  than the one above, and show that your bound is tight by providing an example.