



Unbiased Estimators

Dirk Grunwald
University of Colorado, Boulder

Unbiased Estimators

- Estimates
- Estimators
- Bias vs. Unbiased
- Estimator of the Mean
- Estimator of Variance

Estimates

Given a dataset x_1, x_2, \dots, x_n from a set of random variables X_1, X_2, \dots, X_n ,

Estimates

Given a dataset x_1, x_2, \dots, x_n from a set of random variables X_1, X_2, \dots, X_n ,

Estimates

An *estimate* is a value t that only depends on the dataset x_1, x_2, \dots, x_n ; *i.e.* t is some function of the dataset only:

$$t = h(x_1, x_2, \dots, x_n).$$

Estimates

Given a dataset x_1, x_2, \dots, x_n from a set of random variables X_1, X_2, \dots, X_n ,

Estimates

An *estimate* is a value t that only depends on the dataset x_1, x_2, \dots, x_n ; *i.e.* t is some function of the dataset only:

$$t = h(x_1, x_2, \dots, x_n).$$

The estimate t can be a single number (e.g. $E[X]$), a vector or a complex distribution function.

Estimates

Given a dataset x_1, x_2, \dots, x_n from a set of random variables X_1, X_2, \dots, X_n ,

Estimates

An *estimate* is a value t that only depends on the dataset x_1, x_2, \dots, x_n ; *i.e.* t is some function of the dataset only:

$$t = h(x_1, x_2, \dots, x_n).$$

The estimate t can be a single number (e.g. $E[X]$), a vector or a complex distribution function.

E.g., mean of the Poisson distribution or predicted wood hardness

Estimator

Estimator

Let $t = h(x_1, x_2, \dots, x_n)$ be a estimate based on dataset x_1, x_2, \dots, x_n . Then t is a realization of the random variable

$$T = h(x_1, x_2, \dots, x_n).$$

and random variable T is called an *estimator*.

Estimator

Estimator

Let $t = h(x_1, x_2, \dots, x_n)$ be a estimate based on dataset x_1, x_2, \dots, x_n . Then t is a realization of the random variable

$$T = h(x_1, x_2, \dots, x_n).$$

and random variable T is called an *estimator*.

E.g., two estimators for the same mean of the Poisson distribution or a linear model relating wood hardness and density.

Unbiased Estimators

Unbiased Estimators

A estimator T is called an *unbiased estimator* for the parameter θ if

$$E[T] = \theta$$

irrespective of the value of θ . The difference $E[T] - \theta$ is called the *bias* of T ; if this difference is non-zero, then T is *biased*.

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Model *variance* is refers to the amount the model changes with different data

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Model *variance* is refers to the amount the model changes with different data

Mode *bias* refers to the error introduced by approximating a solution using a simpler model

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Model *variance* is refers to the amount the model changes with different data

Mode *bias* refers to the error introduced by approximating a solution using a simpler model

Models are trade-offs between *variance* and *bias*.

Biased Estimators: Example

- We built two estimators, S and T , to estimate μ of Poisson

Biased Estimators: Example

- We built two estimators, S and T , to estimate μ of Poisson
- Estimator S is *unbiased*, and $E[S] = p_0$ for all p_0 .

Biased Estimators: Example

- We built two estimators, S and T , to estimate μ of Poisson
- Estimator S is *unbiased*, and $E[S] = p_0$ for all p_0 .
- Estimator T is *biased*, and over-predicts p_0 .




Fig. 19.4. $E[T]$ as a function of n .

Biased Estimators: Example

- We built two estimators, S and T , to estimate μ of Poisson
- Estimator S is *unbiased*, and $E[S] = p_0$ for all p_0 .
- Estimator T is *biased*, and over-predicts p_0 .




Fig. 19.4. $E[T]$ as a function of n .

The bias occurs because

$$E[T] = E[e^{-\bar{X}_n}] > e^{E[-\bar{X}_n]}$$

due to Jensen's inequality.

Who cares?

- Two common cases:

Who cares?

- Two common cases:
- Unbiased estimate of the mean:

$$E[\bar{X}_n] = \frac{X_1 + X_2 + \dots + X_n}{n} = \mu$$

Who cares?

- Two common cases:
- Unbiased estimate of the mean:

$$E[\bar{X}_n] = \frac{X_1 + X_2 + \dots + X_n}{n} = \mu$$

- Unbiased estimate of the variance:

$$S_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

Who cares?

- Two common cases:
- Unbiased estimate of the mean:

$$E[\bar{X}_n] = \frac{X_1 + X_2 + \dots + X_n}{n} = \mu$$

- Unbiased estimate of the variance:

$$S_n^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X}_n)^2$$

- Why do we divide by $n-1$ rather than n ?

Unbiased estimate for μ

Suppose $X_1 + X_2 + \dots + X_n$ is a random sample from a distribution with mean μ and variance σ^2 .

$$\begin{aligned} E[\bar{X}_n] &= \frac{X_1 + X_2 + \dots + X_n}{n} \\ &= \frac{1}{n} (E[X_1] + E[X_2] + \dots + E[X_n]) \\ &= \frac{1}{n} (\mu + \mu + \dots + \mu) = \mu \end{aligned}$$

Unbiased estimate for μ

Suppose $X_1 + X_2 + \dots + X_n$ is a random sample from a distribution with mean μ and variance σ^2 .

$$\begin{aligned} E[\bar{X}_n] &= \frac{X_1 + X_2 + \dots + X_n}{n} \\ &= \frac{1}{n} (E[X_1] + E[X_2] + \dots + E[X_n]) \\ &= \frac{1}{n} (\mu + \mu + \dots + \mu) = \mu \end{aligned}$$

This holds for all μ and thus \bar{X}_n is an unbiased estimator for μ .

Variance of Sample vs Estimated Variance

N.B.

$$\text{Var}[\bar{X}_n] \neq S_n^2$$

$\text{Var}[\bar{X}_n]$ is variance of a sample of X_i with mean μ and variance σ^2 .

$$\begin{aligned}\text{Var}[\bar{X}_n] &= \text{Var}\left[\frac{1}{n} (X_1 + X_2 + \dots + X_n)\right] \\ &= \left(\frac{1}{n}\right)^2 \text{Var}[X_1 + X_2 + \dots + X_n] \\ &= \frac{1}{n^2} (\sigma^2 + \dots + \sigma^2) = \frac{\sigma^2}{n}\end{aligned}$$

Variance of Sample vs Estimated Variance

N.B.

$$\text{Var}[\bar{X}_n] \neq S_n^2$$

$\text{Var}[\bar{X}_n]$ is variance of a sample of X_i with mean μ and variance σ^2 .

$$\begin{aligned}\text{Var}[\bar{X}_n] &= \text{Var}\left[\frac{1}{n} (X_1 + X_2 + \dots + X_n)\right] \\ &= \left(\frac{1}{n}\right)^2 \text{Var}[X_1 + X_2 + \dots + X_n] \\ &= \frac{1}{n^2} (\sigma^2 + \dots + \sigma^2) = \frac{\sigma^2}{n}\end{aligned}$$

S_n^2 is an estimator for the σ^2 of X_i .

Estimating S_n^2

We stated:

$$E[S_n^2] = \sigma^2 = \frac{1}{n-1} \sum_{i=1}^n E[(X_i - \bar{X}_n)^2]$$

Estimating S_n^2

We stated:

$$\begin{aligned} E[S_n^2] &= \sigma^2 = \frac{1}{n-1} \sum_{i=1}^n E[(X_i - \bar{X}_n)^2] \\ &= \frac{1}{n-1} \sum_{i=1}^n \text{Var}[X_i - \bar{X}_n] \quad \text{Magic \#1} \end{aligned}$$

Estimating S_n^2

We stated:

$$\begin{aligned} E[S_n^2] &= \sigma^2 = \frac{1}{n-1} \sum_{i=1}^n E[(X_i - \bar{X}_n)^2] \\ &= \frac{1}{n-1} \sum_{i=1}^n \text{Var}[X_i - \bar{X}_n] \quad \text{Magic \#1} \\ &= \frac{1}{n-1} * n * \left(\frac{n-1}{n} \sigma^2 \right) \quad \text{Magic \#2} \\ &= \sigma^2 \end{aligned}$$

Magic #1

We need to show that

$$\text{Var}[X_i - \bar{X}_n] = E[(X_i - \bar{X}_n)^2]$$

Magic #1

We need to show that

$$\text{Var}[X_i - \bar{X}_n] = E[(X_i - \bar{X}_n)^2]$$

Side trip!

Earlier def'n of variance sez:

$$\text{Var}[Y] = E[Y^2] - (E[Y])^2$$

Magic #1

We need to show that

$$\text{Var}[X_i - \bar{X}_n] = E[(X_i - \bar{X}_n)^2]$$

Side trip!

Earlier def'n of variance sez:

$$\text{Var}[Y] = E[Y^2] - (E[Y])^2$$

Let $Y = (X_i - \bar{X}_n)$.

Magic #1

We need to show that

$$\text{Var}[X_i - \bar{X}_n] = E[(X_i - \bar{X}_n)^2]$$

Side trip!

Earlier def'n of variance sez:

$$\text{Var}[Y] = E[Y^2]$$

Let $Y = (X_i - \bar{X}_n)$.

$$E[Y] = E[(X_i - \bar{X}_n)] = E[X_i] - E[\bar{X}_n] = \mu - \mu = 0$$

Magic #1

We need to show that

$$\text{Var}[X_i - \bar{X}_n] = E[(X_i - \bar{X}_n)^2]$$

Side trip!

Earlier def'n of variance sez:

$$\text{Var}[Y] = E[Y^2]$$

Let $Y = (X_i - \bar{X}_n)$.

$$E[Y] = E[(X_i - \bar{X}_n)] = E[X_i] - E[\bar{X}_n] = \mu - \mu = 0$$

Thus,

$$\text{Var}[Y] = \text{Var}[X_i - \bar{X}_n] = E[(X_i - \bar{X}_n)^2]$$

Magic #2

We want to show $\text{Var}[X_i - \bar{X}_n] = \frac{n-1}{n}\sigma^2$.

Magic #2

We want to show $\text{Var}[X_i - \bar{X}_n] = \frac{n-1}{n}\sigma^2$.

Side trip!

$$\begin{aligned} X_i - \bar{X}_n &= \frac{n}{n}X_i - \left(\frac{X_i}{n} + \frac{1}{n} \sum_{j \neq i} X_j \right) \\ &= \frac{n-1}{n}X_i - \frac{1}{n} \sum_{j \neq i} X_j \end{aligned}$$

Magic #2

We want to show $\text{Var}[X_i - \bar{X}_n] = \frac{n-1}{n}\sigma^2$.

$$\text{Var}[X_i - \bar{X}_n] = \text{Var}\left[\frac{n-1}{n}X_i - \frac{1}{n}\sum_{j \neq i} X_j\right]$$

Magic #2

We want to show $\text{Var}[X_i - \bar{X}_n] = \frac{n-1}{n}\sigma^2$.

$$\begin{aligned}\text{Var}[X_i - \bar{X}_n] &= \text{Var}\left[\frac{n-1}{n}X_i - \frac{1}{n}\sum_{j \neq i} X_j\right] \\ &= \left(\frac{n-1}{n}\right)^2 \text{Var}[X_i] + \left(\frac{1}{n}\right)^2 \text{Var}\left[\sum_{j \neq i} X_j\right]\end{aligned}$$

Magic #2

We want to show $\text{Var}[X_i - \bar{X}_n] = \frac{n-1}{n}\sigma^2$.

$$\begin{aligned}\text{Var}[X_i - \bar{X}_n] &= \text{Var}\left[\frac{n-1}{n}X_i - \frac{1}{n}\sum_{j \neq i} X_j\right] \\&= \left(\frac{n-1}{n}\right)^2 \text{Var}[X_i] + \left(\frac{1}{n}\right)^2 \text{Var}\left[\sum_{j \neq i} X_j\right] \\&= \left(\left(\frac{n-1}{n}\right)^2 + \frac{n-1}{n^2}\right)\sigma^2 \\&= \frac{n^2 - 2n + 1 + n - 1}{n^2}\sigma^2 = \frac{n(n-1)}{n * n}\sigma^2 = \frac{n-1}{n}\sigma^2\end{aligned}$$

Back to where we were..

$$\begin{aligned} E[S_n^2] &= \sigma^2 = \frac{1}{n-1} \sum_{i=1}^n E[(X_i - \bar{X}_n)^2] \\ &= \frac{1}{n-1} \sum_{i=1}^n \text{Var}[X_i - \bar{X}_n] \quad \text{Magic \#1} \\ &= \frac{1}{n-1} * n * \left(\frac{n-1}{n} \sigma^2 \right) \quad \text{Magic \#2} \\ &= \sigma^2 \end{aligned}$$

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Model *variance* refers to the amount the model changes with different data

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Model *variance* refers to the amount the model changes with different data

Model *bias* refers to the error introduced by approximating a solution using a simpler model

Bias and Machine Learning

Machine learning (e.g. a regression model) is trying learn a rule from data

It's *estimating* the parameters from the data

Model *variance* refers to the amount the model changes with different data

Model *bias* refers to the error introduced by approximating a solution using a simpler model

Models are trade-offs between *variance* and *bias*.