



Weak Law of Large Numbers

Dirk Grunwald
University of Colorado, Boulder

Weak Law of Large Numbers

Overview: Apply Chebyshev's inequality to the average \bar{X}_n where we use $E[\bar{X}_n] = \mu$ and $Var(\bar{X}_n) = \sigma^2/n$ and where $\epsilon > 0$.

Overview: Apply Chebyshev's inequality to the average \bar{X}_n where we use $E[\bar{X}_n] = \mu$ and $Var(\bar{X}_n) = \sigma^2/n$ and where $\epsilon > 0$.

$$\begin{aligned} P(|\bar{X}_n - \mu| > \epsilon) &= P(|\bar{X}_n - E[\bar{X}_n]| > \epsilon) \\ &\leq \frac{1}{\epsilon^2} Var(\bar{X}_n) \\ &\leq \frac{\sigma^2}{n\epsilon^2} \end{aligned}$$

Definition: Weak Law of Large Numbers

Weak Law of Large Numbers - if \hat{X}_n is the average of n independent random variables with expectation μ and variance σ^2 , then for any $\epsilon > 0$:

$$\lim_{n \rightarrow \infty} P(|\bar{X}_n - \mu| > \epsilon) = 0$$

We can increase the precision of measurements by taking more measurements.

Measuring Things

- Assume we conduct a random experiment n times. Let S_n be the number of times event A occurs.
- We intuitively assume

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = P[A]$$

Measuring Things

- Assume we conduct a random experiment n times. Let S_n be the number of times event A occurs.
- We intuitively assume

$$\lim_{n \rightarrow \infty} \frac{S_n}{n} = P[A]$$

- We can't “know” $P[A] = p$ precisely because we can only know what we have seen. How can we estimate p ?

Weak Law of Large Numbers

- Assuming each observation is independent, then S_n has a Binomial distribution with $E[S_n] = np$ and $Var[S_n] = np(1 - p)$

Weak Law of Large Numbers

- Assuming each observation is independent, then S_n has a Binomial distribution with $E[S_n] = np$ and $Var[S_n] = np(1 - p)$
- Weak Law of Large Numbers - we can increase the precision of measurements by measuring more:

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \frac{p(1 - p)}{n\delta^2}$$

Weak Law of Large Numbers - II

Mean and variance of the Binomial distribution is:

Weak Law of Large Numbers - II

Mean and variance of the Binomial distribution is:

$$E[S_n] = np$$

$$\text{Var}[S_n] = np(1 - p)$$

Weak Law of Large Numbers - II

Mean and variance of the Binomial distribution is:

$$E[S_n] = np$$

$$\text{Var}[S_n] = np(1 - p)$$

Thus..

$$E[S_n/n] = 1/nE[S_n] = p$$

$$\text{Var}[S_n/n] = 1/n^2\text{Var}[S_n] = \frac{p(1 - p)}{n}$$

- Chebyshev:

$$P[|X - E[X]| \geq t] \leq \frac{\sigma^2}{t^2}$$

- Compute bound for S_n/n :

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \frac{p(1-p)}{n\delta^2}$$

- We can make the R.H.S. arbitrarily small by increasing n (measuring more)
- Thus, we can estimate $P[A]$ from S_n/n .

Weak Law of Large Numbers - How large should n be?

- We saw that


$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \frac{p(1-p)}{n\delta^2}$$

Weak Law of Large Numbers - How large should n be?

- We saw that

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \frac{p(1-p)}{n\delta^2}$$

- Because $0 \leq p \leq 1$, $p(1-p)$ has maximum at $p = 1/2$, or $p(1-p) = 1/4$




Weak Law of Large Numbers - How large should n be?

- We saw that

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \frac{p(1-p)}{n\delta^2}$$

- Because $0 \leq p \leq 1$, $p(1-p)$ has maximum at $p = 1/2$, or $p(1-p) = 1/4$
- We can be certain that no matter what the value of p ,



$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \frac{1}{4n\delta^2}$$

Weak Law of Large Numbers - How large should n be?

- We often want to say “I know that 95% of the time, my estimate will be within δ of the mean”.
- Or, “I am willing to have my estimate be more than δ from the mean no more 5% of the time”

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \epsilon$$

Weak Law of Large Numbers - How large should n be?

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \epsilon$$

This is satisfied if

$$\frac{p(1-p)}{n\delta^2} \leq \epsilon$$

Weak Law of Large Numbers - How large should n be?

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \epsilon$$

This is satisfied if

$$\frac{p(1-p)}{n\delta^2} \leq \epsilon$$

Or, if

$$n \geq \frac{p(1-p)}{(\epsilon\delta^2)}$$

Weak Law of Large Numbers - How large should n be?

$$P\left[\left|\frac{S_n}{n} - p\right| \geq \delta\right] \leq \epsilon$$

This is satisfied if

$$\frac{p(1-p)}{n\delta^2} \leq \epsilon$$

Or, if

$$n \geq \frac{p(1-p)}{(\epsilon\delta^2)}$$

Or, since I may not know what p is (since I'm estimating it),

$$n \geq \frac{1}{4\epsilon\delta^2}$$

Weak Law of Large Numbers - Example

- I'm trying to estimate the probability p of something being busy.

Weak Law of Large Numbers - Example

- I'm trying to estimate the probability p of something being busy.
- Assuming my observations are independent, I want to know how many observations n need to be made such that

$$P\left[\left|\frac{S_n}{n} - p\right| \geq 0.1\right] \leq 0.05$$

- In other words, my estimate will be no more than 0.1 off 95% of the time.

Weak Law of Large Numbers - Example

- I'm trying to estimate the probability p of something being busy.
- Assuming my observations are independent, I want to know how many observations n need to be made such that

$$P\left[\left|\frac{S_n}{n} - p\right| \geq 0.1\right] \leq 0.05$$

- In other words, my estimate will be no more than 0.1 off 95% of the time.
- Since I don't know p then $n \geq 1/(4\epsilon\delta^2)$ says I need

$$n \geq 1/(4 * 0.05 * 0.01) = 500$$

measurements

Weak Law of Large Numbers - Make More Assumptions

- If I know more or make more assumptions, I need fewer measurements

Weak Law of Large Numbers - Make More Assumptions

- If I know more or make more assumptions, I need fewer measurements
- For example if I think $p \approx 0.2$, then I need $n \geq p(1 - p)/(\epsilon\delta^2)$ measurements or

$$n \geq (0.2 * 0.8)/(0.05 * 0.01) = 320$$

measurements

Weak Law of Large Numbers - Make More Assumptions

- If I know more or make more assumptions, I need fewer measurements
- For example if I think $p \approx 0.2$, then I need $n \geq p(1 - p)/(\epsilon\delta^2)$ measurements or

$$n \geq (0.2 * 0.8)/(0.05 * 0.01) = 320$$

measurements

- If we're willing to make more assumptions about our measurements, we can make even fewer measurements as shown by the Central Limit Theorem.