



Statistics: Expectation and Variance Of The Mean

Dirk Grunwald
University of Colorado, Boulder

Measurement and Expectation


- Data is generated from experiments through measurements

Measurement and Expectation

- Data is generated from experiments through measurements
- Those measurements have some error, and scientists and engineers realize that repeated measurements establish more accurate estimates

Measurement and Expectation

- Data is generated from experiments through measurements
- Those measurements have some error, and scientists and engineers realize that repeated measurements establish more accurate estimates
- Running example - Michelson's measurements for speed of light..



Expectation of Data

- Assume X_i are random variables from a source with some distribution with “true” mean μ , variance σ^2 and CDF $F_X()$.

Expectation of Data

- Assume X_i are random variables from a source with some distribution with “true” mean μ , variance σ^2 and CDF $F_X()$.
- The average of the first n random variables in the sequence is..

$$\hat{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Expectation of Data

- Assume X_i are random variables from a source with some distribution with “true” mean μ , variance σ^2 and CDF $F_X()$.
- The average of the first n random variables in the sequence is..

$$\hat{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

- Using linearity of expectation

$$E[\hat{X}_n] = \frac{1}{n}E[X_1 + X_2 + \dots + X_n] = \frac{1}{n}(\mu + \mu + \dots + \mu) = \mu$$

Variance of Data

- The average of the first n random variables in the sequence is..

$$\hat{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

Variance of Data

- The average of the first n random variables in the sequence is..

$$\hat{X}_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$

- Using linearity of expectation $E[\hat{X}_n] = \mu$
- Variance:

$$\text{Var}(\hat{X}_n) = \text{Var}\left(\frac{1}{n} (X_1 + X_2 + \dots + X_n)\right) = \frac{1}{n^2} \text{Var}(\sigma^2 + \dots + \sigma^2) = \frac{\sigma^2}{n}$$

Expectation and Variance of An Average

Definition: Expectation and Variance of An Average

If \hat{X}_n is the average of n independent variables with the same expectation μ and variance σ^2 , then

$$E[\hat{X}_n] = \mu \quad \text{and} \quad \text{Var}(\hat{X}_n) = \frac{\sigma^2}{n}$$

The standard deviation of the average is a factor \sqrt{n} less than single measurement.

Example

Mean and variance of the Binomial distribution.

Let S_n be the sum of n binomial samples with parameter p .

Sample Mean:

$$E\left[\frac{S_n}{n}\right] = p$$

Sample Variance:

$$\text{Var}\left[\frac{S_n}{n}\right] = \frac{np(1-p)}{n^2} = \frac{p(1-p)}{n}$$

Example

Mean and variance of the Binomial distribution.

Let S_n be the sum of n binomial samples with parameter p .

Sample Mean:

$$E\left[\frac{S_n}{n}\right] = p$$

Sample Variance:

$$\text{Var}\left[\frac{S_n}{n}\right] = \frac{np(1-p)}{n^2} = \frac{p(1-p)}{n}$$

Mean and Variance of number of events

$$E[S_n] = np \quad \text{Var}[S_n] = np(1-p)$$