

## Math 637: Homework Chapter 3

1. (3.1.1)

Let  $\varphi : G \rightarrow H$  be a homomorphism and let  $E$  be a subgroup of  $H$ . Prove that  $\varphi^{-1}(E) \leq G$ . If  $E \trianglelefteq H$  prove that  $\varphi^{-1}(E) \trianglelefteq G$ . Deduce that  $\ker \varphi \trianglelefteq G$ .

*Proof.* Let  $a, b \in \varphi^{-1}(E)$ . Then  $\varphi(ab^{-1}) = \varphi(a)(\varphi(b))^{-1}$ . Since  $\varphi(a), \varphi(b) \in E$  and  $E$  is a subgroup, then  $\varphi(a)(\varphi(b))^{-1} \in E$ . Hence  $\varphi^{-1}(E) \leq G$ .

Suppose  $E \trianglelefteq H$  and let  $g \in G$ . Let  $x \in \varphi^{-1}(E)$  so that  $\varphi(x) \in E$ . We will see that  $gxg^{-1} \in \varphi^{-1}(E)$  which is the same as  $\varphi(gxg^{-1}) \in E$ . From this, by Theorem 6 part (5) of chapter 3.1, we will have that  $\varphi(E) \trianglelefteq G$ . Note that  $\varphi(gxg^{-1}) = \varphi(g)\varphi(x)(\varphi(g))^{-1}$  is  $\varphi(x)$  conjugate with  $\varphi(g)$ . By  $E \trianglelefteq H$ , we have  $\varphi(g)\varphi(x)(\varphi(g))^{-1} \in E$ .

To show that  $\ker \varphi \trianglelefteq G$  set  $E = \{1\}$ . This is always normal in any group since  $g\{1\}g^{-1} = \{gg^{-1}\} = \{1\}$ . But then  $\ker \varphi = \varphi^{-1}(E)$  in this case, so the above entails that this is normal in  $G$ .  $\square$

2. (3.1.3)

Let  $A$  be an abelian group and  $B \leq A$ . Prove that  $A/B$  is abelian. Give an example of a non-abelian group  $G$  containing a proper normal subgroup  $N$  such that  $G/N$  is abelian.

*Proof.* First we show that  $A/B$  is a group. As noted in the examples after the definition of the natural projection, any abelian group is normal. From the theorem that all normal subgroups are the kernel of some homomorphism, we then know that  $B = \ker \pi$  for some homomorphism  $\pi$ . Now from the theorem that the quotient group is defined for kernels of homomorphisms, we have that  $A/B$  is a group.

Now let  $x, y \in A$  so that  $xB, yB \in A/B$ . Then

$$(xB)(yB) = (xy)B = (yx)B = (yB)(xB)$$

The first and last equality come from the definition of the quotient group operation. The middle equality is by the abelianness of  $A$ .

Here is an example of a non-abelian group with a proper normal subgroup such that the quotient is abelian: Set  $A = D_4$  and  $B = \langle r \rangle$ , which is to say that  $B$  is the group of rotations. Then to see that  $B \trianglelefteq D_4$  observe that for any  $i = 0, 1, 2, 3$  we have  $r^i B r^{-i} = B$  just because rotations commute with each other. If  $r^i s$  is any reflection, and  $r^j \in B$  then we'll show that  $(r^i s) r^j (r^i s)^{-1} \in B$ , which as we've noted earlier, suffices to show that  $B \trianglelefteq A$ .

$$\begin{aligned} (r^i s) r^j (r^i s)^{-1} &= r^i s r^j s r^{-i} \\ &= r^i r^{-j} s s r^{-i} \\ &= r^{i-j-i} = r^{-j} \in B \end{aligned}$$

Finally, to see that  $A/B$  is abelian it is enough to show that it has order 2. Since

$$sB = s\{1, r, r^2, r^3\} = \{s, sr, sr^2, sr^3\}$$

Since there can be no other cosets, the only two elements of  $A/B$  are  $B$  and  $sB$ . Since trivially any group of order 2 is abelian,  $A/B$  is abelian. □

3. (3.1.11(a))

Let  $F$  be a field and  $G = \left\{ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mid a, b, c \in F, ac \neq 0 \right\} \leq GL_2(F)$ .

(a) Prove that the map  $\varphi : \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \mapsto a$  is a surjective homomorphism from  $G$  onto  $F^\times$ . Describe the fibers and kernel of  $\varphi$ .

*Proof.* First that it's a homomorphism:

$$\begin{aligned} \varphi \left[ \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \right] &= \varphi \begin{pmatrix} ax & ay + bz \\ 0 & cz \end{pmatrix} \\ &= ax \\ &= \left[ \varphi \begin{pmatrix} a & b \\ 0 & c \end{pmatrix} \right] \left[ \varphi \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \right] \end{aligned}$$

Next we see that it's surjective. If  $a \in F$  then  $\varphi \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} = a$ . Note that  $1 \in F$  because  $F$  is a field. □

4. (3.1.22(a))

Prove that if  $H$  and  $K$  are normal subgroups of a group  $G$  then their intersection  $H \cap K$  is also a normal subgroup of  $G$ .

*Proof.* If  $g \in G$  then it suffices to show that  $g(H \cap K)g^{-1} \subseteq H \cap K$ . So let  $gxg^{-1} \in g(H \cap K)g^{-1}$  where  $x \in H \cap K$ . Because  $H$  is normal in  $G$  we have  $gxg^{-1} \in H$ , and because  $K$  is normal in  $G$  we have  $gxg^{-1} \in K$ . Hence  $gxg^{-1} \in H \cap K$ . □

5. (3.1.42)

Assume both  $H$  and  $K$  are normal subgroups of  $G$  with  $H \cap K = 1$ . Prove that  $xy = yx$  for all  $x \in H$  and  $y \in K$ .

*Proof.* We first prove  $x^{-1}y^{-1}xy \in H$ . We already know  $x \in H$  therefore  $x^{-1} \in H$ . Because  $H \trianglelefteq G$  then  $y^{-1}x(y^{-1})^{-1} \in H$  since this is conjugation of  $x$  by  $y^{-1}$ . Then  $x^{-1}y^{-1}xy \in H$ .

Next we show  $x^{-1}y^{-1}xy \in K$ . We already know  $y \in K$ . Because  $K \trianglelefteq G$  then  $x^{-1}y(x^{-1})^{-1} \in K$  since this is conjugation of  $y$  by  $x^{-1}$ . Then  $x^{-1}y^{-1}xy \in K$ .

Hence  $x^{-1}y^{-1}xy \in H \cap K$  and therefore  $x^{-1}y^{-1}xy = 1$ . Multiply by  $yx$  and you have

$$xy = yx$$

as desired. □

6. (3.2.4)

Show that if  $|G| = pq$  for some primes  $p$  and  $q$  (not necessarily distinct) then either  $G$  is abelian or  $Z(G) = 1$ .

*Proof.* If  $Z(G) \neq 1$  then the order  $|Z(G)|$  can only be either  $p$  or  $q$  or  $pq$ . If the order is  $pq$  then  $Z(G) = G$  and so  $G$  is abelian. So it suffices to show that the order cannot be either  $p$  or  $q$ .

Without loss of generality, suppose for contradiction that the order of  $Z(G)$  is  $p$ . That is to say  $|Z(G)| = p$ . Let  $g \in G \setminus Z(G)$  so that  $g$  does not commute with some element. As a preliminary remark note that no non-identity element of  $\langle g \rangle$  is in  $Z(G)$  either, since if  $g^i$  commutes with every  $x \in G$  then

$$g^i x = x g^i \Leftrightarrow$$

$$g x g^{i-1} = x g^i \Leftrightarrow$$

$$g x = x g$$

So  $g$  commutes with every  $x \in G$  contrary to assumption. Further, this implies that the order of the group generated by both sets,  $|\langle Z(G), g \rangle|$  is  $|Z(G)||g|$ . This is because for every  $z_1, z_2 \in Z(G)$  and  $h_1, h_2 \in \langle g \rangle$ , the elements  $z_1 h_1 = z_2 h_2$  if and only if  $z_2^{-1} z_1 = h_2 h_1^{-1} \in \langle g \rangle \cap Z(G)$ . So in that case  $z_1 = z_2$  and  $h_1 = h_2$ . Thus every  $z \in Z(G), h \in \langle g \rangle$  corresponds to a unique  $zh \in \langle Z(G), g \rangle$ . The number of such choices is  $|Z(G)||g|$ .

Now the order of  $g$  is  $p$  or  $q$  or  $pq$ . If the order is  $pq$  then  $G$  is cyclic, so abelian, so  $|Z(G)| = pq$  contrary to assumption.

Suppose the order of  $g$  is  $q$ . Thus  $|\langle Z(G), g \rangle| = pq$ . Hence  $G = \langle Z(G), g \rangle$ . Thus if we pick any  $a, b \in G$  then by the observations above,  $a$  must have the form  $z_a h_a$  for some  $z_a \in Z(G)$  and  $h_a \in \langle g \rangle$ . Likewise  $b = z_b h_b$  for some  $z_b \in Z(G)$  and  $h_b \in \langle g \rangle$ . Now  $z_a$  and  $z_b$  commute with everything in  $G$  by definition.  $h_a$  and  $h_b$  commute with everything in  $\langle g \rangle$  trivially. So

$$ab = z_a h_a z_b h_b = z_a z_b h_a h_b = z_a z_b h_b h_a = z_b h_b z_a h_a = ba$$

This shows that  $G$  is abelian, contrary to the assumption that  $|Z(G)| = p < pq$ .

Finally suppose that  $|g| \neq q$  so that  $|g| = p$ . Notice that these two facts entail  $p \neq q$ . Also  $|\langle Z(G), g \rangle| = p^2$ , so  $p^2 | pq$ , so  $p | q$ , which is impossible for distinct primes.  $\square$

7. (3.2.7)

Let  $H \leq G$  and let  $g \in G$ . Prove that if the right coset  $Hg$  equals some left coset of  $H$  in  $G$ , then it equals the left coset  $gH$  and that  $g \in N_G(H)$ .

*Proof.* Suppose  $Hg = xH$  for some  $x \in G$ . Since  $xH = gH$  is equivalent to  $g^{-1}x \in H$ , we make this our goal. First note that  $x \in xH$  since  $1 \in H$ , and hence  $x \in Hg$ . So there exists some  $h \in H$  such that  $x = hg$  and we get  $g^{-1}x = h \in H$  as desired.

From  $Hg = gH$  we can infer  $H = gHg^{-1}$  so that  $g \in N_G(H)$  by definition.  $\square$

8. (3.2.12)

Let  $H \leq G$ . Prove that the map  $x \mapsto x^{-1}$  sends each left coset of  $H$  in  $G$  onto a right coset of  $H$ , and gives a bijection between the set of left cosets and the set of right cosets of  $H$  in  $G$  (hence the number of left cosets equals the number of right cosets).

*Proof.* Let  $gH$  be any left coset and let  $gh \in gH$  be any element. Then  $gh \mapsto (gh)^{-1} = h^{-1}g^{-1}$ . Since  $h^{-1} \in H$  then  $h^{-1}g^{-1} \in Hg^{-1}$ . Thus the map sends every element of  $gH$  to some element of  $Hg^{-1}$ .

To see that this map is onto, let  $hg^{-1} \in Hg^{-1}$ . Since  $h^{-1} \in H$  then  $gh^{-1} \in gH$  and the map sends this to  $(gh^{-1})^{-1} = hg^{-1}$ , as desired.

Now we use this to build a bijection between the set of left cosets,  $G/H$ , and the set of right cosets,  $H \backslash G$ . Namely, let  $\varphi(g_iH) = Hg_i^{-1}$ . We first need to see that this map is well-defined, then that it's a bijection. To see that it's well-defined suppose that  $g_iH = g_jH$  and note that this is equivalent to  $g_j^{-1}g_i \in H$ . Therefore we have  $(g_j^{-1}g_i)^{-1} \in H$ , which is the same as  $g_i^{-1}g_j \in H$ . But this entails  $Hg_i^{-1}g_j = H$  so that  $Hg_i^{-1} = Hg_j^{-1} = \varphi(g_iH) = \varphi(g_jH)$ , and this shows  $\varphi$  is well-defined.

Next we show that  $\varphi$  is injective. If

$$\varphi(g_iH) = \varphi(g_jH) = Hg_i^{-1} = Hg_j^{-1}$$

then

□

9. (3.2.16)

*Proof.*

□

10. (3.3.3)

*Proof.*

□

11. (3.3.8)

*Proof.*

□

12. (3.3.9)

*Proof.*

□

13. (3.4.1)

*Proof.*

□

14. (3.4.11)

*Proof.*

□

15. (3.5.12)

*Proof.*

□