

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/271101950>

# Ultraviolet Pasteurization for Food Industry

Article in *International Journal of Food Science and Nutrition Engineering* · February 2012

DOI: 10.5923/j.food.20120201.03

CITATIONS

38

READS

3,562

2 authors, including:



**Ruplal Choudhary**

Southern Illinois University Carbondale

47 PUBLICATIONS 1,091 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Non thermal food processing [View project](#)



Food Safety [View project](#)

# Ultraviolet Pasteurization for Food Industry

Ruplal Choudhary\*, Srinivasarao Bandla

Department of Plant, Soil and Agricultural Systems, Southern Illinois University, Carbondale, IL 62901, USA

**Abstract** This article reviews the status and scope of ultraviolet (UV) light technology in food processing industry for control of foodborne pathogens and spoilage organisms for food safety and shelf life extension. The literature suggests that there have been extensive studies on application of ultraviolet light for disinfection of apple ciders. FDA acceptable reduction of pathogens and spoilage organisms were reported by UV treatment of for apple cider. Several recent studies on UV treatment of milk suggest that there is more work needed to improve design of UV reactors for raw milk processing. Fresh produce industry will also benefit from UV processing so that occurrence of frequent outbreaks can be avoided. A brief review of recent applications in produce industry has also been reported. Finally a recommendation was made for future direction of UV application research in food processing industry.

**Keywords** Ultraviolet processing, nonthermal processing, food processing, UV, UV-C

## 1. Introduction

Deterioration of foods by pathogenic and spoilage microorganisms can be minimized by food processing and preservation techniques. Frequent outbreaks of foodborne pathogens are associated with fresh produce, milk and fruit juices. Currently, produce industry uses chemicals and fumigants to control pathogens in packing and handling industries. The liquid foods and dairy industry uses thermal pasteurization methods to disinfect food pathogens and extend the shelf-life of products. Though thermal methods are effective in inactivating microorganisms, thermal processing negatively affects foods through losses in vitamins; and changes in sensory properties, such as color, flavor and wholesomeness (Montenegro et al. 2002). Moreover, thermal processing is not practical in fresh produce industry. Thermal pasteurization also affects enzyme inactivation, lipid oxidation, protein denaturation and non-enzymatic browning. Therefore nonthermal processing methods are being studied as alternative food processing techniques. Currently several nonthermal technologies are under research. Some of these technologies are ultrasound (US), high-pressure processing (HPP), pulsed electric fields (PEF), pulsed light treatment (PL), ultraviolet light (UV), and non-thermal atmospheric pressure plasma (NTAP). These novel non-thermal methods may be useful in inactivating foodborne pathogens and spoilage microorganisms from a range of solid and liquid foods (Montenegro et al. 2002; Pereira and Vicente 2009; Song et al. 2009; Ekem and Akan 2006; Purevdorj et al. 2002; Critzer et al. 2007; Deng et al.

2007). UV irradiation is considered one of the effective means of disinfection, which excludes the necessity of heat to get rid of microorganisms (Sastry et al. 2008).

## 2. Ultraviolet Light

There are three regions of ultraviolet light within the electromagnetic spectrum. These are the UV-A (315–400 nm), UV-B (280–315 nm) and UV-C (200–280 nm). UV light exhibits germicidal properties in the UV-C region. The inactivation efficiency follows a bell-shaped curve where maximum inactivation occurs approximately at the range of 254 to 264 nm. However, typical mercury UV lamps deliver at 254 nm maximum. Therefore, it is usually mentioned that UV inactivation is at 254 nm. Several studies suggest that the destruction of microorganisms occurs due to the penetration of UV-C light into the outer membranes of the cells leading to tremendous damage of the DNA owing to the formation of thymine dimers, which prevent the microorganism from undertaking DNA transcription and replication, eventually leading to cell death in the UV-C light disinfection process (Bank et al. 1990; Bintsis et al. 2000; Miller et al. 1999).

## 3. Penetration and Absorption of Ultraviolet Light in Food

UV light penetrates food materials only up to several millimeters depending upon the optical properties of the food. UV light can easily penetrate water since it is transparent to the wavelengths produced. UV light cannot penetrate milk and other turbid foods as well, so opaque foods need to be presented to the system as a thin layer. Guerrero-Beltran and Barbosa-Canovas (2004) stated that the color or the turbidity

\* Corresponding author:

choudhry@siu.edu (Ruplal Choudhary)

Published online at <http://journal.sapub.org/food>

Copyright © 2012 Scientific & Academic Publishing. All Rights Reserved

of the liquid influences the optical absorption coefficient of the liquid. The penetration capacity of the UV light reduces as the absorption coefficient increases. Therefore, it is essential to understand that enhancing the penetrative depth will be beneficial for UV light treatment of foods with higher coefficients of absorption. Presentation of food in a thin film may result in an increase in the efficiency of microorganism inactivation.

#### 4. Effect of UV on Food Components and Quality

UV light exposure initiates free radical oxidation and catalyzes other stages of the oxidation process. Lipid radicals, superoxide radicals (SOR), and H<sub>2</sub>O<sub>2</sub> are formed due to UV light (Kolakowska 2003). SOR can further induce carbohydrate cross linking, protein cross linking, protein fragmentation, peroxidation of unsaturated fatty acid, and loss of membrane fluidity function. Denaturation of components such as proteins, enzymes, and amino acids (especially amino acids with aromatic compounds) in milk may occur with UV radiation, thereby also bringing about textural changes. Water also absorbs UV photons and produces OH- and H<sup>+</sup> radicals, which in turn aids changes in other food components. Kolakowska (2003) reported that there are obvious changes in the chemical composition of food components and product quality deterioration when the UV light treatment is applied in high doses. Therefore, it is mandatory to properly optimize the disinfection process so that the quality of the food products is maintained and its safety is ensured. Normally, microbial inactivation can be achieved within seconds to minutes depending upon the opacity of the food products and microorganism type.

In general, using UV light treatment for food has been found not to cause any adverse effects, especially if UV light is applied in moderate amounts (Krishnamurthy 2006). However, the modification and optimization of the UV light treatment might be necessary for successful implementation of the process with regard to some foods.

#### 5. UV Disinfection of Liquid Foods

New regulations imposed by FDA stipulate the implementation of Hazard Analysis and Critical Control Point (HACCP) in fruit and vegetable juice processing operations (66 FR 6137). This regulation requiring 5-log reduction of pathogens has necessitated the development of affordable technology. The efficacy of UV irradiation to control pathogens in apple cider has been the focus of recent studies (Hanes et al. 2002; Wright et al. 2000; Basaran et al. 2004).

Wright and colleagues (2000) inoculated apple cider with a cocktail of five *Escherichia coli* O157:H7 strains to an approximate level of 10<sup>6</sup> CFU/ml and placed the stained apple cider in thin films through the Cider-10uv model (Ideal Horizons, Poultney, VT) UV disinfection unit through an

UV radiation at 254 nm (Wright et al. 2000). The flow rates ranged from 60 to 90 l/h to generate UV doses between 9.4 and 61 mJ/cm<sup>2</sup>. The mean log reduction was 3.8 log CFU/ml (Wright et al. 2000). They suggested that if the apparatus was modified to increase the intensity of UV irradiation in addition to a maximal flow rate, *E. coli* reduction would be higher and at faster rates. The levels of yeast and mold counts in the cider also influenced the reduction of microbial counts (Wright et al. 2000).

Choi and Nielsen (2005) demonstrated that UV pasteurized apple cider were superior in color and overall sensory scores compared to thermally pasteurized apple cider. UV-irradiated samples were lower in soluble solids for the first 7 days and showed no significant difference in consumer acceptability (Choi and Nielsen 2005).

Studies using apple cider that utilized the CiderSure 3500 UV apparatus (FPE, Inc., Rochester, NY) confirmed the ability of the apparatus to achieve a 5-log reduction of *Cryptosporidium parvum* and *E. coli* (Hanes et al. 2002; Basaran et al. 2004). The apparatus was designed to allow the apple cider to pass through a series of eight germicidal UV lamps in thin films. The wavelength and intensity inside the CiderSure apparatus was 254 nm and 14.3 mJ/cm<sup>2</sup> of UV irradiation, respectively, with exposure times between 1.2 and 1.9 sec (Hanes et al. 2002). A computer-monitored UV sensor was placed within the apparatus to adjust the flow rate in accordance with the sensor readings to ensure that all of the cider received the correct amount of UV light.

In a study conducted by Basaran et al. (2004), ciders with different solid compositions and concentrations were subjected to different filtration treatments that produced lighter and darker ciders. These variables were deliberately included to test the UV apparatus for its ability to overcome such differences and still achieve a 5-log reduction in *E. coli* O157:H7. Three strains of *E. coli* O157:H7 were used in the study (ATCC 43889, 43895, and 43933). The apple cider samples were inoculated with each of the three strains and then passed through the CiderSure UV-irradiation unit at 4°C. The results showed a 6.12 ± 0.36, 5.83 ± 0.11, and 5.87 ± 0.11 log reduction for the three *E. coli* O157:H7 strains, respectively (Basaran et al. 2004). They also found >5 log reduction of a surrogate organism *E. coli* ATCC 25922 in a validation test at a cider mill production setting. Quintero-Ramos et al. (2004) showed that apple cider's pH did not have a significant influence on reduction of *E. coli* 25922 in apple cider. The results showed existence of a nonlinear relationship between *E. coli* survival rate and UV dose.

Koutchama et al. (2004) studied the efficacy of UV light on the destruction of *E. coli* K-12 in apple juice using laminar and turbulent flow UV reactors. They concluded that increase in flow rate of UV radiation increases the inactivation of *E. coli* in apple juice when turbulent flow UV reactors are used because of better mixing conditions inside the UV reactor. On examining the physical and chemical parameters of the apple juices, they found that absorbance consistently affected inactivation of *E. coli* K 12 in the apple juice.

Matak (2004) studied the efficacy of UV light in inacti-

vating *E. coli* K 12 for different fat percentages of milk at different temperatures (4°C and 20°C) using CiderSure 3500 apparatus. They found that a 0.73 to 2.29 log reduction of *E. coli* in different fat percentages of milk. *Listeria monocytogenes* was inactivated to non-detectable levels from 107 CFU/ml in goat milk when UV light at cumulative UV dose of  $15.8 \pm 1.6$  mJ/cm<sup>2</sup> was used for an exposure time of 18 sec (Re- 5187) (Matak *et al.* 2005).

Reinemann *et al.* (2006) achieved 3-log reduction of natural flora in raw cow milk with UV dose of 1.5 J/ml using UV reactors pure version 1 and 2 in their laboratory. These reactors contain low pressure mercury UV lamp inside the quartz glass sleeve and this enclosed in a stainless steel chamber.

Most recently our group (Choudhary *et al.*, 2011a; Bandla *et al.*, 2012) examined efficiency of Dean flow UV reactors on inactivation of *E. coli* W 1485 and *Bacillus cereus* endospores in raw cow milk, commercially processed skimmed milk, and soymilk. We found >7 log reduction of *E. coli* W 1485 in skimmed milk and >5 Log reduction of the same organism for soymilk using a Dean flow reactor with 1.6 mm diameter with a UV dose of 0.05 J/ml. With raw cow milk using the same reactor at the same UV dose, we found a 4 log reduction of *E. coli* W 1485. A higher UV dose for raw cow milk than skimmed milk and soymilk was recommended due to the lower UV transmission of raw cow milk.

## 6. UV Treatment of Fresh Produce

There are several non-thermal pathogen reduction techniques for fresh produce. These alternative treatment processes may be chemical or non-chemical in nature. Most chemical treatment techniques include chlorine, bromine, iodine, trisodium phosphate, hydrogen peroxide, or ozone. Non-chemical treatments include microwave, ultrasound, ionizing radiation and ultraviolet (UV). UV is a promising technology of surface decontamination because it is safe and does not leave any residual effect in treated food products. In addition to being germicidal, UV treatments have been found to induce desirable changes in health constituents of fruits and vegetables such as increased antioxidant capacity and increased shelf life (Wang *et al.*, 2009). UV systems are affordable as they require low initial investment and a lower operating cost of treatment (Yuan *et al.* 2004).

Several studies on UV treatment of fruits and vegetables were conducted for increasing shelf life and improving product quality (Lu *et al.* 1991; Bialka and Demirci 2007, Pombo *et al.* 2009). Lu *et al.* (1991) studied the effect of low levels of UVC radiation (130-4000 mJ/cm<sup>2</sup>) on the shelf life of peaches and tomatoes, and reported reduced post-harvest rots and delayed ripening.

Bialka and Demirci (2007) reported using UV treatments for decontamination of *E. coli* and *Salmonella enterica* on blueberries. After 60 s of pulsed UV treatment, they reported a maximum reduction of 4.3 and 2.9 log CFU/g for *Salmonella* and *E. coli* respectively. Pulsed UV is more expensive

than continuous-wave UV. The low initial cost of continuous-wave UV as well as the lack of extensive safety equipment may benefit those with little capital to invest, which applies to most commercial blueberry packinghouses.

## 7. Conclusions

Promising opportunity exists for adopting ultraviolet processing in a small or large scale food and dairy processing industry. With the approval of FDA, several new applications of UV processing are being tested and validated by the dairy and food industries in the United States of America. With potential for offering superior organoleptic qualities of food products at lower initial investment and operating costs, the authors foresee a great success for adoption of UV processing technology by the food processing industry.

## ACKNOWLEDGEMENTS

The authors acknowledge the support of Department of Plant, Soil and Agricultural Systems, College of Agricultural Sciences, Southern Illinois University, Carbondale, IL.

## REFERENCES

- [1] Bandla S., R. Choudhary, D.W. Watson, J. Haddock. (2012). UV-C treatment of soymilk in coiled tube UV reactors for inactivation of *Escherichia coli* W1485 and *Bacillus cereus* endospores. *LWT-Food Science and Technology* 46(1) 71-76.
- [2] Bank H. L., J. L. Schmehl, and R. J. Dratch. (1990). Bacteriocidal effectiveness of modulated UV light. *Applied Environmental Microbiology* 56: 3888-89.
- [3] Basaran N., A. Quintero-Ramos, M.M. Moake, J.J. Churey and R.W. Worobo (2004). Influence of apple cultivars on inactivation of different strains of *Escherichia coli* O157:H7 in apple cider by UV irradiation. *Applied Environmental Microbiology* 70: 6061-6065.
- [4] Bialka K.L and A. Demirci (2007). Decontamination of *Escherichia coli* O157:H7 and *Salmonella enterica* on blueberries using ozone and pulsed UV-light. *Journal of Food Science* 72(9): M931-M936.
- [5] Bintsis T., E. Litopoulou-Tzanetaki, and R. Robinson. (2000). Existing and potential applications of the ultraviolet light in the food industry- A critical review. *Journal of the Science of Food and Agriculture* 80: 637-45.
- [6] Choi L. H. and S. S. Nielsen (2005). The effects of thermal and nonthermal processing methods on apple cider quality and consumer acceptability. *Journal of Food quality* 28: 13-29.
- [7] Choudhary, R., S. Bandla, D.G. Watson, J. Haddock, A. Abughazaleh and B. Bhattacharya (2011). Performance of coiled tube ultraviolet reactors to inactivate *Escherichia coli* W1485 and *Bacillus cereus* endospores in raw cow milk and

commercially processed skimmed cow milk. *Journal of Food Engineering* doi:10.1016/j.jfoodeng.2011.06.009

- [8] Critzer, F. J., K. K. Winterberg, S. L. South, and D. A. Golden. (2007). Atmospheric plasma inactivation of foodborne pathogens on fresh produce surfaces. *Journal of Food Protection* 70: 2290–96.
- [9] Deng S., R. Ruan R, K. C. Mok, G. Huang, X. Lin, and P. Chen. (2007). Inactivation of *Escherichia coli* on almonds using nonthermal plasma. *Journal of Food Science* 72: M62–66.
- [10] Ekem N., and T. Akan. (2006). Sterilization of *Staphylococcus aureus* by atmospheric pressure pulsed plasma. *Surface Coating and Technology* 201:993–97.
- [11] Guerrero-Beltran, J.A., G.V. Barbosa-Canovas. (2004). Review: Advantages and limitations on processing foods by UV light. *Food Science and Technology International* 10(3):137-147.
- [12] Hanes, D. E., P. A. Orlandi, D. H. Burr, M. D. Miliotis, M. G. Robi, J. W. Bier, G. J. Jackson, M. J. Arrowood, J. J. Churey, and R. W. Worobo. (2002). Inactivation of *Cryptosporidium parvum* oocysts in fresh apple cider using ultraviolet irradiation. *Applied Environmental Microbiology* 68:4168–72.
- [13] Kolakowska A. (2003). Lipid oxidation in food systems. In *Chemical and functional properties of food lipids*, 133–168. New York: CRC Press.
- [14] Koutchama T., S. Keller S, S. Chirtel S, and B. Parisi. (2004). Ultraviolet disinfection of juice products in laminar and turbulent flow reactors. *Innovative Food Science and Emerging Technologies* 5:179–89.
- [15] Krishnamurthy, K. (2006). Decontamination of milk and water by pulsed UV-light and infrared heating. PhD thesis. Pennsylvania State University, chapters 2, pp. 23–44.
- [16] Lu J., Y.C. Stevens, V. A. Khan, and M. Kabwe. (1991). The effect of ultraviolet irradiation on shelf- life and ripening of peaches and apples. *Journal of Food Quality* 14 (4): 299-305.
- [17] Matak, K. E. (2004). Effects of UV irradiation on the reduction of bacterial pathogens and chemical indicators of milk. PhD thesis. Virginia Polytechnic Institute and State University.
- [18] Matak, K.E., J. J. Churey, R. W. Worobo, S. S. Sumner, E. Hovingh, C. R. Hackney, and M. D. Pierson. (2005). Efficacy of UV light for the reduction of *Listeria monocytogenes* in goat's milk. *Journal of Food Protection* 68:2212–16.
- [19] Miller R., W. Jeffrey, D. Mitchell, and M. Elasri. (1999). Bacterial responses to ultraviolet light. *ASM News* 65:534–41.
- [20] Montenegro, J., R. Ruan, H. Ma, and P. Chen. (2002). Inactivation of *E. coli* O157:H7 using a pulsed nonthermal plasma system. *Journal of Food Science* 67:646–48.
- [21] Pereira, R. N., and A. A. Vicente (2009). Environmental impact of novel thermal and non-thermal technologies in food processing. *Journal of Food Research* 1–8.
- [22] Pombo M.A., M.C. Dotto, G.A. Martinez, and P.M. Civello. (2009). UV-C irradiation delays strawberry fruit softening and modifies the expression of genes involved in cell wall degradation. *Postharvest Biology and Technology* 51:141-148.
- [23] Purevdorj D., N. Igura, I. Hayakawa, and O. Ariyada. (2002). Inactivation of *Escherichia coli* by microwave induced low temperature argon plasma treatments. *Journal of Food Engineering* 53: 341–46.
- [24] Quintero-Ramos, A., J. J. Churney, P. Hartman, J. Barnard, and R. W. Worobo. (2004). Modeling of *Escherichia coli* inactivation by UV irradiation at different pHs in apple cider. *Journal of Food Protection* 67(6):1153–56.
- [25] Reinemann, D.J., P. Gouws, T. Cilliers, K. Houck, and J. R. Bishop. (2006). New methods for UV treatment of milk for improved food safety and product quality. ASABE presentation. Paper No. 066088.
- [26] Sastry, S. K., A. K. Datta, and R. W. Worobo. (2008). Ultraviolet light. *Journal of Food Science Supplement* 90–2.
- [27] Song, P. H., B. Kim, H. J. Choe, S. Jung, Y. S. Moon, and W. Choe (2009). Evaluation of atmospheric pressure plasma to improve the safety of sliced cheese and ham inoculated by 3-strain cocktail *Listeria monocytogenes*. *Journal of Food Microbiology* 26:432–36.
- [28] Wang, C. Y., C. T. Chen, and S.Y. Wang. (2009). Changes of flavonoid content and antioxidant capacity in blueberries after illumination with UV-C. *Food Chemistry*. Accepted manuscript: doi:10.1016/j.foodchem.2009.04.037
- [29] Wright, J. R., S. S. Sumner, C. R. Hackney, M. D. Pierson, and B. W. Zoecklein. (2000). Efficacy of ultraviolet light for reducing *Escherichia coli* O157:H7 in unpasteurized apple cider. *Journal of Food Protection* 63:563–67.
- [30] Yaun B.R, Sumner S. S., Eifert J.D., Marcy J.E. (2004). Inhibition of pathogens on fresh produce by ultraviolet energy. *International Journal of Food Microbiology* 90 (1): 1-8.