Pharmacovigilance Internship – Task 1

ADR Identification Basics

1. Objective

To identify potential adverse drug reactions by analysing patient symptoms, medication history and the timeline of drug administration.

Adverse Drug Reactions

Adverse drug reaction may be defined as any response to a drug that is noxious and unintended and that occurs at doses normally used in humans for prophylaxis, diagnosis or therapy of disease, or for the modification of physiological function

2. Patient Information

- Patient: Female
- Age: 45 years
- Symptoms: Rash, itching, nausea and dizziness.

3. Medication History

Medicine	Medication history
Amoxicillin	Started 5 days ago
Paracetamol	Taken as needed

4. Suspected Adverse Drug Reaction

The patient developed rash, itching, nausea and dizziness after starting amoxicillin.

Amoxicillin is a penicillin-class antibiotic that can cause hypersensitivity reactions, including skin rash and itching.

5. Timeline Analysis

- Day -5: Amoxicillin started.
- Day -2: Symptoms appeared.
- Day 0: Symptoms are being evaluated.

6. Suspected Drug

Amoxicillin is the primary suspected drug because the symptoms developed after its administration and rash and itching are recognised adverse reactions.

Paracetamol should also be considered as a possible contributing drug.

7. Assessment

The available information suggests a possible drug-related reaction. However, additional details about previous allergies, exact doses, symptom progression and response to drug withdrawal are needed to establish causality.

8. Conclusion

A suspected adverse drug reaction associated with amoxicillin has been identified. Early identification and reporting of ADRs help improve patient safety and rational medicine use.

9. Learning Outcome

- How to review patient symptoms
- What is ADR
- How to report an adverse drug reaction
- Types of Adverse Drug Reactions
- Pharmacovigilance