

Research Article

Deep RetinaNet for Dynamic Left Ventricle Detection in Multiview Echocardiography Classification

Meijun Yang,¹ Xiaoyan Xiao,² Zhi Liu^{1,3}, Longkun Sun,⁴ Wei Guo,³ Lizhen Cui,³ Dianmin Sun,⁵ Pengfei Zhang⁴, and Guang Yang⁶

¹School of Information Science and Engineering, Shandong University, Qingdao 266237, China

²Department of Nephrology, Qilu Hospital, Cheeloo College of Medicine, Shandong University, No. 107 Wenhua Road, Jinan 250012, China

³Joint SDU-NTU Centre for Artificial Intelligence Research (C-FAIR), Shandong University, Jinan 250101, China

⁴Department of Cardiology, Qilu Hospital, Cheeloo College of Medicine, Shandong University, No. 107 Wenhua Road, Jinan, Shandong Province, China

⁵Department of Thoracic Surgery, Shandong Cancer Hospital and Institute, Shandong First Medical University and Shandong Academy of Medical Sciences, Jinan 250117, China

⁶National Heart and Lung Institute, Imperial College London, SW7 2AZ, London, UK

Correspondence should be addressed to Zhi Liu; liuzhi@sdu.edu.cn and Pengfei Zhang; pengf-zhang@163.com

DW[hW #* 8WdgSk \$ " \$ - DM[eW % # ? Sk \$ " \$ - 3UWfW \$ + - g ` W\$ " \$ - BgT'feZW # 3gYgef \$ " \$ "

Academic Editor: Chenxi Huang

Copyright © 2020 Meijun Yang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background. Currently, echocardiography has become an essential technology for the diagnosis of cardiovascular diseases. Accurate classification of apical two-chamber (A2C), apical three-chamber (A3C), and apical four-chamber (A4C) views and the precise detection of the left ventricle can significantly reduce the workload of clinicians and improve the reproducibility of left ventricle segmentation. In addition, left ventricle detection is significant for the three-dimensional reconstruction of the heart chambers. **Method.** RetinaNet is a one-stage object detection algorithm that can achieve high accuracy and efficiency at the same time. RetinaNet is mainly composed of the residual network (ResNet), the feature pyramid network (FPN), and two fully convolutional networks (FCNs); one FCN is for the classification task, and the other is for the border regression task. **Results.** In this paper, we use the classification subnetwork to classify A2C, A3C, and A4C images and use the regression subnetworks to detect the left ventricle simultaneously. We display not only the position of the left ventricle on the test image but also the view category on the image, which will facilitate the diagnosis. We used the mean intersection-over-union (mIOU) as an index to measure the performance of left ventricle detection and the accuracy as an index to measure the effect of the classification of the three different views. Our study shows that both classification and detection effects are noteworthy. The classification accuracy rates of A2C, A3C, and A4C are 1.000, 0.935, and 0.989, respectively. The mIOU values of A2C, A3C, and A4C are 0.858, 0.794, and 0.838, respectively.

1. Introduction

Heart disease is a common circulatory disease that not only seriously affects the function of the cardiovascular system but also causes certain damage to the respiratory system [1]. In severe cases, heart failure can endanger life [2]. Therefore, it is important to use more advanced methods to observe cardiac symptoms or exercise [3]. In the clinical diagnosis of heart disease, echocardiography is the most commonly used tool [4]. Echocardiography is a noninvasive technique for

examining the anatomy and functional status of the heart and large blood vessels using ultrasound [5]. It uses pulsed ultrasound to measure the periodic activities of the underlying walls, ventricles, and valves through the chest wall and soft tissues [6].

The left ventricle is the focal part of the heart, and the symptoms of the left ventricle are an important basis for the diagnosis of heart disease [7]. Therefore, accurate information regarding the left ventricle extracted from echocardiography is crucial for further clinical procedures and

