

# Atrial Scar Segmentation from 3D Late Gadolinium Enhanced Datasets: Effect of Time After Contrast Injection

Guang Yang, Jun Chen, Heye Zhang, Rick Wage, Jack Allen, Tom Wong, Raad Mohiaddin, David Firmin, Jennifer Keegan

**Background:** Late gadolinium enhancement (LGE) imaging enables visualization and quantification of native and ablation induced atrial scars. A recent study reported a marked increase in measured atrial scar percentage (AS%) in data acquired at different times after contrast agent injection (median [IQR]: 17.9% [11.7%–28.5%] at 20 mins vs. 24.1% [19.2%–35.2%] at 30 mins [1]) as the blood-scar contrast increased. This reflects issues with the analysis technique, which determined scar as a fixed threshold above blood signal intensity. We have used a newly developed deep learning-based method to analyze AS% at two different time points after gadolinium administration. We hypothesized that this fully automated method would be less dependent on the contrast characteristics and that the measured AS% would be more reproducible.

**Methods:** Transverse navigator-gated 3D LGE imaging [2][3] was performed in 20 atrial fibrillation (AF) patients (4 pre-ablation and 16 three months post-ablation) using an inversion prepared segmented gradient echo sequence (TE/TR=2.2/5.2ms). Imaging was performed 20 mins after gadolinium administration (Scan 1) and then repeated immediately (~30 mins post-injection, depending on the respiratory efficiency) (Scan 2). A joint multiview two-task (MVTT) segmentation method [4] which had been trained on 184 previously acquired datasets (20 mins after post-injection) was used to simultaneously segment both the left atrium (LA) and the AS tissue in these 40 datasets and AS% ( $100 \times \text{scar volume} / \text{LA volume}$ ) was calculated. Segmentation time was ~0.3 seconds per 3D dataset on a state-of-the-art GPU. Intraclass correlation coefficient of the AS% and apparent contrast-to-noise (aCNR) ratios were also evaluated.

**Results:** The mean ( $\pm$ SD) acquisition times of Scans 1 and Scans 2 were  $17.6\pm 1.3$ mins and  $27.6\pm 3.3$ mins post injection, respectively. The aCNR was significantly lower in Scan 1 than Scan 2 (median [IQR]: 14.10 [10.36–19.30] vs. 18.42 [12.53–24.42],  $P<0.05$ ). There was a small change in measured AS% with time after contrast administration (median [IQR]: Scan 1–24.11% [21.26%–26.82%] vs. Scan 2–21.34% [18.74%–22.65%]) as shown in Figure 1 but this did not reach statistical significance (paired Wilcoxon rank sum test,  $P=0.0531$ ). Visualization of the delineated scar demonstrated good subjective agreement between scans at the two timepoints (example cases shown in Figure 2). The AS% intraclass correlation coefficient (absolute agreement) was 0.901.

**Conclusion:** Our fast fully automatic MVTT model results in AS% measurements that are relatively independent of the image contrast changes that occur with time after contrast administration. Further studies using multi-center and multi-scanner datasets with different acquisition protocols are underway.

## Reference

- [1] H. Chubb *et al.*, “Optimization of late gadolinium enhancement cardiovascular magnetic resonance imaging of post-ablation atrial scar: a cross-over study.,” *J. Cardiovasc. Magn. Reson.*, vol. 20, no. 1, p. 30, 2018.
- [2] R. S. Oakes *et al.*, “Detection and quantification of left atrial structural remodeling with delayed-enhancement magnetic resonance imaging in patients with atrial fibrillation,” *Circulation*, vol. 119, no. 13, pp. 1758–1767, 2009.
- [3] D. C. Peters *et al.*, “Recurrence of Atrial Fibrillation Correlates With the Extent of Post-Procedural Late Gadolinium Enhancement. A Pilot Study,” *JACC Cardiovasc. Imaging*, vol. 2, no. 3, pp. 308–316, 2009.
- [4] G. Yang *et al.*, “Simultaneous left atrium anatomy and scar segmentations via deep learning in multiview information with attention,” *Futur. Gener. Comput. Syst.*, vol. 107, pp. 215–228, Jun. 2020.




Figure 1: Measured scar percentage calculated in 20 patients scanned 20 mins (Scan 1, green) and 30 mins (Scan 2, orange) after gadolinium administration.




Figure 2: Visualization of the delineated LA and AS for two AF patients scanned 20 mins (Scan 1) and 30 mins (Scan 2) after the gadolinium administration.