

The Central Distribution: A Minimax-KL Approach to Noninformative Priors and Feature Selection

Behrouz Ayati* 

Abstract. We introduce the *central distribution*, a probability distribution that represents a set, or more generally a hierarchy, of probability distributions by minimizing the worst-case Kullback–Leibler (KL) divergence to the admissible models. This provides a principled method for converting partial knowledge and set-valued uncertainty into a single probability measure suitable for Bayesian inference.

We show that central distributions admit a minimax characterization and can be obtained as solutions to convex–concave saddle point problems. For parametric models, we establish a sufficient fixed-point condition for priors whose induced marginal distributions are central, establishing a formal connection with Bernardo’s reference prior methodology. We extend the framework to hierarchical uncertainty structures and propose an asymptotic notion of centrality for infinite sequences.

Finally, we illustrate how uncertainty about relevant feature subsets can be represented hierarchically, and how central distributions can be used to aggregate such structural uncertainty into a single predictive model. This suggests an information-theoretic framework for constructing noninformative priors and performing Bayesian feature selection.

MSC2020 subject classifications: Primary 62F15; secondary 62B10, 62H30.

Keywords: Central distribution, noninformative prior, minimax-KL, feature selection, partial knowledge, hierarchical uncertainty.

1 Introduction

We begin by revisiting a fundamental decision-making problem in Bayesian decision theory. An agent must choose from a set of available actions. The loss incurred depends on an unknown state of the environment and is quantified by a real-valued loss function. Specifically, let a denote an action and x the unknown state. The loss associated with choosing action a when the state is x is given by the loss function $\ell(a, x)$.

We assume that some evidence e about the unknown state is available and can be used to guide the agent’s decision. In other words, the agent aims to find an optimal strategy, where a strategy is a conditional distribution over actions given the observed evidence, denoted by $S(A | E)$.

*Independent Researcher, aybehrouz@gmail.com

In Bayesian decision theory, the optimal strategy is the one that minimizes the expected loss:

$$S_{\text{opt}} = \arg \min_S \sum_a \sum_{x,e} \ell(a, x) S(a | e) P(x, e).$$

One can show that there always exists a deterministic optimal strategy, namely a strategy that, for each evidence e , selects the action minimizing the expected loss conditioned on e :

$$S_{\text{opt}}(e) = \arg \min_a \mathbb{E}_P [\ell(a, x) | e].$$

Now suppose that, instead of basing our decisions on the original evidence E , we choose to use a transformed version E' . In practice, the evidence available to the agent is fixed by the environment. However, the agent may choose to base its decisions on a deterministic transformation $E' = f(E)$. This is analogous to feature selection in machine learning, where a high-dimensional observation is replaced by a lower-dimensional representation.

Could such a transformation improve the agent's strategy? Within the framework of Bayesian decision theory, this question has a straightforward answer. Any two strategies can be compared through their expected loss: the strategy with the lower expected loss is preferred. However, a fundamental result of Bayesian decision theory states that the optimal strategy based on E is always at least as good as any strategy based on E' .

One might argue that adopting a strategy that uses simpler evidence may discard information and is therefore not justified by Bayesian decision theory. The key point here is that the space of strategies using E' is a subset of the space of strategies using E . When we search for the optimal strategy based on E , we are simultaneously considering all strategies that depend on E' , since every such strategy can also be expressed as a strategy based on E . This helps explain why standard Bayesian inference does not inherently favor simpler representations of the evidence.

With an appropriate choice of prior distribution, the Bayesian framework may effectively favor strategies that depend only on relevant aspects of the evidence. In this sense, the prior can induce a form of feature selection. However, many commonly used noninformative priors, such as flat priors, do not appear to naturally promote feature selection.

In this paper, we introduce the central distribution, a minimax-KL representative of a set, or more generally a hierarchy, of probability distributions. This provides an information-theoretic framework for converting set-valued and hierarchical uncertainty into a single probability distribution suitable for Bayesian inference. We establish existence, uniqueness, and optimization-theoretic characterizations of central distributions, and develop the corresponding notion of central priors for parametric models. We further identify a sufficient fixed-point condition linking central priors to objective Bayesian reference analysis [2]. Finally, we investigate asymptotic extensions of the framework and illustrate how hierarchical central distributions can be used to represent uncertainty about model structure in a feature-selection setting.

1.1 Related Work

The problem of constructing objective or noninformative priors has a long history in Bayesian statistics. Among the most influential approaches are Jeffreys priors, reference priors, and maximum-entropy based methods [5, 1, 4].

Jeffreys introduced a general rule for constructing invariant noninformative priors based on the Fisher information matrix [5]. The resulting priors are widely used because of their parameterization invariance and their connection to asymptotic likelihood theory. However, Jeffreys priors are derived from local geometric properties of the parameter space and do not directly address the problem of representing a set of admissible probability distributions by a single distribution.

The closest connection to the present work is Bernardo’s reference prior methodology [2, 1]. Reference priors are constructed by maximizing the expected information gained from the data and are intended to represent a state of minimal prior information. In Section 2.3, we show that central priors satisfy a fixed-point condition closely related to the reference prior construction. In particular, for a single parameter and a single set of distributions, the resulting central prior coincides with the corresponding reference prior.

Several authors have also studied objective Bayesian inference through information-theoretic principles [4, 7]. Examples include maximum entropy methods, minimum discrimination information principles, and related divergence-based approaches. These methods typically select a distribution by optimizing an information criterion subject to constraints. In contrast, the central distribution is defined as a minimax representative of a set or hierarchy of distributions, minimizing the worst-case Kullback–Leibler divergence to the admissible models.

The present work is also related to robust Bayesian statistics and imprecise probability [8, 9]. In these frameworks uncertainty is represented by sets of probability distributions rather than a single model. Existing approaches generally focus on performing inference directly with sets of distributions. The central distribution framework instead seeks a single representative distribution that summarizes the available set-valued or hierarchical uncertainty while preserving its information-theoretic structure.

The central distribution framework differs from these approaches in two respects. First, it is defined as a minimax-KL representative of a set, or more generally a hierarchy, of probability distributions. Second, it provides a mechanism for converting set-valued and hierarchical uncertainty into a single probability distribution suitable for Bayesian inference. While related to objective Bayesian inference, robust Bayesian analysis, and information-theoretic methods, its primary goal is not the direct specification of a prior distribution for parameters or inference with sets of distributions, but rather the construction of a representative distribution that preserves the informational structure of the underlying uncertainty.

2 Central Distribution

Consider a random variable X . In Bayesian modeling, our knowledge about X is represented by a probability distribution, which serves as a basis for making inferences.

However, in many situations, our information may be too limited to justify choosing a single distribution. Instead, we may be able to specify a set of plausible distributions. Such a set represents some form of knowledge: although it does not identify a single distribution, it constrains the range of plausible probabilistic models.

A well-known example of this idea is the exchangeability assumption used in de Finetti's theorem. A sequence of random variables $X_1, X_2, \dots$ is said to be exchangeable if the ordering of the variables carries no information. That is, the joint probability distribution remains invariant under permutations of the variables. This assumption imposes a constraint, effectively introducing a set of plausible distributions. Thus, exchangeability can be seen as a specific form of partial knowledge.

In this work, we adopt the view that a set of probability distributions can represent a meaningful state of *uncertainty* or *partial knowledge*, and that such set-valued uncertainty may itself be used as the basis for inference.

More generally, this partial knowledge may possess a hierarchical structure. In the simplest case, uncertainty is represented by a single set of candidate distributions. In richer settings, however, this uncertainty may be organized into nested collections, where elements of a set may themselves be sets of distributions. Such nesting can continue recursively, yielding a hierarchical structure.

Crucially, the choice of hierarchical organization is itself informative: grouping distributions into subsets expresses that certain alternatives are regarded as belonging to a common epistemic category and should therefore be aggregated before being compared with alternatives in other groups. Therefore, the hierarchy encodes information beyond the collection of leaf distributions alone.

This organization may be interpreted as introducing multiple layers of latent structural decisions, where higher-level choices restrict the admissible lower-level alternatives.

Within a Bayesian framework, inference ultimately requires a single probability measure. Hence, to use hierarchical partial knowledge in Bayesian inference, we require a principled method for transforming a nested set-valued uncertainty into a single probability distribution while preserving the informational structure encoded by the hierarchy.

From a minimax perspective, when uncertainty is represented by a set of candidate distributions, a natural representative distribution is one that minimizes the worst-case divergence to the members of the set. Such a distribution may be viewed as the most conservative or robust single-distribution summary of the available knowledge.

Notation 2.1 (Kullback–Leibler Divergence). For probability distributions P and Q , $\mathbb{D}_{\text{KL}}[Q(X) \parallel P(X)]$ denotes the Kullback–Leibler divergence between P and Q over a random variable X . For a discrete random variable X we have

$$\mathbb{D}_{\text{KL}}[Q(X) \parallel P(X)] = \sum_x Q(x) \ln \frac{Q(x)}{P(x)}.$$

When the underlying random variable is clear from context, we omit the argument and write $\mathbb{D}_{\text{KL}}[Q \parallel P]$.

We assume that the KL divergence is defined for every pair of probability distributions P and Q , taking the value $+\infty$ whenever the support of Q is not contained in the support of P .

Definition 2.2. Let X be a random variable and $\mathcal{Q}$ be a set of probability distributions over X . We call a distribution C a *central distribution* of $\mathcal{Q}$ for X if, for every probability distribution P , we have

$$\sup_{Q \in \mathcal{Q}} \mathbb{D}_{\text{KL}}[Q(X) \parallel C(X)] \leq \sup_{Q \in \mathcal{Q}} \mathbb{D}_{\text{KL}}[Q(X) \parallel P(X)], \quad (1)$$

By convention, any probability distribution is regarded as its own central distribution.

Under this definition, the central distribution of $\mathcal{Q}$ is not necessarily an element of $\mathcal{Q}$. That is, the distribution that best represents the set in the specified sense may lie outside the set it summarizes.

Sometimes partial knowledge is represented by a hierarchical structure of sets of distributions. Our current definition of the central distribution does not account for this case. The following recursive definitions extend the concept to handle such situations. While not directly about feature selection, this extension plays a key role in enabling the connection between central distributions and feature selection, a connection that will become clearer later.

Definition 2.3 (Hierarchy of Distributions). A *hierarchy of distributions* over X is defined recursively as follows:

- Any probability distribution over X is a hierarchy of distributions over X .
- Any set whose elements are hierarchies of distributions over X is itself a hierarchy of distributions over X .

Intuitively, a central distribution acts as a representative or summary of a set of distributions. While it may not belong to the set itself, it reflects the overall properties of the set in a way that is meaningful for inference. It also provides a way to evaluate probability distributions under partial knowledge, motivating the definition of a corresponding central divergence.

The concept of a central distribution is not intended to yield a unique object; rather, all distributions satisfying the definition can be regarded as equally good representatives. However, when multiple central distributions exist, our constructions should not depend on an arbitrary choice among them. To avoid such dependence, we define the central divergence using the best divergence attained over the set of all central distributions.

Definition 2.4 (Central Divergence). For any distribution P , we define the *central divergence* of P from a hierarchy of distributions $\mathcal{H}$, with respect to a random variable

X , as follows:

$$\mathbb{D}_C[\mathcal{H}(X) \parallel P(X)] = \inf_C \mathbb{D}_{\text{KL}}[C(X) \parallel P(X)],$$

where C denotes a central distribution of $\mathcal{H}$ for X .

So far, this definition applies only when the hierarchy is a single set of distributions. The following definition extends the construction recursively to general hierarchical uncertainty structures.

Definition 2.5 (Central Distribution). Let X be a random variable, and let $\mathcal{H}^*$ be a hierarchy of distributions over X . We call a distribution C a *central distribution* of $\mathcal{H}^*$ if, for every probability distribution P , we have:

$$\sup_{\mathcal{H} \in \mathcal{H}^*} \mathbb{D}_C[\mathcal{H}(X) \parallel C(X)] \leq \sup_{\mathcal{H} \in \mathcal{H}^*} \mathbb{D}_C[\mathcal{H}(X) \parallel P(X)].$$

Definition 2.5 generalizes Definition 2.2, which is recovered when the hierarchy is a set of probability distributions.

For later use, we introduce a compact notation for the minimax objective appearing in Definition 2.2.

Notation 2.6. For a set of probability distributions $\mathcal{Q}$ over X , we define the functional

$$\mathbb{S}_{\mathcal{Q}}[P(X)] := \sup_{Q \in \mathcal{Q}} \mathbb{D}_{\text{KL}}[Q(X) \parallel P(X)].$$

When $\mathcal{Q}$ and X are clear from the context, we may simply write $\mathbb{S}[P]$.

This notation allows Definition 2.2 to be written compactly as

$$C \in \arg \min_P \mathbb{S}_{\mathcal{Q}}[P(X)].$$

A simple example illustrates that hierarchical organization can affect the resulting representative distribution even when the underlying admissible sets remain unchanged.

Example. Let X be a random variable with three possible outcomes x_1, x_2, x_3 . For each i , let S_i denote the set of all probability distributions over X that assign zero probability to x_i .

By symmetry, the central distribution of each set S_i is the uniform distribution over its support:

$$C_{S_1} = (0, 0.5, 0.5), \quad C_{S_2} = (0.5, 0, 0.5), \quad C_{S_3} = (0.5, 0.5, 0).$$

If the three sets S_1, S_2, S_3 are treated symmetrically as elements of a single set, then their central distribution is

$$C_{\{S_1, S_2, S_3\}} = \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3} \right).$$

Now consider instead a hierarchical organization in which S_1 and S_2 are grouped together, while S_3 forms its own separate branch.

By numerically solving the minimax problem in Definition 2.5, we obtain

$$C_{\{S_1, S_2\}} = (0.25, 0.25, 0.5).$$

Applying the same procedure to the two distributions $C_{\{S_1, S_2\}}$ and C_{S_3} yields

$$C_{\{\{S_1, S_2\}, S_3\}} = (0.4, 0.4, 0.2).$$

Although the underlying admissible sets are identical in both constructions, the resulting representative distribution differs.

This demonstrates that hierarchical structure carries information beyond the collection of leaf distributions alone. Consequently, flattening a hierarchy into a single unstructured set may discard meaningful epistemic structure.

In many learning problems, uncertainty concerns not only the parameters of a predictive model but also its structural form. In particular, when the relevance of potential features is unknown, each candidate feature subset determines a distinct set of plausible models in which the target variable depends only on that feature subset.

Feature selection may therefore be viewed as uncertainty over a collection of distributions. The hierarchical central distribution framework provides a principled mechanism for representing and aggregating such uncertainty. By organizing candidate models hierarchically, one can encode preferences over structural hypotheses, such as favoring sparse subsets, grouping related attributes, or privileging particular interactions.

Candidate feature subsets typically induce overlapping families of admissible distributions. Distributions representing simpler dependency structure may belong simultaneously to many such families, whereas more complex dependency patterns belong to fewer. As a result, hierarchical aggregation through central distributions can induce a preference toward structurally simpler models by naturally assigning more weight to distributions compatible with multiple structural hypotheses.

2.1 Minimax Characterization

With the concept of hierarchical central distributions in place, we now develop the theoretical machinery needed to characterize and compute them. Throughout this subsection, X denotes a discrete random variable with finite support and $\mathcal{Q}$ denotes a finite set of probability distributions over X .

As we shall see, central distributions arise naturally as solutions to a convex-concave optimization problem. By formulating the appropriate objective function, we obtain a saddle-point characterization that enables the use of standard techniques from convex optimization.

Definition 2.7 (Saddle Point). Consider a function $f : \mathcal{W} \times \mathcal{Z} \rightarrow \mathbb{R}$. We refer to a pair (w^*, z^*) as a *saddle point* for f if we have

$$f(w^*, z) \leq f(w^*, z^*) \leq f(w, z^*),$$

for all $w \in \mathcal{W}$ and $z \in \mathcal{Z}$.

Lemma 2.8. *Define:*

$$f(P, R) = \sum_i \mathbb{D}_{\text{KL}}[Q_i(X) \parallel P(X)] R(i),$$

where P and R are probability distributions, $Q_i \in \mathcal{Q}$ and the index i ranges over all elements of $\mathcal{Q}$. Then there exists a saddle point for f .

Proof. The Kullback–Leibler divergence is convex and lower-semicontinuous in its second argument. Therefore, for any fixed probability distribution R , the function $f(\cdot, R)$ is a finite convex combination of lower-semicontinuous convex functions and is therefore itself convex and lower-semicontinuous in P .

For fixed P , the function $f(P, \cdot)$ is affine in R , and hence concave and continuous.

The set of probability distributions over a discrete random variable with finite support is a compact convex simplex. Therefore, the assumptions of Sion’s minimax theorem are satisfied [6], and

$$\min_P \sup_R f(P, R) = \sup_R \min_P f(P, R).$$

Since $f(P, \cdot)$ is continuous and the feasible set for R is compact, the supremum is attained for every P . Hence the minimax solution exists and f admits a saddle point. $\square$

Lemma 2.9. *Let f be the function defined in Lemma 2.8. If (P^*, R^*) is a saddle point for f , then P^* is a central distribution of $\mathcal{Q}$ for X .*

Proof. Let P be an arbitrary probability distribution over X , and (P^*, R^*) be a saddle point for f . As P is a probability distribution, $f(P, R^*)$ is defined, and by the definition of a saddle point

$$f(P^*, R^*) \leq f(P, R^*). \quad (2)$$

Since R^* is a probability distribution, the definition of f implies:

$$f(P, R^*) \leq \mathbb{S}[P].$$

On the other hand, by the definition of a saddle point:

$$f(P^*, R^*) = \max_R f(P^*, R) = \mathbb{S}[P^*].$$

Hence, Equation 2 implies

$$\mathbb{S}[P^*] \leq \mathbb{S}[P].$$

This holds for any arbitrary P . Therefore, by Definition 2.2, P^* is a central distribution of $\mathcal{Q}$ for X . $\square$

Lemma 2.10. *Let f be the function defined in Lemma 2.8. The point (P^*, R^*) is a saddle point for f , if and only if there is a constant λ , and a function μ , such that for every x and i the following conditions hold:*

$$\sum_x Q_i(x) \ln \frac{Q_i(x)}{P^*(x)} + \mu(i) = \lambda, \quad (3)$$

$$P^*(x) = \sum_i R^*(i) Q_i(x) \quad (4)$$

$$\sum_i R^*(i) = 1, \quad (5)$$

$$R^*(i) \geq 0, \quad (6)$$

$$\mu(i) \geq 0, \quad (7)$$

$$\mu(i) R^*(i) = 0. \quad (8)$$

Furthermore, for a fixed probability distribution R , a distribution $\hat{P}$ minimizes $f(\cdot, R)$, if and only if for all x it satisfies

$$\hat{P}(x) = \sum_i Q_i(x) R(i). \quad (9)$$

Proof. We extend f to a function $\hat{f}$ defined as follows:

$$\hat{f}(P, R) = \sum_i R(i) \sum_x Q_i(x) \ln \frac{Q_i(x)}{P(x)},$$

where the domain is extended by relaxing the constraints that P and R be probability distributions. This enables us to formulate the saddle points of f as solutions to a constrained optimization problem defined for $\hat{f}$.

Let (P^*, R^*) be a simultaneous solution to the following two optimization problems:

$$\begin{aligned} & \text{minimize} && \hat{f}(P, R) && \text{w.r.t. } P \\ & \text{subject to} && \sum_x P(x) = 1, \\ & && && \\ & \text{maximize} && \hat{f}(P, R) && \text{w.r.t. } R \\ & \text{subject to} && R(i) \geq 0, && \text{for all } i \\ & && \sum_i R(i) = 1 \end{aligned} \quad (10)$$

Clearly, if (P^*, R^*) exists, it will be a saddle point for f .

Both optimization problems are convex and differentiable; see [3]. To establish this, consider that all the constraint functions are affine. Additionally, both $\hat{f}(P, \cdot)$ and

$\hat{f}(\cdot, R)$ are differentiable on their domains. For any fixed P , $\hat{f}(P, \cdot)$ is an affine function and is concave. For a fixed solution to the second optimization problem denoted by $\hat{R}$, we can write

$$\hat{f}(P, \hat{R}) = c - \sum_{i,x} \hat{R}(i) Q_i(x) \ln P(x), \quad (11)$$

where c is not a function of P . Since $\hat{R}$ is a solution to the second optimization problem, it satisfies the corresponding constraints and is therefore a valid probability distribution. Hence, for all i, x we have $\hat{R}(i) Q_i(x) \geq 0$, and $\hat{f}(\cdot, \hat{R})$ is the negative of a linear combination of concave $\ln P(x)$ functions with nonnegative coefficients. This function is therefore convex.

Thus, both optimization problems are convex and differentiable, with affine equality and inequality constraints. In this setting, Slater's condition reduces to feasibility [3]. The feasible set for the first problem is the probability simplex, while the feasible set for the second problem is the strictly positive probability simplex. Both sets are nonempty. Therefore, Slater's condition holds, and the KKT conditions are necessary and sufficient for optimality. In other words, if there exists a point (P^*, R^*) that satisfies the KKT conditions for both problems, it will be a saddle point for f , and vice versa.

The KKT conditions are as follows. There exist constants λ, η and a function μ , such that for every x and i :

$$\sum_i R^*(i) Q_i(x) = \eta P^*(x) \quad (12)$$

$$\sum_x P^*(x) = 1, \quad (13)$$

$$\begin{aligned} \sum_x Q_i(x) \ln \frac{Q_i(x)}{P^*(x)} + \mu(i) &= \lambda, \\ \sum_i R^*(i) &= 1, \\ R^*(i) &\geq 0, \\ \mu(i) &\geq 0, \\ \mu(i) R^*(i) &= 0. \end{aligned} \quad (14)$$

By summing Equation 12 over x and using Equation 14, we obtain $\eta = 1$. Substituting this into Equation 12 yields Equation 4, which in turn implies Equation 13.

To prove the second part of the lemma, we note that Equations 12 and 13 are the KKT conditions for the minimization problem in Equation 10. As shown above, the KKT conditions are necessary and sufficient for optimality, and when R is a probability distribution, these conditions simplify to

$$\hat{P}(x) = \sum_i Q_i(x) R(i). \quad \square$$

Proof. Since $Q_\theta \in \mathcal{Q}$ is a probability distribution, for every θ there exists x such that $Q_\theta(x) > 0$. Equation 19 therefore implies $c(\theta | x) > 0$, and consequently $c(\theta) > 0$.

Using Equation 19 and $c(\theta) > 0$, we obtain

$$\begin{aligned} c(\theta) &= \exp \left\{ -\lambda + \sum_x Q(x | \theta) \ln \frac{Q(x | \theta) c(\theta)}{C(x)} \right\} \\ &= \exp \{ -\lambda + \mathbb{D}_{\text{KL}} [Q_\theta(X) \| C(X)] + \ln c(\theta) \} \\ &= c(\theta) \exp \{ -\lambda + \mathbb{D}_{\text{KL}} [Q_\theta(X) \| C(X)] \}, \end{aligned}$$

and it follows that

$$\mathbb{D}_{\text{KL}} [Q_\theta(X) \| C(X)] = \lambda.$$

Since this holds for every θ , and λ is constant, a parametric extension of Theorem 2.12 implies that C , the marginal distribution induced by c , is a central distribution of $\mathcal{Q}$ for X . Therefore, by Definition 2.17, c is a central prior of $\mathcal{Q}$ for X . $\square$

This theorem is proved using Theorem 2.12. However, Theorem 2.14 suggests an alternative route to establishing the connection with reference priors. Since reference priors are defined by maximizing missing information, the optimization characterization provided by Theorem 2.14 also offers another perspective on this relationship.

2.4 Asymptotic Central Distributions

Many probabilistic models are naturally formulated in terms of a sequence of random variables whose length is not fixed in advance. For example, when a coin is tossed repeatedly, there is no natural upper bound on the number of tosses, and any finite sample can be viewed as an initial segment of an infinite sequence.

Motivated by such examples, let S_n denote the first n variables of an infinite sequence of random variables:

$$S_n = (X_1, X_2, \dots, X_n).$$

The collection $\{S_n\}_{n=1}^\infty$ therefore defines a second sequence of random variables.

In many settings, our uncertainty about S_n can be represented by a central distribution. A prominent example arises when the sequence is assumed to be exchangeable. As discussed earlier, exchangeability can be viewed as a form of partial knowledge that defines a set of plausible distributions, from which a central distribution for S_n can be constructed.

However, this central distribution generally depends on the sequence length n . In this work, we adopt the principle that the probability measure used to represent prior beliefs about the environment should depend only on the available information about the environment itself, and not on arbitrary aspects of the experimental setup, such as the number of variables under consideration.

Importantly, a distribution defined over a longer sequence induces marginal distributions over all shorter initial segments. This suggests that, rather than constructing a

By Theorem 2.21 and Equation 21, we can derive a central distribution for $\{S_n\}_{n=1}^\infty$, which induces the following distribution over S_n :

$$C(S_n) = \frac{\Gamma\left(\frac{m}{2}\right)}{\Gamma\left(n + \frac{m}{2}\right)} \prod_{y=1}^m \frac{\Gamma\left(n_y + \frac{1}{2}\right)}{\Gamma\left(\frac{1}{2}\right) |f^{-1}(y)|^{n_y}}.$$

When dealing with layered ignorance and seeking the central distribution of a hierarchy of distributions, we typically encounter situations in which a given level of the hierarchy contains only finitely many alternatives. In such cases, and under appropriate regularity conditions, if there exists a consistent estimator for the index identifying these alternatives, then the central distribution reduces to an equally weighted average of the corresponding representative central distributions.

3 An Application to Feature Selection

A natural application of hierarchical central distributions arises in feature selection. When the relevant set of features is unknown, each candidate feature set induces a distinct family of admissible distributions. Feature selection can therefore be viewed as uncertainty over a hierarchy of probabilistic models, making it a natural setting for the central distribution framework.

To illustrate this, we provide a proof of concept experiment using synthetic data. The purpose of this section is not to establish a competitive feature selection procedure, but to illustrate how hierarchical central distributions can be used to represent uncertainty over model structure. A comprehensive empirical evaluation against established feature selection methods, such as Spike-and-Slab priors, on real-world datasets lies beyond the scope of this foundational paper and is left for future work.

We consider a classification problem where the goal is to predict a target class, denoted by Y , from a set of observed nominal attributes $(A_1, A_2, \dots, A_n)$. We assume that Y may depend on an unknown subset of the attributes, rather than the full set.

To capture this relationship, we use the fused multinomial model introduced in Section 2.4. As an example, assume that Y depends only on A_2 and A_5 . Then, we define the labeling function f as follows:

$$f(A_1, A_2, \dots, A_n, Y) = (A_2, A_5, Y).$$

In a fused multinomial model with this labeling function, the target class Y is conditionally independent of all other attributes given A_2, A_5 .

For every possible subset of attributes, a distinct fused multinomial model of this form is constructed. Each such model corresponds to a set of plausible probability distributions. Collecting these sets yields our overall uncertainty representation, which forms a one-level hierarchy of distributions (Definition 2.3).

Let $C_{\mathcal{H}^*}$ denote a central distribution of the full hierarchy, and let C_{f_i} denote the central distribution of the fused multinomial model associated with the labeling function

Table 1: Mappings of 3-bit inputs under class functions Y_1 and Y_2

Input (bits)	$Y_1(\text{Input})$	$Y_2(\text{Input})$
000	a	A
001	b	B
010	c	A
011	d	C
100	e	D
101	f	D
110	g	D
111	h	E

f_i . As discussed earlier, because the number of candidate models is finite, we have

$$C_{\mathcal{H}^*}(S_n) \propto \sum_i C_{f_i}(S_n). \quad (22)$$

To justify this, observe that we can construct a consistent estimator for the model indicator I by examining equality relationships among the observed frequencies of outcomes. Specifically, the estimator identifies outcomes that occur with equal frequencies and returns the indicator corresponding to the model that supports those equalities.

The optimal Bayesian decision rule depends on the choice of loss function. In this example, we use the zero-one loss, under which the optimal strategy predicts the target class with the highest posterior probability. These posterior probabilities are computed using the central distribution of Equation 22. It turns out that each subset-specific fused multinomial model is assigned its corresponding Jeffreys' prior. Consequently, the resulting predictive distribution can be interpreted as Bayesian model averaging over all candidate feature subsets, with each model equipped with its Jeffreys' prior. The central distribution framework provides a minimax-KL justification for this aggregation.

To see how this approach works in practice, we carried out a synthetic experiment with the following setup. We considered binary attributes and generated each attribute independently with equal probability for 0 and 1. The target class was defined as a deterministic function of a fixed subset of 3 out of 12 total attributes, as shown in Table 1. The same subset of relevant attributes and class assignment function were used across all trials.

For each experiment, we generated a test set of size 100. A larger test set was used to reduce variability in performance estimates, taking advantage of the synthetic nature of the data. To assess robustness to irrelevant features, we progressively removed one attribute at a time from the full set of 12, starting with irrelevant ones and stopping just before removing any of the three relevant attributes.

At each step, we trained and tested two versions of the central distribution: one assuming that the target class is a deterministic function of the attributes (CD-Det), and one allowing the target class to follow a conditional probability distribution given

- URL <https://www.sciencedirect.com/science/article/pii/S0169716105250022> 2, 3, 16, 19
- [3] Boyd, S. and Vandenberghe, L. (2004). *Convex Optimization*. Cambridge University Press.
URL <https://books.google.com/books?id=IUZdAAAAQBAJ> 9, 10
- [4] Jaynes, E. T. (1957). “Information Theory and Statistical Mechanics.” *Phys. Rev.*, 106: 620–630.
URL <https://link.aps.org/doi/10.1103/PhysRev.106.620> 3
- [5] Jeffreys, H. (1946). “An invariant form for the prior probability in estimation problems.” *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 186(1007): 453–461.
URL <https://doi.org/10.1098/rspa.1946.0056> 3
- [6] Komiya, H. (1988). “Elementary proof for Sion’s minimax theorem.” *Kodai Mathematical Journal*, 11(1): 5 – 7.
URL <https://doi.org/10.2996/kmj/1138038812> 8
- [7] Kullback, S. and Leibler, R. A. (1951). “On Information and Sufficiency.” *The Annals of Mathematical Statistics*, 22(1): 79 – 86.
URL <https://doi.org/10.1214/aoms/1177729694> 3
- [8] Rios Insua, D. and Ruggeri, F. (2000). *Robust Bayesian Analysis*, volume 152 of *Lecture Notes in Statistics*. New York: Springer-Verlag. 3
- [9] Walley, P. (1991). *Statistical Reasoning with Imprecise Probabilities*. Chapman & Hall/CRC Monographs on Statistics & Applied Probability. Taylor & Francis.
URL <https://books.google.com/books?id=-hbvAAAAAAAJ> 3