

MATHS 130 – Assignment 3

University of Auckland

Aidan Webster

aweb904@aucklanduni.ac.nz

June 1, 2023

Contents

Question 1 1

Question 2 2

Question 3 3

Question 4 4

Question 1

Consider the equation $f(x) = 0$, on \mathbb{R} , where

$$f(x) := 7x^3 + 2x - \sin(x) - 1.$$

- (a) Prove that there is at least one solution to the equation.
- (b) Use the mean value theorem to prove that there is only one solution.

Question 2

Determine whether the function

$$h(x) = \frac{1}{x^2 - 4x}$$

has any absolute extrema on the interval $(0, 4)$. If there are, find them and state where they occur, otherwise prove that there are none.

Question 3

Consider the function of two real variables given by the formula

$$f(x, y) = \frac{1}{\sqrt{x^2 - 9y^2}}$$

- (a) Determine the natural domain on f .
- (b) Determine and sketch the level curves of f for the values $k = 0$, and $k = 1$.
- (c) Compute the gradient ∇f at a general point (x, y) and then at the point $(4, 1)$.
- (d) Compute the directional derivative of f at $(4, 1)$ in the direction of the vector $u = (9, 4)$.
- (e) At the point $(4, 1)$ find the unit vector of a direction in which the function increases most rapidly. What is the directional derivative in that direction?

Question 4

Consider the integral

$$g(x) := \int_{-7}^x |t|^{\frac{1}{2}} \cdot e^{t^2} dt$$

- (a) Without calculating it, explain why $g'(x)$ exists for all $x \in \mathbb{R}$.
- (b) Compute $g'(x)$, naming any results/theorems that you use.
- (c) Use the mean value theorem to prove that g is strictly increasing on \mathbb{R} .
- (d) Use **proof by contradiction** to prove that the second derivative of h does not exist at 0.