






OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	2
1.4	Final three nucleotides: frequency of each observed triplet	2
1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele	4
3	Allele usage in potential haplotype anchor genes	9
4	Haplotype plots	10
5	Configuration settings	11



























1.5 CDR3 length distribution, in assignments to novel alleles























2 Variation from germline, in assignments to each allele












3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /work/jenkins/PRJEB26509/S2/S2/2_IGH/2_IGH/e1/f83f8e7fccd36160a86ab49ad48e3e/2_Fina
##
## Germline reference file: /work/jenkins/PRJEB26509/S2/S2/2_IGH/2_IGH/e1/f83f8e7fccd36160a86ab49ad48e3
##
## Novel allele file: /work/jenkins/PRJEB26509/S2/S2/2_IGH/2_IGH/e1/f83f8e7fccd36160a86ab49ad48e3e/nove
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```