


OGRDBstats Report


Contents













1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	2
1.4	Final three nucleotides: frequency of each observed triplet	2
1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele	4
3	Allele usage in potential haplotype anchor genes	9
4	Haplotype plots	10
5	Configuration settings	11













1.5 CDR3 length distribution, in assignments to novel alleles






















2 Variation from germline, in assignments to each allele












3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJEB26509/S91/S91/91_IGH/91_IGH/65/438eff1f32a73d27f8458b618334
##
## Germline reference file: /misc/work/jenkins/PRJEB26509/S91/S91/91_IGH/91_IGH/65/438eff1f32a73d27f845
##
## Novel allele file: /misc/work/jenkins/PRJEB26509/S91/S91/91_IGH/91_IGH/65/438eff1f32a73d27f8458b6183
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```