


OGRDBstats Report






Contents

1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	4
1.3	Whole-sequence composition of each assigned read	6
1.4	Final three nucleotides: frequency of each observed triplet	7
1.5	CDR3 length distribution, in assignments to novel alleles	8
2	Variation from germline, in assignments to each allele	10
3	Allele usage in potential haplotype anchor genes	18
4	Haplotype plots	19
5	Configuration settings	20


1 Novel sequence analysis






1.1 End-nucleotide composition







1.2 Per-nucleotide consensus where previous nucleotides match the consensus









1.3 Whole-sequence composition of each assigned read


















1.4 Final three nucleotides: frequency of each observed triplet


















1.5 CDR3 length distribution, in assignments to novel alleles









































2 Variation from germline, in assignments to each allele












































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots



5 Configuration settings

```
## Repertoire file: /work/jenkins/PRJNA381394/B13_N/B13_N/4_IGH/4_IGH/79/fe6844cf000ddd79a6f99a1c2898fe.  
##  
## Germline reference file: /work/jenkins/PRJNA381394/B13_N/B13_N/4_IGH/4_IGH/79/fe6844cf000ddd79a6f99a.  
##  
## Novel allele file: /work/jenkins/PRJNA381394/B13_N/B13_N/4_IGH/4_IGH/79/fe6844cf000ddd79a6f99a1c2898.  
##  
## Species: Homosapiens  
##  
## Chain: IGHV  
##  
## Segment: V
```