


OGRDBstats Report






Contents

1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	4
1.3	Whole-sequence composition of each assigned read	6
1.4	Final three nucleotides: frequency of each observed triplet	7
1.5	CDR3 length distribution, in assignments to novel alleles	8
2	Variation from germline, in assignments to each allele	10
3	Allele usage in potential haplotype anchor genes	17
4	Haplotype plots	18
5	Configuration settings	19


1 Novel sequence analysis






1.1 End-nucleotide composition







1.2 Per-nucleotide consensus where previous nucleotides match the consensus







1.3 Whole-sequence composition of each assigned read










1.4 Final three nucleotides: frequency of each observed triplet


















1.5 CDR3 length distribution, in assignments to novel alleles









































2 Variation from germline, in assignments to each allele





































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /work/jenkins/PRJNA381394/B14_N/B14_N/5_IGH/5_IGH/bf/45df78b478df2e732d98991e7ad472,
##
## Germline reference file: /work/jenkins/PRJNA381394/B14_N/B14_N/5_IGH/5_IGH/bf/45df78b478df2e732d9899
##
## Novel allele file: /work/jenkins/PRJNA381394/B14_N/B14_N/5_IGH/5_IGH/bf/45df78b478df2e732d98991e7ad4
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```