


OGRDBstats Report


Contents

1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	3
1.3	Whole-sequence composition of each assigned read	5
1.4	Final three nucleotides: frequency of each observed triplet	6
1.5	CDR3 length distribution, in assignments to novel alleles	7
2	Variation from germline, in assignments to each allele	8
3	Allele usage in potential haplotype anchor genes	15
4	Haplotype plots	16
5	Configuration settings	17


1 Novel sequence analysis








1.1 End-nucleotide composition







1.2 Per-nucleotide consensus where previous nucleotides match the consensus






1.3 Whole-sequence composition of each assigned read







1.4 Final three nucleotides: frequency of each observed triplet



























1.5 CDR3 length distribution, in assignments to novel alleles



























2 Variation from germline, in assignments to each allele






































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots



5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA381394/B16_N/B16_N/6_IGH/6_IGH/67/55c527889a780821fb140cbd0
##
## Germline reference file: /misc/work/jenkins/PRJNA381394/B16_N/B16_N/6_IGH/6_IGH/67/55c527889a780821f
##
## Novel allele file: /misc/work/jenkins/PRJNA381394/B16_N/B16_N/6_IGH/6_IGH/67/55c527889a780821fb140cb
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```