


OGRDBstats Report









Contents

1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	4
1.3	Whole-sequence composition of each assigned read	6
1.4	Final three nucleotides: frequency of each observed triplet	8
1.5	CDR3 length distribution, in assignments to novel alleles	9
2	Variation from germline, in assignments to each allele	11
3	Allele usage in potential haplotype anchor genes	18
4	Haplotype plots	19
5	Configuration settings	20


1 Novel sequence analysis









1.1 End-nucleotide composition










1.2 Per-nucleotide consensus where previous nucleotides match the consensus






1.3 Whole-sequence composition of each assigned read










1.4 Final three nucleotides: frequency of each observed triplet
















1.5 CDR3 length distribution, in assignments to novel alleles









































2 Variation from germline, in assignments to each allele





































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots



5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA381394/B18_N/B18_N/8_IGH/8_IGH/e7/189f5512bd6d78399165da49b
##
## Germline reference file: /misc/work/jenkins/PRJNA381394/B18_N/B18_N/8_IGH/8_IGH/e7/189f5512bd6d78399
##
## Novel allele file: /misc/work/jenkins/PRJNA381394/B18_N/B18_N/8_IGH/8_IGH/e7/189f5512bd6d78399165da4
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```