


OGRDBstats Report


Contents













1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	2
1.4	Final three nucleotides: frequency of each observed triplet	2
1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele	4
3	Allele usage in potential haplotype anchor genes	9
4	Haplotype plots	10
5	Configuration settings	11













1.5 CDR3 length distribution, in assignments to novel alleles

























2 Variation from germline, in assignments to each allele












3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /work/jenkins/PRJNA280743/TET3_45/TET3_45/772/772/6e/e612274aae76c985b260fdad149fa2
##
## Germline reference file: /work/jenkins/PRJNA280743/TET3_45/TET3_45/772/772/6e/e612274aae76c985b260fd
##
## Novel allele file: /work/jenkins/PRJNA280743/TET3_45/TET3_45/772/772/6e/e612274aae76c985b260fdad149f
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
```