


# OGRDBstats Report


## Contents

|          |                                                                                   |           |
|----------|-----------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Novel sequence analysis</b>                                                    | <b>2</b>  |
| 1.1      | End-nucleotide composition . . . . .                                              | 2         |
| 1.2      | Per-nucleotide consensus where previous nucleotides match the consensus . . . . . | 3         |
| 1.3      | Whole-sequence composition of each assigned read . . . . .                        | 4         |
| 1.4      | Final three nucleotides: frequency of each observed triplet . . . . .             | 5         |
| 1.5      | CDR3 length distribution, in assignments to novel alleles . . . . .               | 6         |
| <b>2</b> | <b>Variation from germline, in assignments to each allele</b>                     | <b>8</b>  |
| <b>3</b> | <b>Allele usage in potential haplotype anchor genes</b>                           | <b>15</b> |
| <b>4</b> | <b>Haplotype plots</b>                                                            | <b>16</b> |
| <b>5</b> | <b>Configuration settings</b>                                                     | <b>17</b> |


# 1 Novel sequence analysis


## 1.1 End-nucleotide composition







## 1.2 Per-nucleotide consensus where previous nucleotides match the consensus







### 1.3 Whole-sequence composition of each assigned read















## 1.4 Final three nucleotides: frequency of each observed triplet






## 1.5 CDR3 length distribution, in assignments to novel alleles









































## 2 Variation from germline, in assignments to each allele

















































### 3 Allele usage in potential haplotype anchor genes



## 4 Haplotype plots



## 5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA280743/SLE1_126/SLE1_126/770/770/9d/5c07d14dd05681a3eaf4d5f
##
## Germline reference file: /misc/work/jenkins/PRJNA280743/SLE1_126/SLE1_126/770/770/9d/5c07d14dd05681a
##
## Novel allele file: /misc/work/jenkins/PRJNA280743/SLE1_126/SLE1_126/770/770/9d/5c07d14dd05681a3eaf4d
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
##
## Novel sequence(s) IGHV3_23_01_C113G IGHV3_23_04_C113G are not listed in the genotype and will be ign
##
##
## No sequences found for IGHV3_23_01_C113G(check SEQUENCE_IMGT in input file)
##
## No sequences found for IGHV3_23_04_C113G(check SEQUENCE_IMGT in input file)
```