






OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	3
1.3	Whole-sequence composition of each assigned read	4
1.4	Final three nucleotides: frequency of each observed triplet	4
1.5	CDR3 length distribution, in assignments to novel alleles	5
2	Variation from germline, in assignments to each allele	6
3	Allele usage in potential haplotype anchor genes	13
4	Haplotype plots	14
5	Configuration settings	15

1 Novel sequence analysis


1.1 End-nucleotide composition




1.2 Per-nucleotide consensus where previous nucleotides match the consensus




1.3 Whole-sequence composition of each assigned read




1.4 Final three nucleotides: frequency of each observed triplet



























1.5 CDR3 length distribution, in assignments to novel alleles



























2 Variation from germline, in assignments to each allele

































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA324093/Donor 5/122 S049/5_IGH/5_IGH/b1/3d9482ed20c896007856
##
## Germline reference file: /misc/work/jenkins/PRJNA324093/Donor 5/122 S049/5_IGH/5_IGH/b1/3d9482ed20c8
##
## Novel allele file: /misc/work/jenkins/PRJNA324093/Donor 5/122 S049/5_IGH/5_IGH/b1/3d9482ed20c8960078
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
##
## Unexpected N in reference sequence IGHV1 24 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV1 24 01_A315G_C317G: replacing with gap
##
##
## Unexpected N in reference sequence IGHV3 30 03: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3 30 03_T288C: replacing with gap
##
##
## Unexpected N in reference sequence IGHV4 34 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV4 34 01_T300C: replacing with gap
##
##
## Unexpected N in reference sequence IGHV4 59 12: replacing with gap
##
##
## Unexpected N in novel sequence IGHV4 59 12_C300T: replacing with gap
##
##
## Unexpected N in reference sequence IGHV4 NL1 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV4 NL1 01_C300T: replacing with gap
```