


OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	3
1.4	Final three nucleotides: frequency of each observed triplet	3
1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele	4
3	Allele usage in potential haplotype anchor genes	10
4	Haplotype plots	11
5	Configuration settings	12

1 Novel sequence analysis


1.1 End-nucleotide composition




1.2 Per-nucleotide consensus where previous nucleotides match the consensus




1.3 Whole-sequence composition of each assigned read




1.4 Final three nucleotides: frequency of each observed triplet



























1.5 CDR3 length distribution, in assignments to novel alleles



























2 Variation from germline, in assignments to each allele


























3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA324093/Donor 162/036 S027/165_IGH/165_IGH/5d/a4d7eb5f40b4aa
##
## Germline reference file: /misc/work/jenkins/PRJNA324093/Donor 162/036 S027/165_IGH/165_IGH/5d/a4d7eb
##
## Novel allele file: /misc/work/jenkins/PRJNA324093/Donor 162/036 S027/165_IGH/165_IGH/5d/a4d7eb5f40b4
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
##
## Unexpected N in reference sequence IGHV1 18 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV1 18 04_G107C_A110T_G112A_G119C_G126C_C134T_A144G_G147A_G153A_A16
##
##
## Unexpected N in reference sequence IGHV4 34 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV4 34 01_T300C: replacing with gap
##
##
## Unexpected N in reference sequence IGHV4 59 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV4 59 12_C300T: replacing with gap
```