


OGRDBstats Report


Contents

1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	3
1.3	Whole-sequence composition of each assigned read	4
1.4	Final three nucleotides: frequency of each observed triplet	5
1.5	CDR3 length distribution, in assignments to novel alleles	6
2	Variation from germline, in assignments to each allele	8
3	Allele usage in potential haplotype anchor genes	15
4	Haplotype plots	16
5	Configuration settings	17


1 Novel sequence analysis


1.1 End-nucleotide composition







1.2 Per-nucleotide consensus where previous nucleotides match the consensus







1.3 Whole-sequence composition of each assigned read















1.4 Final three nucleotides: frequency of each observed triplet


















1.5 CDR3 length distribution, in assignments to novel alleles






























2 Variation from germline, in assignments to each allele


































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots



5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA491287/M64 013/M64 013/M64 013/M64 013/2c/8d48bcccd08bbacc6
##
## Germline reference file: /misc/work/jenkins/PRJNA491287/M64 013/M64 013/M64 013/M64 013/2c/8d48bcccd
##
## Novel allele file: /misc/work/jenkins/PRJNA491287/M64 013/M64 013/M64 013/M64 013/2c/8d48bcccd08bbacc
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
##
## Unexpected N in reference sequence IGHV1 8 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV1 8 01_G107C: replacing with gap
##
##
## Unexpected N in reference sequence IGHV1 69 2 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV1 24 01_A315G_C317G: replacing with gap
##
##
## Unexpected N in reference sequence IGHV3 30 03: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3 30 03_C75G_T288C: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3 30 03_T288C: replacing with gap
##
##
## Unexpected N in reference sequence IGHV3 30 3 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3 30 3 01_T288C: replacing with gap
##
##
## Unexpected N in reference sequence IGHV3 30 18: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3 33 06_G170C_G171A: replacing with gap
##
##
## Unexpected N in reference sequence IGHV3 74 01: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3 74 01_A315G: replacing with gap
##
##
## Unexpected N in reference sequence IGHV4 59 01: replacing with gap
```

```
##
##
## Unexpected N in novel sequence IGHV4 59 01_A138G: replacing with gap
##
##
## Unexpected N in reference sequence IGHV4 61 02: replacing with gap
##
##
## Unexpected N in novel sequence IGHV4 61 02_A291G_C300T: replacing with gap
```