


OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	2
1.4	Final three nucleotides: frequency of each observed triplet	2
1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele	4
3	Allele usage in potential haplotype anchor genes	11
4	Haplotype plots	12
5	Configuration settings	13

1 Novel sequence analysis


1.1 End-nucleotide composition




1.2 Per-nucleotide consensus where previous nucleotides match the consensus




1.3 Whole-sequence composition of each assigned read




1.4 Final three nucleotides: frequency of each observed triplet



























1.5 CDR3 length distribution, in assignments to novel alleles



























2 Variation from germline, in assignments to each allele






































3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJNA491287/M64_029/M64_029/M64_029/db/b4fad6eb41f73cdf1.
##
## Germline reference file: /misc/work/jenkins/PRJNA491287/M64_029/M64_029/M64_029/db/b4fad6eb4.
##
## Novel allele file: /misc/work/jenkins/PRJNA491287/M64_029/M64_029/M64_029/db/b4fad6eb41f73cd.
##
## Species: Homosapiens
##
## Chain: IGHV
##
## Segment: V
##
## Unexpected N in reference sequence IGHV3_23_01_04: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3_23_01_04_C288T_C300T_A303G_A317G: replacing with gap
##
##
## Unexpected N in novel sequence IGHV3_23_01_04_C300T_A303G_A317G: replacing with gap
```