


OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	3
1.4	Final three nucleotides: frequency of each observed triplet	3
1.5	CDR3 length distribution, in assignments to novel alleles	4
2	Variation from germline, in assignments to each allele	5
3	Allele usage in potential haplotype anchor genes	10
4	Haplotype plots	11
5	Configuration settings	12

1 Novel sequence analysis


1.1 End-nucleotide composition




1.2 Per-nucleotide consensus where previous nucleotides match the consensus







1.3 Whole-sequence composition of each assigned read















1.4 Final three nucleotides: frequency of each observed triplet



























1.5 CDR3 length distribution, in assignments to novel alleles



















2 Variation from germline, in assignments to each allele












3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJEB26509/S22/S22/22_IGK/22_IGK/b9/67e7cdb94c1d83d1fe52c3c9983e
##
## Germline reference file: /misc/work/jenkins/PRJEB26509/S22/S22/22_IGK/22_IGK/b9/67e7cdb94c1d83d1fe52
##
## Novel allele file: /misc/work/jenkins/PRJEB26509/S22/S22/22_IGK/22_IGK/b9/67e7cdb94c1d83d1fe52c3c998
##
## Species: Homosapiens
##
## Chain: IGKV
##
## Segment: V
##
## Warning: Over 90% of reference genes are called in the repertoire. This suggests that either the ref
```