


OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	3
1.4	Final three nucleotides: frequency of each observed triplet	3
1.5	CDR3 length distribution, in assignments to novel alleles	4
2	Variation from germline, in assignments to each allele	5
3	Allele usage in potential haplotype anchor genes	8
4	Haplotype plots	9
5	Configuration settings	10

1 Novel sequence analysis


1.1 End-nucleotide composition




1.2 Per-nucleotide consensus where previous nucleotides match the consensus




1.3 Whole-sequence composition of each assigned read




1.4 Final three nucleotides: frequency of each observed triplet



























1.5 CDR3 length distribution, in assignments to novel alleles




2 Variation from germline, in assignments to each allele







3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJEB26509/S33/S33/33_IGK/33_IGK/07/25333c341c06208f33b1393fb291
##
## Germline reference file: /misc/work/jenkins/PRJEB26509/S33/S33/33_IGK/33_IGK/07/25333c341c06208f33b1
##
## Novel allele file: /misc/work/jenkins/PRJEB26509/S33/S33/33_IGK/33_IGK/07/25333c341c06208f33b1393fb2
##
## Species: Homosapiens
##
## Chain: IGKV
##
## Segment: V
```