


# OGRDBstats Report

## Contents


|          |                                                                                   |           |
|----------|-----------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Novel sequence analysis</b>                                                    | <b>2</b>  |
| 1.1      | End-nucleotide composition . . . . .                                              | 2         |
| 1.2      | Per-nucleotide consensus where previous nucleotides match the consensus . . . . . | 2         |
| 1.3      | Whole-sequence composition of each assigned read . . . . .                        | 2         |
| 1.4      | Final three nucleotides: frequency of each observed triplet . . . . .             | 2         |
| 1.5      | CDR3 length distribution, in assignments to novel alleles . . . . .               | 3         |
| <b>2</b> | <b>Variation from germline, in assignments to each allele</b>                     | <b>4</b>  |
| <b>3</b> | <b>Allele usage in potential haplotype anchor genes</b>                           | <b>9</b>  |
| <b>4</b> | <b>Haplotype plots</b>                                                            | <b>10</b> |
| <b>5</b> | <b>Configuration settings</b>                                                     | <b>11</b> |

# 1 Novel sequence analysis


## 1.1 End-nucleotide composition




## 1.2 Per-nucleotide consensus where previous nucleotides match the consensus




## 1.3 Whole-sequence composition of each assigned read




## 1.4 Final three nucleotides: frequency of each observed triplet



























## 1.5 CDR3 length distribution, in assignments to novel alleles

















## 2 Variation from germline, in assignments to each allele














### 3 Allele usage in potential haplotype anchor genes



## 4 Haplotype plots

## 5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/PRJEB26509/S84/S84/84_IGK/84_IGK/5d/f181c22c7290cfb7cc64384afb87
##
## Germline reference file: /misc/work/jenkins/PRJEB26509/S84/S84/84_IGK/84_IGK/5d/f181c22c7290cfb7cc64
##
## Novel allele file: /misc/work/jenkins/PRJEB26509/S84/S84/84_IGK/84_IGK/5d/f181c22c7290cfb7cc64384afb
##
## Species: Homosapiens
##
## Chain: IGKV
##
## Segment: V
##
## Warning: Over 90% of reference genes are called in the repertoire. This suggests that either the ref
```