






OGRDBstats Report

Contents


1	Novel sequence analysis	2
1.1	End-nucleotide composition	2
1.2	Per-nucleotide consensus where previous nucleotides match the consensus	2
1.3	Whole-sequence composition of each assigned read	2
1.4	Final three nucleotides: frequency of each observed triplet	2
1.5	CDR3 length distribution, in assignments to novel alleles	3
2	Variation from germline, in assignments to each allele	4
3	Allele usage in potential haplotype anchor genes	9
4	Haplotype plots	10
5	Configuration settings	11



























1.5 CDR3 length distribution, in assignments to novel alleles


















2 Variation from germline, in assignments to each allele












3 Allele usage in potential haplotype anchor genes



4 Haplotype plots

5 Configuration settings

```
## Repertoire file: /misc/work/jenkins/unpublished_PRJEB58016/I35/I35/I35_IGK/I35_IGK/3e/f6d359b6c71859
##
## Germline reference file: /misc/work/jenkins/unpublished_PRJEB58016/I35/I35/I35_IGK/I35_IGK/3e/f6d359
##
## Novel allele file: /misc/work/jenkins/unpublished_PRJEB58016/I35/I35/I35_IGK/I35_IGK/3e/f6d359b6c718
##
## Species: Homosapiens
##
## Chain: IGKV
##
## Segment: V
##
## Warning: Over 90% of reference genes are called in the repertoire. This suggests that either the ref
```