

# The Foundations: Logic and Proofs

Chapter 1, Part I: Propositional Logic

# Discrete Mathematics

Discrete mathematics is the part of mathematics devoted to the study of discrete objects (Kenneth H. Rosen, 7th edition).

- How to think (and argument) mathematically.
- Learn mathematical facts and their applications.
- **Discrete mathematics** is the mathematical study of properties, and relationships among discrete objects.
- **Discrete mathematics** is the study of mathematical structures that are fundamentally discrete rather than continuous.

# Discrete Vs Continuous

- **Continuous Data:** A set of data is said to be continuous if the values belonging to the set can take on any value within a finite or infinite interval.
  - **Continuous data** is information that can be measured on a continuum or scale. , e.g., [ 0, 70].
  - **Continuous data** can have almost any numeric value and can be meaningfully subdivided into finer and finer increments, depending upon the precision of the measurement system.
- 
- **Discrete Data:** A set of data is said to be discrete if the values belonging to the set are distinct and separate. It is counted e.g., {1,2,3,4,5,6}


# Discrete vs Continuous

- Examples of discrete Data

- Number of boys in the class.
  - Number of candies in a packet.
  - Number of suitcases lost by an airline.

- Examples of continuous Data

- Height of a person.
  - Time in a race.
  - Distance traveled by a car.



## Why DM?






- Digital computers are based on discrete “atoms” (bits).
- Therefore, both a computer’s
  - structure (circuits) and
  - operations (execution of algorithms)can be described by discrete mathematics.

# Applications

- How many ways are there to choose a valid password on a computer system?
- What is the probability of winning a lottery?
- Is there a link between two computers in a network?
- How can I identify spam e-mail messages?
- How can I encrypt a message so that no unintended recipient can read it?
- What is the shortest path between two cities using a transportation system?
- How can a list of integers be sorted so that the integers are in increasing order?
- How many steps are required to do such a sorting?
- How can it be proved that a sorting algorithm correctly sorts a list?
- How can a circuit that adds two integers be designed?
- How many valid Internet addresses are there?

# Course Outline

- Logic and Proofs
- Sets and Functions
- Relations
- Counting and Recurrence Relations
- Graphs
- Trees
- Number Theory
- Discrete Probability and Finite State Automata







## **TEXT Book:**

- Discrete Mathematics & its Applications, 7<sup>th</sup> edition. By Kenneth H. Rosen.

## **References Book:**

- Invitation to Discrete Maths, 2nd edition. By Matousek and Nesetril.
- Discrete Mathematics. By Lovasz, Pelikan and Vesztergombi.


# Logic



- Logic is fundamental because it allows us to understand meaning of statements, deduce information about mathematical structures and uncover further structures.
- The rules of logic specify the meaning of mathematical statements.
- These rules are used to distinguish between valid and invalid arguments.


# Applications: Logic

## Hardware and software specifications




One-bit Full Adder  
with  
Carry-In and

Carry-Out



Formal: Input\_wire\_A  
value in {0, 1}



Example 1: Adder


### Example 2: System Specification:

- The router can send packets to the edge system only if it supports the new address space.
- For the router to support the new address space it's necessary that the latest software release be installed.
- The router can send packets to the edge system if the latest software release is installed.
- The router does not support the new address space.

How to write these specifications in a rigorous / formal way? Use Logic


# Applications: Number Theory

## RSA and Public-key Cryptography




Alice and Bob have never met but they would like to exchange a message. Eve would like to eavesdrop.  
E.g. between you and the Bank of America.

They could come up with a good encryption algorithm and exchange the **encryption key** – but how to do it without Eve getting it? (If Eve gets it, all security is lost.)





CS folks found the solution:  
*public key encryption*. Quite remarkable that is feasible.

# Applications: Coloring a Map



How to color this map so that no two adjacent regions have the same color?

# Applications: Graph representation



Coloring the nodes of the graph:

What's the minimum number of colors such that any two nodes connected by an edge have different colors?

# Applications: Scheduling of Final Exams

- How can the final exams at FAST be scheduled so that no student has two exams at the same time?

## Graph:

A vertex correspond to a course.


An edge between two vertices denotes that there is at least one common

student in the courses they represent.

Each time slot for a final exam is represented by a different color.

A coloring of the graph corresponds to a valid schedule of the exams.

# Applications: Scheduling of Final Exams



What are the constraints between courses?  
Find a valid coloring

# Logic

- Logic is fundamental because it allows us to understand meaning of statements, deduce information about mathematical structures and uncover further structures.
- The rules of logic specify the meaning of mathematical statements.
- These rules are used to distinguish between valid and invalid arguments.

# Propositional Logic

## Introduction

- A **proposition** is a **declarative sentence** (a sentence that declares a fact) that is either **true or false**, but not both.
- Are the following sentences propositions?
  - Islamabad is the capital of Pakistan. **(Yes)**
  - Read this carefully. **(No)**
  - $1+2=3$  **(Yes)**
  - $x+1=2$  **(No)**
  - What time is it? **(No)**

# Proposition

## Definition

**proposition** (or **statement**):

a declarative sentence that is either true or false

- **law of the excluded middle:**

a proposition cannot be partially true or partially false

- **law of contradiction:**

a proposition cannot be both true and false

## propositions

- The Moon revolves around the Earth.
- Elephants can fly.
- $3 + 8 = 11$

## not propositions

- What time is it?
- Exterminate!
- $x < 43$

# Examples: Propositions

Is the following sentence a proposition? If it is a proposition, determine whether it is true or false.

Islamabad is the capital of Pakistan.

This makes a declarative statement, and hence is a proposition. The proposition is TRUE (T).

Can Ali come with you?.

This is a question not the declarative sentence and hence not a proposition.



Take two aspirins.

This is an imperative sentence not the declarative sentence and therefore not a proposition.

$$x+4 > 9.$$

Because this is true for certain values of  $x$  (such as  $x = 6$ ) and false for other values of  $x$  (such as  $x = 5$ ), it is not a proposition.

He is a college student.

Because truth or falsity of this proposition depend on the reference for the pronoun *he*. it is not a proposition.

# Activity 1



**Write down at least 5 examples of propositions and  
Non-propositions.**

# Propositional Variable

- **propositional variable:**  
a name that represents the proposition


## examples

- $p_1$ : The Moon revolves around the Earth. ( $T$ )
- $p_2$ : Elephants can fly. ( $F$ )
- $p_3$ :  $3 + 8 = 11$  ( $T$ )

## Notations

- The small letters are commonly used to denote the propositional variables, that is, variables that represent propositions, such as,  $p, q, r, s, \dots$
- The **truth value of a proposition** is true, denoted by **T or 1**, if it is a true proposition and false, denoted by **F or 0**, if it is a false proposition.

# Compound Propositions



Logical operators are used to form new propositions also called compound propositions from two or more existing propositions.

- compound propositions are obtained by applying logical operators

The logical operators are also called connectives.

- truth table:

a table that lists the truth value of the compound proposition for all possible values of its variables

Propositional Logic – the area of logic that deals with propositions

# 1. Negation:

## DEFINITION 1:

Let  $p$  be a proposition. The negation of  $p$ , denoted by  $\neg p$ , is the statement “It is not the case that  $p$ .”

The proposition  $\neg p$  is read “not  $p$ .” The truth value of the negation of  $p$ ,  $\neg p$  is the opposite of the truth value of  $p$ .

- Examples
  - Find the negation of the proposition “Today is Friday.” and express this in simple English.  
**Solution:** The negation is “It is not the case that *today is Friday*.”  
In simple English, “Today is not Friday.” or “It is not Friday today.”
  - Find the negation of the proposition “At least 10 mm of rain fell today in Karachi.” and express this in simple English.  
**Solution:** The negation is “It is not the case that *at least 10 mm of rain fell today in Karachi*.”  
In simple English, “Less than 10 mm of rain fell today in Karachi.”

# Negation:

- Note: Always assume fixed times, fixed places, and particular people unless otherwise noted.
- Truth table:

The Truth Table for the Negation of a Proposition.

$p$	$\neg p$
T	F
F	T

## examples

- $\neg p_1$ : The Moon does not revolve around the Earth.  
 $\neg T : F$
- $\neg p_2$ : Elephants cannot fly.  
 $\neg F : T$

# 2. Conjunction:

## DEFINITION 2

Let  $p$  and  $q$  be propositions. The *conjunction* of  $p$  and  $q$ , denoted by  $p \wedge q$ , is the proposition “ $p$  and  $q$ ”. The conjunction  $p \wedge q$  is true when both  $p$  and  $q$  are true and is false otherwise.

- **Examples**

- Find the conjunction of the propositions  $p$  and  $q$  where  $p$  is the proposition “Today is Friday.” and  $q$  is the proposition “It is raining today.”, and the truth value of the conjunction.

**Solution:** The conjunction is the proposition “Today is Friday and it is raining today.” The proposition is true on rainy Fridays.

$p \wedge q$		
$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

### examples

- $p_1 \wedge p_2$ : The Moon revolves around the Earth and elephants can fly.  
 $T \wedge F : F$
- $p_1 \wedge p_3$ : The Moon revolves around the Earth and  $3 + 8 = 11$ .  
 $T \wedge T : T$


### 3. Disjunction:

#### DEFINITION 3

Let  $p$  and  $q$  be propositions. The *disjunction* of  $p$  and  $q$ , denoted by  $p \vee q$ , is the proposition “ $p$  or  $q$ ”. The *disjunction*  $p \vee q$  is false when both  $p$  and  $q$  are false and is true otherwise.

- Note:
  - inclusive or* : The disjunction is true when at least one of the two propositions is true.
    - E.g. “Students who have taken calculus or computer science can take this class.” – those who take one or both classes.
  - exclusive or* : The disjunction is true only when one of the proposition is true.
    - E.g. “Students who have taken calculus or computer science, **but not both**, can take this class.” – only those who take one of them.
- Definition 3 uses *inclusive or*.

# Disjunction (OR)



$p \vee q$

$p$	$q$	$p \vee q$
$T$	$T$	$T$
$T$	$F$	$T$
$F$	$T$	$T$
$F$	$F$	$F$

## example

- $p_1 \vee p_2$ : The Moon revolves around the Earth or elephants can fly.  
 $T \vee F : T$

## 4. Exclusive OR:

### DEFINITION 4

Let  $p$  and  $q$  be propositions. The *exclusive or* of  $p$  and  $q$ , denoted by  $p \oplus q$ , is the proposition that is true when exactly one of  $p$  and  $q$  is true and is false otherwise.

The Truth Table for the Conjunction of Two Propositions.

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

The Truth Table for the Disjunction of Two Propositions.

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

The Truth Table for the Exclusive Or (XOR) of Two Propositions.

$p$	$q$	$p \oplus q$
T	T	F
T	F	T
F	T	T
F	F	F

## Examples

1. Find the *exclusive or* of the propositions  $p$  and  $q$ , where

$p$  : Atif will pass the course CSC102.

$q$  : Atif will fail the course CSC102.

The *exclusive or* is

$p \oplus q$  : Atif will pass or fail the course CSC102.

---

The following proposition uses the (English) connective “or”. Determine from the context whether “or” is intended to be used in the inclusive or exclusive sense.

1. “Nabeel has one or two brothers”.

A person cannot have both one and two brothers.  
Therefore, “or” is used in the exclusive sense.

## Examples (OR vs XOR)

2. To register for BSC you must have passed the qualifying exam or be listed as an Math major.

Presumably, if you have passed the qualifying exam and are also listed as an Math major, you can still register for BCS. Therefore, “or” is inclusive.

# 5. Implication / Conditional Statements:

## DEFINITION 5

Let  $p$  and  $q$  be propositions. The *conditional statement*  $p \rightarrow q$ , is the proposition “if  $p$ , then  $q$ .” The conditional statement  $p \rightarrow q$  is false when  $p$  is true and  $q$  is false, and true otherwise. In the conditional statement  $p \rightarrow q$ ,  $p$  is called the *hypothesis* (or *antecedent* or *premise*) and  $q$  is called the *conclusion* (or *consequence*).

- A conditional statement is also called an implication.
- Example: “If I am elected, then I will lower taxes.”       $p \rightarrow q$

implication:

elected, lower taxes.

T      T      | T

not elected, lower taxes.

F      T      | T

not elected, not lower taxes.

F      F      | T

elected, not lower taxes.

T      F      | F

# Example: Conditional Statements

- Example:
  - Let  $p$  be the statement “Maria learns discrete mathematics.” and  $q$  the statement “Maria will find a good job.” Express the statement  $p \rightarrow q$  as a statement in English.

**Solution:** Any of the following -

“If Maria learns discrete mathematics, then she will find a good job.

“Maria will find a good job when she learns discrete mathematics.”

“For Maria to get a good job, it is sufficient her to learn discrete mathematics.”

“Maria will find a good job unless she does not learn discrete mathematics.”

# Examples: Implication

Examples of implications:

If you stand in the rain, then you'll get wet.

If you got an A in this class, I gave you \$5.

An implication  $P \implies Q$  is false only when  $P$  is true and  $Q$  is false. For example, the first statement would be false only if you stood in the rain but didn't get wet. The second statement above would be false only if you got an "A," yet I didn't give you \$5.

Here is the truth table for  $P \implies Q$ :

$P$	$Q$	$P \implies Q$	$\neg P \vee Q$
$T$	$T$	$T$	$T$
$T$	$F$	$F$	$F$
$F$	$T$	$T$	$T$
$F$	$F$	$T$	$T$

---

<sup>1</sup> $P$  is also called the *antecedent* and  $Q$  the *consequent*.

# Examples: Implication

Note that  $P \implies Q$  is always true when  $P$  is false. This means that many statements that sound nonsensical in English are true, mathematically speaking. Examples are statements like: “If pigs can fly, then horses can read” or “If 14 is odd then  $1 + 2 = 18$ .“ When an implication is stupidly true because the hypothesis is false, we say that it is **vacuously true**. Note also that  $P \implies Q$  is logically equivalent to  $\neg P \vee Q$ , as can be seen in the above truth table.


$P \implies Q$  is the most common form mathematical theorems take. Here are some of the different ways of saying it:

- (1) If  $P$ , then  $Q$ .
- (2)  $Q$  if  $P$ .
- (3)  $P$  only if  $Q$ .
- (4)  $P$  is sufficient for  $Q$ .
- (5)  $Q$  is necessary for  $P$ .

## Some other cases of implications:

“if $p$ , then $q$ ”	“ $p$ implies $q$ ”
“if $p, q$ ”	“ $p$ only if $q$ ”
“ $p$ is sufficient for $q$ ”	“a sufficient condition for $q$ is $p$ ”
“ $q$ if $p$ ”	“ $q$ whenever $p$ ”
“ $q$ when $p$ ”	“ $q$ is necessary for $p$ ”
“a necessary condition for $p$ is $q$ ”	“ $q$ follows from $p$ ”
“ $q$ unless $\neg p$ ”	

# Implication Example



- "If I weigh over 70 kg, then I will exercise."

## Implication Example

## Implication Examples

- "If I weigh over 70 kg, then I will exercise."

- $p_4: 3 < 8, p_5: 3 < 14, p_6: 3 < 2, p_7: 8 < 6$

- $p$ : I weigh over 70 kg.
- $q$ : I exercise.
- when is this claim false?

$p \rightarrow q$		
$p$	$q$	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

- $p_4 \rightarrow p_5$ :  
if  $3 < 8$ , then  $3 < 14$   
 $T \rightarrow T : T$
- $p_6 \rightarrow p_4$ :  
if  $3 < 2$ , then  $3 < 8$   
 $F \rightarrow T : T$
- $p_4 \rightarrow p_6$ :  
if  $3 < 8$ , then  $3 < 2$   
 $T \rightarrow F : F$
- $p_6 \rightarrow p_7$ :  
if  $3 < 2$ , then  $8 < 6$   
 $F \rightarrow F : T$

- Other conditional statements:

- Converse of  $p \rightarrow q$  :  $q \rightarrow p$
- Contrapositive of  $p \rightarrow q$  :  $\neg q \rightarrow \neg p$
- Inverse of  $p \rightarrow q$  :  $\neg p \rightarrow \neg q$

The contrapositive of “If you got an A in this class, I gave you \$5,” is “If I did not give you \$5, you didn’t get an A in this class.” The converse is “If I gave you \$5 you must have received an A in this class.” Does the contrapositive say the same thing as the original statement? Does the converse?

Let’s look at the truth table:

$P$	$Q$	$\neg P$	$\neg Q$	$P \Rightarrow Q$	$Q \Rightarrow P$	$\neg Q \Rightarrow \neg P$	$P \Leftrightarrow Q$
$T$	$T$	$F$	$F$	$T$	$T$	$T$	$T$
$T$	$F$	$F$	$T$	$F$	$T$	$F$	$F$
$F$	$T$	$T$	$F$	$T$	$F$	$T$	$F$
$F$	$F$	$T$	$T$	$T$	$T$	$T$	$T$

Note that the contrapositive of  $P \Rightarrow Q$  has the same truth values, while the converse does not. Many students unreasonably assume that the converse is true, but the above truth table shows that it is not necessarily the case. When two propositional forms have the same truth values, they are said to be **logically equivalent** – they mean the same thing. We’ll see next time how useful this can be for proving theorems.

# Converse, Contrapositive, and Inverse

- From  $p \rightarrow q$  we can form new conditional statements .
  - $q \rightarrow p$  is the **converse** of  $p \rightarrow q$
  - $\neg q \rightarrow \neg p$  is the **contrapositive** of  $p \rightarrow q$
  - $\neg p \rightarrow \neg q$  is the **inverse** of  $p \rightarrow q$

**Example:** Find the converse, inverse, and contrapositive of “It is raining is a sufficient condition for my not going to town.”

**Solution:**

**converse:** If I do not go to town, then it is raining.

**inverse:** If it is not raining, then I will go to town.

**contrapositive:** If I go to town, then it is not raining.


# Contrapositive:

- Contrapositive of  $p \rightarrow q$  is  $\neg q \rightarrow \neg p$
- Any proposition and its contrapositive are logically equivalent (have the same truth table values) – Check with the truth table.
- E.g. The contrapositive of “If you get 100% in this course, you will get an A+” is “If you do not get an A+ in this course, you did not get 100%”.

# Converse:

- Converse of  $p \rightarrow q$  is  $q \rightarrow p$
- Both are not logically equivalent.
- Ex 1: “If you get 100% in this course, you will get an A+” and “If you get an A+ in this course, you scored 100%” are not equivalent.
- Ex 2: If you won the lottery, you are rich.

## Inverse:



- Inverse of  $p \rightarrow q$  is  $\neg p \rightarrow \neg q$
- Both are not logically equivalent.
- Ex1 : “If you get 100% in this course, you will get an A+” and “If you didn’t 100%, then won’t have an A+ in this course.” are not equivalent.
- Ex2: You can not ride the roller coaster if you are under 4 feet. What is its inverse statement?

# Example of converse (1/2)

- Find the converse of the following statement:
- R: ‘Raining tomorrow is a sufficient condition for my not going to town.’
- Step 1: Assign propositional variables to component propositions
- P: It will rain tomorrow
- Q: I will not go to town

## Example of converse (2/2)

- Step 2: Symbolize the assertion  $R: P \rightarrow Q$
- Step 3: Symbolize the converse  $Q \rightarrow P$
- Step 4: Convert the symbols back into words  
‘If I don’t go to town then it will rain tomorrow’

or

‘My not going to town is a sufficient condition for it raining tomorrow.’

# Practice questions

Let  $p$ ,  $q$ , and  $r$  be the propositions

$p$ : You get an A on the final exam.  $q$ : You do every exercise in this book.  $r$ : You get an A in this class.

Write these propositions using  $p$ ,  $q$ , and  $r$  and logical connectives.

- a) You get an A in this class, but you do not do every exercise in this book.
- b) You get an A on the final, you do every exercise in this book, and you get an A in this class.
- c) To get an A in this class, it is necessary for you to get an A on the final.
- d) You get an A on the final, but you don't do every exercise in this book; nevertheless, you get an A in this class.
- e) Getting an A on the final and doing every exercise in this book is sufficient for getting an A in this class.
- f) You will get an A in this class if and only if you either do every exercise in this book or you get an A on the final.

# Solution

a) You get an A in this class, but you do not do every exercise in this book.

Solution:  $r \wedge \neg q$

b) You get an A on the final, and you do every exercise in this book, and you get an A in this class.

Solution:  $p \wedge q \wedge r$

d) You get an A on the final, but you don't do every exercise in this book; nevertheless, you get an A in this class.

Solution:  $p \wedge \neg q \wedge r$

e) Getting an A on the final and doing every exercise in this book is sufficient for getting an A in this class.

Solution:  $(p \wedge q) \rightarrow r$

f) You will get an A in this class if and only if you either do every exercise in this book or you get an A on the final.

Solution:  $r \leftrightarrow (q \vee p)$

# Practice Question

Suppose that Smartphone A has 256MB RAM and 32GB ROM, and the resolution of its camera is 8 MP;

Smartphone B has 288 MB RAM and 64 GB ROM, and the resolution of its camera is 4 MP; and

Smartphone C has 128 MB RAM and 32 GB ROM, and the resolution of its camera is 5 MP.

Express the given statements using Logical Connectives. Also determine the truth value of each of these propositions.

- a) Smartphone B has the most RAM of these three smartphones.
- b) Smartphone C has more ROM or a higher resolution camera than Smartphone B.
- c) Smartphone B has more RAM, more ROM, and a higher resolution camera than Smartphone A.
- d) If Smartphone B has more RAM and more ROM than Smartphone C, then it also has a higher resolution camera.
- e) Smartphone A has more RAM than Smartphone B if and only if Smartphone B has more RAM than Smartphone A.

# Solution

.....

	Smart Phone A	Smart Phone B	Smart Phone C
RAM	256 MB	288 MB	128 MB
ROM	32 GB	64 GB	32 GB
Camera resolution	8 MP	4 MP	5 MP

# Solution

a) Smartphone B has the most RAM of these three smartphones.

Solution: True Statement: p

p: Smartphone B has the most RAM of these three smartphones.

b) Smartphone C has more ROM or a higher resolution camera than Smartphone B.

Solution: True Statement:  $p \vee q$  (p OR q)

First identify propositions:

p: C has more ROM than B. FALSE q: C has a higher resolution camera than B TRUE

c) Smartphone B has more RAM, more ROM, and a higher resolution camera than Smartphone A.

Solution: False Statement:  $p \wedge q \wedge r$  (p AND q AND r )

First identify propositions:

p: B has more RAM than A. TRUE q: B has more ROM than A. TRUE

r: B has a higher resolution camera than A. FALSE

Cont...

d) If Smartphone B has more RAM and more ROM than Smartphone C, then it also has a higher resolution camera.

Solution: False Statement:  $(p \wedge q) \rightarrow r$  ( IF (p AND q) THEN r )

p: B has more RAM than C. TRUE q: B has more ROM than C. TRUE

r: B has a higher resolution camera than C. FALSE

e) Smartphone A has more RAM than Smartphone B if and only if Smartphone B has more RAM than Smartphone A.

Solution: False Statement:  $(p \leftrightarrow q)$  (p IF AND ONLY IF q)

p: A has more RAM than B. FALSE q: B has more RAM than A. TRUE

# Practice Questions

Write each of these statements in the form “if p, then q” in English. [Hint: Refer to the list of common ways to express conditional statements provided in this section.]

- a) You send me an e-mail message only if I will remember to send you the address.
- b) To be a citizen of this country, it is sufficient that you were born in the United States.
- c) If you keep your textbook, it will be a useful reference in your future courses.
- d) The Red Wings will win the Stanley Cup if their goalie plays well.
- e) That you get the job implies that you had the best credentials.
- f) The beach erodes whenever there is a storm.
- h) You will reach the summit unless you begin your climb too late.

# Solution

a) You send me an e-mail message only if I will remember to send you the address. “p only if q”

Solution: If you send me an e-mail message, then I will remember to send you the address.

b) To be a citizen of this country, it is sufficient that you were born in the United States.

“a sufficient condition for q is p”

Solution: If you were born in the United States, then you can be a citizen of the United States.

c) If you keep your textbook, it will be a useful reference in your future courses.

Solution :If you keep your textbook, then it will be a useful reference in your future courses.

# Solution

d) The Red Wings will win the Stanley Cup if their goalie plays well. “q if p”

Solution: If the Red Wings's goalie plays well, they will win the Stanley Cup.

e) That you get the job implies that you had the best credentials. “p implies q”

Solution: If you get the job, then you had the best credentials.

f) The beach erodes whenever there is a storm. “q whenever p”

Solution: If there is a storm, then beach erodes.

h) You will reach the summit unless you begin your climb too late. “q unless  $\neg p$ ”

Solution: If you begin your climb too late, then you will not reach the summit.

because  $p =$  you not begin your climb too late

# 6. Bi-implications:

## DEFINITION 6

Let  $p$  and  $q$  be propositions. The *biconditional statement*  $p \leftrightarrow q$  is the proposition “ $p$  if and only if  $q$ .” The biconditional statement  $p \leftrightarrow q$  is true when  $p$  and  $q$  have the same truth values, and is false otherwise. Biconditional statements are also called *bi-implications*.

- $p \leftrightarrow q$  has the same truth value as  $(p \rightarrow q) \wedge (q \rightarrow p)$
  - “*if and only if*” can be expressed by “*iff*”
  - Example:
    - Let  $p$  be the statement “You can take the flight” and let  $q$  be the statement “You buy a ticket.” Then  $p \leftrightarrow q$  is the statement “You can take the flight if and only if you buy a ticket.”
- Implication:**
- If you buy a ticket you can take the flight.
- If you don't buy a ticket you cannot take the flight.

# Bi-implications:

If  $p$  denotes “I am at home.” and  $q$  denotes “It is raining.” then  $p \leftrightarrow q$  denotes “I am at home if and only if it is raining.”

If both  $P \Rightarrow Q$  and  $Q \Rightarrow P$  are true, then we say “ $P$  if and only if  $Q$ ” (abbreviated  $P$  iff  $Q$ ). Formally, we write  $P \Leftrightarrow Q$ .  $P$  if and only if  $Q$  is true only when  $P$  and  $Q$  have the same truth values.

For example, if we let  $P$  be “3 is odd,”  $Q$  be “4 is odd,” and  $R$  be “6 is even,” then  $P \Rightarrow R$ ,  $Q \Rightarrow P$  (vacuously), and  $R \Rightarrow P$ . Because  $P \Rightarrow R$  and  $R \Rightarrow P$ ,  $P$  if and only if  $R$ .

Given an implication  $P \Rightarrow Q$ , we can also define its

- (a) Contrapositive:  $\neg Q \Rightarrow \neg P$
- (b) Converse:  $Q \Rightarrow P$

The contrapositive of “If you got an A in this class, I gave you \$5,” is “If I did not give you \$5, you didn’t get an A in this class.” The converse is “If I gave you \$5 you must have received an A in this class.” Does the contrapositive say the same thing as the original statement? Does the converse?

The Truth Table for  
the Biconditional  $p$

$\leftrightarrow q$ .

$p$	$q$	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

# Expressing the Biconditional

- Some alternative ways “ $p$  if and only if  $q$ ” is expressed in English:
  - **$p$  is necessary and sufficient for  $q$**
  - **$if p$  then  $q$  , and conversely**
  - **$p$  iff  $q$**

**Without changing their meanings, convert each of the following sentences into a sentence having the form  
“p iff q”**

For a matrix to be invertible, it is necessary and sufficient that its determinant is not zero.

**Answer:** A matrix is invertible if and only if its determinant is not zero.

If  $xy = 0$  then  $x = 0$  or  $y = 0$ , and conversely.

**Answer:**  $xy = 0$  if and only if  $x = 0$  or  $y = 0$

For an occurrence to become an adventure, it is necessary and sufficient for one to recount it.

**Answer:** An occurrence becomes an adventure if and only if one recounts it.

# Truth Tables of Compound Propositions

- We can use connectives to build up complicated compound propositions involving any number of propositional variables, then use truth tables to determine the truth value of these compound propositions.
- Example: Construct the truth table of the compound proposition


$$(p \vee \neg q) \rightarrow (p \wedge q).$$

The Truth Table of $(p \vee \neg q) \rightarrow (p \wedge q)$ .					
$p$	$q$	$\neg q$	$p \vee \neg q$	$p \wedge q$	$(p \vee \neg q) \rightarrow (p \wedge q)$
T	T	F	T	T	T
T	F	T	T	F	F
F	T	F	F	F	T
F	F	T	T	F	F

# Example Truth Table

- Construct a truth table for  
 $p \vee q \rightarrow \neg r$

p	q	r	$\neg r$	$p \vee q$	$p \vee q \rightarrow \neg r$
T	T	T	F	T	F
T	T	F	T	T	T
T	F	T	F	T	F
T	F	F	T	T	T
F	T	T	F	T	F
F	T	F	T	T	T
F	F	T	F	F	T
F	F	F	T	F	T



## Problem:

- How many rows are there in a truth table with  $n$  propositional variables?

**Solution:**  $2^n$  We will see how to do this in Chapter 6.

- Note that this means that with  $n$  propositional variables, we can construct  $2^n$  distinct (i.e., not equivalent) propositions.

# Precedence of Logical Operators

- We can use parentheses to specify the order in which logical operators in a compound proposition are to be applied.
- To reduce the number of parentheses, the precedence order is defined for logical operators.

Precedence of Logical Operators.	
Operator	Precedence
$\neg$	1
$\wedge$	2
$\vee$	3
$\rightarrow$	4
$\leftrightarrow$	5

$$\text{E.g. } \neg p \wedge q = (\neg p) \wedge q$$

$$p \wedge q \vee r = (p \wedge q) \vee r$$

$$p \vee q \wedge r = p \vee (q \wedge r)$$

# Translating English Sentence

- (1) If  $P$ , then  $Q$ .
- (2)  $Q$  if  $P$ .
- (3)  $P$  only if  $Q$ .
- (4)  $P$  is sufficient for  $Q$ .
- (5)  $Q$  is necessary for  $P$ .

- Steps to convert an English sentence to a statement in propositional logic
  - Identify atomic propositions and represent using propositional variables.
  - Determine appropriate logical connectives
- “If I go to Harry’s or to the country, I will not go shopping.”
  - $p$ : I go to Harry’s
  - $q$ : I go to the country.
  - $r$ : I will go shopping.

If  $p$  or  $q$  then not  $r$ .  
$$(p \vee q) \rightarrow \neg r$$

# Translating English Sentences

- Example: How can this English sentence be translated into a logical expression?

“You can access the Internet from campus only if you are a computer science major or you are not a freshman.”

**Solution:** Let  $a$ ,  $c$ , and  $f$  represent “You can access the Internet from campus,” “You are a computer science major,” and “You are a freshman.” The sentence can be translated into:

$$a \rightarrow (c \vee \neg f).$$

- (1) If  $P$ , then  $Q$ .
- (2)  $Q$  if  $P$ .
- (3)  $P$  only if  $Q$ .
- (4)  $P$  is sufficient for  $Q$ .
- (5)  $Q$  is necessary for  $P$ .

# Translating English Sentences

- (1) If  $P$ , then  $Q$ .
- (2)  $Q$  if  $P$ .
- (3)  $P$  only if  $Q$ .
- (4)  $P$  is sufficient for  $Q$ .
- (5)  $Q$  is necessary for  $P$ .

- English (and every other human language) is often ambiguous. Translating sentences into compound statements removes the ambiguity.
- Example: How can this English sentence be translated into a logical expression?

“You cannot ride the roller coaster if you are under 4 feet tall unless you are older than 16 years old.”

**Solution:** Let  $q$ ,  $r$ , and  $s$  represent “You can ride the roller coaster,”

“You are under 4 feet tall,” and “You are older than 16 years old.” The sentence can be translated into:

$t \square \neg q$  where  $t = \neg s \square r$  or

$$(r \wedge \neg s) \rightarrow \neg q.$$

You cannot ride the coaster if You are under 4 feet tall and you are not older than 16 Years old.

# System Specifications

- System and Software engineers take requirements in English and express them in a precise specification language based on logic.

**Example:** Express in propositional logic:

“The automated reply cannot be sent when the file system is full”

Solution let  $p$  denote “The file system is full ,and  $q$  denotes the automated reply can be send then  $p \square \neg q$

but what if I suppose  $p$  “The automated reply can be sent” and  $q$  to “The file system is full.”

$$q \rightarrow \neg p$$

# 1.1 Propositional Logic

## Logic and Bit Operations

- Computers represent information using bits.
- A **bit** is a symbol with two possible values, 0 and 1.
- By convention, 1 represents T (true) and 0 represents F (false).
- A variable is called a Boolean variable if its value is either true or false.
- Bit operation – replace true by 1 and false by 0 in logical operations.

Table for the Bit Operators *OR*, *AND*, and *XOR*.

$x$	$y$	$x \vee y$	$x \wedge y$	$x \oplus y$
0	0	0	0	0
0	1	1	0	1
1	0	1	0	1
1	1	1	1	0

# 1.1 Propositional Logic

## DEFINITION 7

A *bit string* is a sequence of zero or more bits. The *length* of this string is the number of bits in the string.

- Example: Find the bitwise *OR*, bitwise *AND*, and bitwise *XOR* of the bit string 01 1011 0110 and 11 0001 1101.

**Solution:**

01 1011 0110

11 0001 1101

-----

11 1011 1111    bitwise *OR*

01 0001 0100    bitwise *AND*

10 1010 1011    bitwise *XOR*

# Propositional Equivalences

## DEFINITION 1






A compound proposition that is always true, no matter what the truth values of the propositions that occurs in it, is called a *tautology*.

A compound proposition that is always false is called a *contradiction*.

A compound proposition that is neither a tautology or a contradiction is called a *contingency*.

Examples of a Tautology and a Contradiction.

$p$	$\neg p$	$p \vee \neg p$	$p \wedge \neg p$
T	F	T	F
F	T	T	F



## Tautologies and Contradictions

- Tautology is a statement that is always true regardless of the truth values of the individual logical variables
- Examples:
- $R \vee (\neg R)$
- $\neg(P \wedge Q) \leftrightarrow (\neg P) \vee (\neg Q)$

# Tautologies and Contradictions

---

- A Contradiction is a statement that is always false regardless of the truth values of the individual logical variables

## Examples

- $R \wedge (\neg R)$
- $\neg(\neg(P \wedge Q) \leftrightarrow (\neg P) \vee (\neg Q))$
- The negation of any tautology is a contradiction, and the negation of any contradiction is a tautology.

# Propositional Equivalences

## DEFINITION 2

The compound propositions  $p$  and  $q$  are called *logically equivalent* if  $p \leftrightarrow q$  is a tautology. The notation  $p \equiv q$  denotes that  $p$  and  $q$  are logically equivalent.

- Compound propositions that have the same truth values in all possible cases are called **logically equivalent**.
- Example: Show that  $\neg p \vee q$  and  $p \rightarrow q$  are logically equivalent.

Truth Tables for  $\neg p \vee q$  and  $p \rightarrow q$  .

$p$	$q$	$\neg p$	$\neg p \vee q$	$p \rightarrow q$
T	T	F	T	T
T	F	F	F	F
F	T	T	T	T
F	F	T	T	T

# Propositional Equivalences

- In general,  $2^n$  rows are required if a compound proposition involves  $n$  propositional variables in order to get the combination of all truth values.
  - Prove that  $\neg(\neg p) \equiv p$

## Solution

$p$	$\neg p$	$\neg(\neg p)$
T	F	T
F	T	F

As you can see the corresponding truth values of  $p$  and  $\neg(\neg p)$  are same, hence equivalence is justified.

# Propositional Equivalences

## Example

Show that the proposition forms  $\neg(p \wedge q)$  and  $\neg p \vee \neg q$  are NOT logically equivalent.

$p$	$q$	$\neg p$	$\neg q$	$(p \wedge q)$	$\neg(p \wedge q)$	$\neg p \vee \neg q$
T	T	F	F	T	F	F
T	F	F	T	F	T	F
F	T	T	F	F	T	F
F	F	T	T	F	T	T

Here the corresponding truth values differ and hence equivalence does not hold

# Applications: Boolean Searches

- Logical connectives are used extensively in searches of large collections of information.
  - Example: indexes of Web pages.
- AND - used to match records that contain both of two search terms.
- OR - used to match one or both of two search terms.
- NOT - used to exclude a particular search term.
- Read about: Web Page Searching

# Applications: Logic Puzzles

- Puzzles (**important job interview question**) that can be solved using logical reasoning
- [Sm78] Smullyan: An island that has two kinds of inhabitants.
  - knights, who always tell the truth.
  - knaves, who always lie.
- You encounter two people A and B.
- What are A and B if:
  - A says “B is a knight” and
  - B says “The two of us are opposite types?”

## Example 1:

- $p$ : A is a knight       $\neg p$ : A is a knave
- $q$ : B is a knight       $\neg q$ : B is a knave
- Consider the possibility that A is a knight;
  - So,  $p$  is true. And he is telling truth.
  - So,  $q$  is true. So, A and B are the same type.
- However, if B is a knight, then B's statement that A and B are of opposite types, the statement  $(p \wedge \neg q) \vee (\neg p \wedge q)$ , would have to be true, which it is not, because A and B are both knights. Consequently, we can conclude that A is not a knight, that is, that  $p$  is false.

## Example 1: Solution (cont..)


- Consider the possibility that A is a knave,
  - everything a knave says is false; q is true, is a lie.
  - So, q is false. B is also a knave.
  - B's statement that A and B are opposite types is a lie, which is consistent with both A and B being knaves.
- We can conclude that both A and B are knaves.

## Example

2:

- A father tells his two children, a boy and a girl, to play in their backyard without getting dirty.
- However, while playing, both children get mud on their foreheads. When the children stop playing, the father says “At least one of you has a muddy forehead,” and then asks the children to answer “Yes” or “No” to the question: “Do you know whether you have a muddy forehead?”
- The father asks this question twice. What will the children answer each time this question is asked, assuming that a child can see whether his or her sibling has a muddy forehead, but cannot see his or her own forehead?
- Assume that both children are honest and that the children answer each question simultaneously.

## Example 2:



Solution:  $S$  denotes "Son has a muddy forehead" and  $D$  denotes "Daughter has a muddy forehead". The father states that  $S \vee D$  is True. Boy can know  $D$  is True but can't know  $S$ . Girl can know  $D$  is True but can't know  $S$ . So no for the first time. After that they can conclude that both  $D$  and  $S$  are True. Since one of them will say yes for the first time if one of  $D$  and  $S$  is not True.

## Example 3:

6. Determine whether these system specifications are consistent:
- ”the diagnostic message is stored in the buffer or it is retransmitted”
  - ”the diagnostic message is not stored in the buffer”
  - ”if the diagnostic message is stored in the buffer, then it is retransmitted”

Solution:  $p$  denotes ”the diagnostic message is stored in the buffer”,  $q$  denotes ”the diagnostic message is retransmitted”. Then the specifications can be written as  $p \vee q$ ,  $\neg p$  and  $p \rightarrow q$ .  $\neg p$  is True, so  $p$  is False.  $p \vee q$  is True, so  $q$  is True. So  $p \rightarrow q$  is True. They are consistent.

# Propositional Satisfiability

- A compound proposition is satisfiable if there is an assignment of truth values to its variables that makes it true.
- When no such assignments exists, that is, when the compound proposition is false for all assignments of truth values to its variables, the compound proposition is unsatisfiable.
- A truth table can be used to determine whether a compound proposition is satisfiable, or equivalently, whether its negation is a tautology.

## De Morgan's laws

De Morgan's laws state that:

The negation of an **and** proposition is logically equivalent to the **or** proposition in which each component is negated.

$$1. \neg(p \wedge q) \equiv \neg p \vee \neg q$$

The negation of an **or** proposition is logically equivalent to the **and** proposition in which each component is negated.

$$2. \neg(p \vee q) \equiv \neg p \wedge \neg q$$

# Propositional Equivalences

## Constructing New Logical Equivalences

- Example: Show that  $\neg(p \rightarrow q)$  and  $p \wedge \neg q$  are logically equivalent.

Solution:

$$\begin{aligned}\neg(p \rightarrow q) &\equiv \neg(\neg p \vee q) && \text{by example discussed in slide 66} \\ &\equiv \neg(\neg p) \wedge \neg q && \text{by the second De Morgan law} \\ &\equiv p \wedge \neg q && \text{by the double negation law}\end{aligned}$$

- Example: Show that  $(p \wedge q) \rightarrow (p \vee q)$  is a tautology.

Solution: To show that this statement is a tautology, we will use logical equivalences to demonstrate that it is logically equivalent to T.

$$\begin{aligned}(p \wedge q) \rightarrow (p \vee q) &\equiv \neg(p \wedge q) \vee (p \vee q) && \text{by example already discussed} \\ &\equiv (\neg p \vee \neg q) \vee (p \vee q) && \text{by the first De Morgan law} \\ &\equiv (\neg p \vee p) \vee (\neg q \vee q) && \text{by the associative and} \\ &&& \text{communicative law for disjunction} \\ &\equiv T \vee T \\ &\equiv T\end{aligned}$$

- Note: The above examples can also be done using truth tables.

## Applying De-Morgan's Law

Question: Negate the following compound Propositions

1. John is six feet tall and he weights at least 200 pounds.
2. The bus was late or Tom's watch was slow.

## Applying De-Morgan's Law

Question: Negate the following compound Propositions

1. John is six feet tall and he weights at least 200 pounds.
2. The bus was late or Tom's watch was slow.

## Solution

- a) John is not six feet tall or he weighs less than 200 pounds.
- b) The bus was not late and Tom's watch was not slow.

## Inequalities and De Morgan's Laws

Question Use De Morgan's laws to write the negation of

$$-1 < x \leq 4$$

Solution: The given proposition is equivalent to

$$-1 < x \text{ and } x \leq 4,$$

By De Morgan's laws, the negation is

$$-1 \geq x \text{ or } x > 4.$$

## Example

Show that the proposition form  $p \vee \neg p$  is a tautology and the proposition form  $p \wedge \neg p$  is a contradiction.

$p$	$\neg p$	$p \vee \neg p$	$p \wedge \neg p$
T	F	T	F
F	T	T	F

Exercise: If  $t$  is a tautology and  $c$  is contradiction, show that  $p \vee t \equiv p$  and  $p \wedge c \equiv c$ ?

# Laws of Logic

## 1. Commutative laws

$$p \wedge q \equiv q \wedge p ; p \vee q \equiv q \vee p$$

## 2. Associative laws

$$p \wedge (q \wedge r) \equiv (p \wedge q) \wedge r ; p \vee (q \vee r) \equiv (p \vee q) \vee r$$


## 3. Distributive laws

$$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$$

$$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$$

# Laws of Logic

---



## 4. Identity laws

$$p \wedge t \equiv p ; p \vee c \equiv p$$

## 5. Negation laws

$$p \vee \neg p \equiv t ; p \wedge \neg p \equiv c$$

## 6. Double negation law

$$\neg(\neg p) \equiv p$$

## 7. Idempotent laws


$$p \wedge p \equiv p ; p \vee p \equiv p$$

## Laws of Logic

### 8. Universal bound laws

$$p \vee t \equiv t ; p \wedge c \equiv c$$

(empty)



### 9. Absorption laws

$$p \wedge (p \vee q) \equiv p ; p \vee (p \wedge q) \equiv p$$

### 10. Negation of $t$ and $c$

$$\neg t \equiv c ; \neg c \equiv t$$

## Exercise

Using laws of logic, show that

$$\neg(\neg p \wedge q) \wedge (p \vee q) \equiv p.$$

### Solution

Take  $\neg(\neg p \wedge q) \wedge (p \vee q)$

$$\equiv (\neg(\neg p) \vee \neg q) \wedge (p \vee q), \text{ (by De Morgan's laws)}$$

$$\equiv (p \vee \neg q) \wedge (p \vee q), \text{ (by double negative law)}$$

$$\equiv p \vee (\neg q \wedge q), \text{ (by distributive law)}$$

$$\equiv p \vee (q \wedge \neg q), \text{ (by the commutative law)}$$

$$\equiv p \vee c, \text{ (by the negation law)}$$

$$\equiv p, \text{ (by the identity law)}$$

Skill in simplifying proposition forms is useful in constructing logically efficient computer programs and in designing digital circuits.

# Exercise

Prove that  $\neg[r \vee (q \wedge (\neg r \rightarrow \neg p))] \equiv \neg r \wedge (p \vee \neg q)$

$$\begin{aligned}& \neg[r \vee (q \wedge (\neg r \rightarrow \neg p))] \\&\equiv \neg r \wedge \neg(q \wedge (\neg r \rightarrow \neg p)), \\&\equiv \neg r \wedge \neg(q \wedge (\neg r \vee \neg p)), \\&\equiv \neg r \wedge \neg(q \wedge (r \vee \neg p)), \\&\equiv \neg r \wedge (\neg q \vee \neg(r \vee \neg p)), \\&\equiv \neg r \wedge (\neg q \vee (\neg r \wedge p)), \\&\equiv (\neg r \wedge \neg q) \vee (\neg r \wedge (\neg r \wedge p)), \\&\equiv (\neg r \wedge \neg q) \vee ((\neg r \wedge \neg r) \wedge p), \\&\equiv (\neg r \wedge \neg q) \vee (\neg r \wedge p), \\&\equiv \neg r \wedge (\neg q \vee p), \\&\equiv \neg r \wedge (p \vee \neg q),\end{aligned}$$

De Morgan's law

Conditional rewritten as disjunction

Double negation law

De Morgan's law

De Morgan's law, double negation

Distributive law

Associative law

Idempotent law

Distributive law

Commutative law

# Course Description:

- This class teaches students
  - how to think logically and mathematically.
  - It stresses mathematical reasoning and different ways to solve problems.  
It enhance your ability to formulate and solve applied problems, to analyze and interpret algorithms and functions and to use them effectively.
- Five important themes are interwoven in this class:
  - mathematical reasoning,
  - combinatorial analysis,
  - discrete structures,
  - algorithmic thinking,
  - applications and modeling

# The Foundations: Logic and Proofs

Chapter 1, Part II: Predicate Logic

# Summary

- Predicate Logic (First-Order Logic (FOL),  
Predicate Calculus)
  - The Language of Quantifiers
  - Logical Equivalences
  - Nested Quantifiers
  - Translation from Predicate Logic to English
  - Translation from English to Predicate Logic

# Predicates and Quantifiers

Section 1.4

# Section Summary

- Predicates
- Variables
- Quantifiers
  - Universal Quantifier
  - Existential Quantifier
- Negating Quantifiers
  - De Morgan's Laws for Quantifiers
- Translating English to Logic
- Logic Programming (*optional*)

# Propositional Logic Not Enough

- “Every computer connected to the university network is functioning properly.”
- Is this a proposition ??
- No rules of propositional logic allow us to conclude the truth of the statement.
- “MATH3 is functioning properly,” where MATH3 is one of the computers connected to the university network.

# Predicates AND Predicate logic

- Predicates are the statements involving variable which are neither true nor false until or unless the values of variables are specified.
- F.g “x is animal”  
-> This sentence involves variables that's why we cannot identify truth of this statement until or unless give value to this variable.
- Here x is subject (some thing we are discussing about) “is an animal” is a predicate (refers to property that the subject of statement can have).
- The predicate logic a statement is divided into two parts.
  1. subject
  - 2.predicateusually we denote such statements with a shorthand notation.

“x is greater than 3” can be represented by  $G(x)$ , where  $G()$ denotes the predicate “greater than 3” and x denotes the subject or variable.

After assigning the value of variable x the statement  $G(x)$  become a proposition and has truth value (T or F).

$G(5)$  : 5 is greater than 3 => true

$G(2)$  : ???

# Introducing Predicate Logic

- Predicate logic uses the following new features:
  - Variables:  $x, y, z$
  - Predicates:  $P(x), M(x)$
  - Quantifiers (*to be covered in a few slides*):
- *Propositional functions* are a generalization of propositions.
  - They contain variables and a predicate, e.g.,  $P(x)$
  - Variables can be replaced by elements from their *domain*.

# Propositional Functions

- Propositional functions become propositions (and have truth values) when their variables are each replaced by a value from the *domain* (or *bound* by a quantifier, as we will see later).
- The statement  $P(x)$  is said to be the value of the propositional function  $P$  at  $x$ .
- For example, let  $P(x)$  denote “ $x > 0$ ” and the domain be the integers. Then:
  - P(-3) is false.
  - P(0) is false.
  - P(3) is true.
- Often the domain is denoted by  $U$ . So in this example  $U$  is the integers.

# Examples of Propositional Functions

- Let “ $x + y = z$ ” be denoted by  $R(x, y, z)$  and  $U$  (for all three variables) be the integers. Find these truth values:

$R(2, -1, 5)$

**Solution:** F

$R(3, 4, 7)$

**Solution:** T

$R(x, 3, z)$

**Solution: Not a Proposition**

- Now let “ $x - y = z$ ” be denoted by  $Q(x, y, z)$ , with  $U$  as the integers. Find these truth values:

$Q(2, -1, 3)$

**Solution:** T

$Q(3, 4, 7)$

**Solution:** F

$Q(x, 3, z)$

**Solution: Not a Proposition**

# Example of your book

**EXAMPLE 2** Let  $A(x)$  denote the statement “Computer  $x$  is under attack by an intruder.” Suppose that of the computers on campus, only CS2 and MATH1 are currently under attack by intruders. What are truth values of  $A(\text{CS1})$ ,  $A(\text{CS2})$ , and  $A(\text{MATH1})$ ?

*Solution:* We obtain the statement  $A(\text{CS1})$  by setting  $x = \text{CS1}$  in the statement “Computer  $x$  is under attack by an intruder.” Because CS1 is not on the list of computers currently under attack, we conclude that  $A(\text{CS1})$  is false. Similarly, because CS2 and MATH1 are on the list of computers under attack, we know that  $A(\text{CS2})$  and  $A(\text{MATH1})$  are true. ◀

# Compound Expressions

- Connectives from propositional logic carry over to predicate logic.
- If  $P(x)$  denotes “ $x > 0$ ,” find these truth values:
  - $P(3) \vee P(-1)$     **Solution:** T
  - $P(3) \wedge P(-1)$     **Solution:** F
  - $P(3) \rightarrow P(-1)$     **Solution:** F
  - $P(3) \rightarrow P(1)$     **Solution:** T
- Expressions with variables are not propositions and therefore do not have truth values. For example,
  - $P(3) \wedge P(y)$
  - $P(x) \rightarrow P(y)$Consider the statement  
If  $x > 0$  then  $x = x + 1$ ;  
-> When we have n variables in predicate function is n-ary predicate
- When used with quantifiers (to be introduced next), these expressions (propositional functions) become propositions.

# Quantifiers



Charles Peirce (1839-1914)

- Quantification express the extent to which a predicate is true over a range of elements(domain ).
- It tells for how many elements a given predicate is true.
- We need *quantifiers* to express the meaning of English words including “all ”and “some ”,are used to express the quantifier without giving an exact number.
- The two most important quantifiers are:
  - *Universal Quantifier*, which tells us that a predicate is true for every element under consideration “For all,” symbol:  $\forall$
  - *Existential Quantifier*, which tells us that there is one or more element under consideration “There exists,” symbol:  $\exists$
  - Example: let  $P(x)$  be a statement  $x+1 > x$   
what will be truth value for , $P(1) : 1+1 > 1$  ;  $P(2) : 2+1 > 2 \dots\dots$
  - This is true for all +ve integer  $x : P(x)$  is true or another way to saying of this sentence  $\forall x P(x)$  , for all quantifier here  $x$  represent positive integer.

## Example 2

- Let  $Q(x)$  be the statement  $x < 2$

$Q(1) : 1 < 2 , Q(2) : 2 < 2 , Q(3) : 3 < 2 \dots\dots$

this will be only true for  $Q=1$  and for rest of values it will be false.

-> Is there some values of  $x$  for which  $Q(x)$  is true.

$\exists x P(x)$ , there exist some  $x$  for which  $Q(x)$  is true.

# Quantifier

- We write as in  $\forall x P(x)$  and  $\exists x P(x)$ .
- $\forall x P(x)$  asserts  $P(x)$  is true for every  $x$  in the *domain*.
- $\exists x P(x)$  asserts  $P(x)$  is true for some  $x$  in the *domain*.
- The quantifiers are said to bind the variable  $x$  in these expressions.

# Quantifiers

- *Universal Quantifier*, “For all,” symbol:  $\forall$
- *Existential Quantifier*, “There exists,” symbol:  $\exists$

TABLE 1 Quantifiers.

<i>Statement</i>	<i>When True?</i>	<i>When False?</i>
$\forall xP(x)$	$P(x)$ is true for every $x$ .	There is an $x$ for which $P(x)$ is false.
$\exists xP(x)$	There is an $x$ for which $P(x)$ is true.	$P(x)$ is false for every $x$ .

# Remember

The truth  
depends

**Remark:** Generally, an implicit assumption is made that all domains of discourse for quantifiers are nonempty. Note that if the domain is empty, then  $\forall xP(x)$  is true for any propositional function  $P(x)$  because there are no elements  $x$  in the domain for which  $P(x)$  is false.

Besides “for all” and “for every,” universal quantification can be expressed in many other ways, including “all of,” “for each,” “given any,” “for arbitrary,” “for each,” and “for any.”

~~negation. There is not any reason to avoid using negation.~~

A statement  $\forall xP(x)$  is false, where  $P(x)$  is a propositional function, if and only if  $P(x)$  is not always true when  $x$  is in the domain. One way to show that  $P(x)$  is not always true when  $x$  is in the domain is to find a counterexample to the statement  $\forall xP(x)$ . Note that a single counterexample is all we need to establish that  $\forall xP(x)$  is false. Example 9 illustrates how counterexamples are used.

# Counter example

**EXAMPLE 9** Let  $Q(x)$  be the statement “ $x < 2$ .” What is the truth value of the quantification  $\forall x Q(x)$ , where the domain consists of all real numbers?

*Solution:*  $Q(x)$  is not true for every real number  $x$ , because, for instance,  $Q(3)$  is false. That is,  $x = 3$  is a counterexample for the statement  $\forall x Q(x)$ . Thus,

$$\forall x Q(x)$$

**EXAMPLE 10** Suppose that  $P(x)$  is “ $x^2 > 0$ .” To show that the statement  $\forall x P(x)$  is false where the universe of discourse consists of all integers, we give a counterexample. We see that  $x = 0$  is a counterexample because  $x^2 = 0$  when  $x = 0$ , so that  $x^2$  is not greater than 0 when  $x = 0$ . 

**EXAMPLE 12** What is the truth value of  $\forall x (x^2 \geq x)$  if the domain consists of all real numbers? What is the truth value of this statement if the domain consists of all integers?

# Universal Quantifier

$\forall x P(x)$  is read as “For all  $x$ ,  $P(x)$ ” or “For every  $x$ ,  $P(x)$ ”

## Examples:

- 1) If  $P(x)$  denotes “ $x > 0$ ” and  $U$  is the integers, then  $\forall x P(x)$  is false.
- 2) If  $P(x)$  denotes “ $x > 0$ ” and  $U$  is the positive integers, then  $\forall x P(x)$  is true.
- 3) If  $P(x)$  denotes “ $x$  is even” and  $U$  is the integers, then  $\forall x P(x)$  is false.

# Existential Quantifier

- $\exists x P(x)$  is read as “For some  $x$ ,  $P(x)$ ”, or as “There is an  $x$  such that  $P(x)$ ,” or “For at least one  $x$ ,  $P(x)$ .”

## Examples:

1. If  $P(x)$  denotes “ $x > 0$ ” and  $U$  is the integers, then  $\exists x P(x)$  is true. It is also true if  $U$  is the positive integers.
2. If  $P(x)$  denotes “ $x < 0$ ” and  $U$  is the positive integers, then  $\exists x P(x)$  is false.
3. If  $P(x)$  denotes “ $x$  is even” and  $U$  is the integers, then  $\exists x P(x)$  is true.

# Example

## Existential Quantifier - Examples

Example 2: Let  $P(x)$  be the statement " $x = x^2$ ." If the domain consists of the integers.  
What is the truth value of  $\exists x P(x)$ ?

Example 3: Let  $Q(x)$  be the statement " $x + 1 > 2x$ ." If the domain consists of all integers,  
What is the truth value of  $\exists x Q(x)$ ?

Example 4: Determine the truth value of each of these statements if the domain consists of all real numbers.

a)  $\exists x (x^3 = -1)$       b)  $\exists x (x^4 < x^2)$

### Homework Problem:

Determine the truth value for each of these statements if the domain consists of all real numbers.

a)  $\exists x (x^2 = 2)$

b)  $\exists x (x^3 = -1)$

c)  $\exists x (x^2 < 0)$

Cont...

Observe that the statement  $\exists xP(x)$  is false if and only if there is no element  $x$  in the domain for which  $P(x)$  is true. That is,  $\exists xP(x)$  is false if and only if  $P(x)$  is false for every element of the domain. We illustrate this observation in Example 14.

---

**EXAMPLE 14** Let  $Q(x)$  denote the statement “ $x = x + 1$ .” What is the truth value of the quantification  $\exists xQ(x)$ , where the domain consists of all real numbers?

*Solution:* Because  $Q(x)$  is false for every real number  $x$ , the existential quantification of  $Q(x)$ , which is  $\exists xQ(x)$ , is false. ◀

**Remark:** Generally, an implicit assumption is made that all domains of discourse for quantifiers are nonempty. If the domain is empty, then  $\exists xQ(x)$  is false whenever  $Q(x)$  is a propositional function because when the domain is empty, there can be no element  $x$  in the domain for which  $Q(x)$  is true.

# Uniqueness Quantifier (*optional*)

- $\exists !x P(x)$  means that  $P(x)$  is true for one and only one  $x$  in the universe of discourse.
- This is commonly expressed in English in the following equivalent ways:
  - “There is a unique  $x$  such that  $P(x)$ .”
  - “There is one and only one  $x$  such that  $P(x)$ ”
- Examples:
  1. If  $P(x)$  denotes “ $x + 1 = 0$ ” and  $U$  is the integers, then  $\exists !x P(x)$  is true.
  2. But if  $P(x)$  denotes “ $x > 0$ ,” then  $\exists !x P(x)$  is false.
- The uniqueness quantifier is not really needed as the restriction that there is a unique  $x$  such that  $P(x)$  can be expressed as:

$$\exists x (P(x) \wedge \forall y (P(y) \rightarrow y = x))$$

# Thinking about Quantifiers

- The universal quantification  $\forall x P(x)$  is conjunction same as for existential quantification is disjunction

$$P(x_1) \wedge P(x_2) \wedge \cdots \wedge P(x_n),$$

because this conjunction is true if and only if  $P(x_1), P(x_2), \dots, P(x_n)$  are all true.

**EXAMPLE 15** What is the truth value of  $\forall x P(x)$ , where  $P(x)$  is the statement “ $x^2 < 10$ ” and the domain consists of the positive integers not exceeding 4?

*Solution:* The statement  $\forall x P(x)$  is the same as the conjunction

$$P(1) \wedge P(2) \wedge P(3) \wedge P(4),$$

because the domain consists of the integers 1, 2, 3, and 4. Because  $P(4)$ , which is the statement “ $4^2 < 10$ ,” is false, it follows that  $\forall x P(x)$  is false. 

# Thinking about Quantifiers

- When the domain of discourse is finite, we can think of quantification as looping through the elements of the domain.
- To evaluate  $\forall x P(x)$  loop through all  $x$  in the domain.
  - If at every step  $P(x)$  is true, then  $\forall x P(x)$  is true.
  - If at a step  $P(x)$  is false, then  $\forall x P(x)$  is false and the loop terminates.
- To evaluate  $\exists x P(x)$  loop through all  $x$  in the domain.
  - If at some step,  $P(x)$  is true, then  $\exists x P(x)$  is true and the loop terminates.
  - If the loop ends without finding an  $x$  for which  $P(x)$  is true, then  $\exists x P(x)$  is false.
- Even if the domains are infinite, we can still think of the quantifiers this fashion, but the loops will not terminate in some cases.

# Properties of Quantifiers

- The truth value of  $\exists x P(x)$  and  $\forall x P(x)$  depend on both the propositional function  $P(x)$  and on the domain  $U$ .
- **Examples:**
  1. If  $U$  is the positive integers and  $P(x)$  is the statement “ $x < 2$ ”, then  $\exists x P(x)$  is true, but  $\forall x P(x)$  is false.
  2. If  $U$  is the negative integers and  $P(x)$  is the statement “ $x < 2$ ”, then both  $\exists x P(x)$  and  $\forall x P(x)$  are true.
  3. If  $U$  consists of 3, 4, and 5, and  $P(x)$  is the statement “ $x > 2$ ”, then both  $\exists x P(x)$  and  $\forall x P(x)$  are true. But if  $P(x)$  is the statement “ $x < 2$ ”, then both  $\exists x P(x)$  and  $\forall x P(x)$  are false.

# Precedence of Quantifiers

- The quantifiers  $\forall$  and  $\exists$  have higher precedence than all the logical operators.
- For example,  $\forall x P(x) \vee Q(x)$  means  $(\forall x P(x)) \vee Q(x)$
- $\forall x (P(x) \vee Q(x))$  means something different.
- Unfortunately, often people write  $\forall x P(x) \vee Q(x)$  when they mean  $\forall x (P(x) \vee Q(x))$ .

# Expressing Quantification in English

- Example:

**EXAMPLE 11** What does the statement  $\forall xN(x)$  mean if  $N(x)$  is “Computer  $x$  is connected to the network” and the domain consists of all computers on campus?

*Solution:* The statement  $\forall xN(x)$  means that for every computer  $x$  on campus, that computer  $x$  is connected to the network. This statement can be expressed in English as “Every computer on campus is connected to the network.” ◀

# Equivalences in Predicate Logic

- Statements involving predicates and quantifiers are *logically equivalent* if and only if they have the same truth value
  - for every predicate substituted into these statements and
  - for every domain of discourse used for the variables in the expressions.
- The notation  $S \equiv T$  indicates that  $S$  and  $T$  are logically equivalent.

# Example: Equivalences in Predicate Logic

- $\forall x (P(x) \wedge Q(x)) = \forall x P(x) \wedge \forall x Q(x)$   
Domain: all student of xyz university.  
 $P(x)$  x has studied discrete mathematics  
 $Q(x)$  x has scored more than 60% marks in exam
- The statement for logical expression on left side will be: Every student in the xyz university has studied discrete mathematics and has scored more than 60% marks in exam.
- The statement for logical expression on right side will be: Every student in the xyz university has studied discrete mathematics and Every student in the xyz university has scored more than 60% marks in exam.
- $\forall x (P(x) \vee Q(x)) = \forall x (P(x) \vee \forall x Q(x)) ??????? No$
- $\exists x (P(x) \wedge Q(x)) = \exists x (P(x) \wedge \exists x Q(x)) ??????$
- $\exists x (P(x) \vee Q(x)) = \exists x (P(x) \vee \exists x Q(x)) ????????$

# Equivalences in Predicate Logic

- $\exists x (P(x) \vee Q(x)) = \exists x (P(x) \vee \exists x Q(x)) ????????$
- The statement for logical expression on left side will be: some student in the xyz university has studied discrete mathematics or has scored more than 60% marks in exam.
- The statement for logical expression on right side will be : some student in the xyz university has studied discrete mathematics or some student in the xyz university has scored more than 60% marks in exam.

# Equivalences in Predicate Logic

- $\exists x (P(x) \wedge Q(x)) = \exists x (P(x) \wedge \exists x Q(x))$  no
- The statement for logical expression on left side will be:  
some student in the xyz university has studied discrete mathematics and has scored more than 60% marks in exam.
- The statement for logical expression on right side will be :  
some student in the xyz university has studied discrete mathematics and some student in the xyz university has scored more than 60% marks in exam.

# Translating from English to Logic

- Let  $P(x)$  be the “where the domain forstatement “x spend more than five hours every weekday in class x consists of all students.
- Express  $\forall x P(x)$  in English.
- Different representation of : for all  $x P(X)$   
for every  $x P(x)$

for each  $x P(x)$

$P(x)$

for any  $x$

All of  $x P(x)$

- For every  $x$  in the domain of all students , $x$  spends more than five hours every weekday in class.  
Every student spends more than 5 hours every weekday in class.
- Let  $N(x)$  be the statement “ $x$  has visited North Korea” where the domain consists of the students in your school. Express  $\forall x P(x)$  in English.

# Translating from English to Logic

- “For every student  $x$  in this class,  $x$  has studied calculus.”  $s(x)$
- “For every person  $x$ , person  $x$  is a student in this class, then  $x$  has studied calculus.”
- $\forall x(S(x) \rightarrow C(x))$ . Caution! Our statement cannot be expressed as  $\forall x(S(x) \wedge C(x))$ .
- $C(x))$  because this statement says that all people are students in this class and have studied calculus!]

# Translating from English to Logic

**Example 1:** Translate the following sentence into predicate logic: “Some student in this class has visited Mexico”

**Solution:**

The statement “Some student in this class has visited Mexico” means that “There is a student  $x$  in this class with the property that the student  $x$  has visited Mexico.”

We introduce  $M(x)$ , which is the statement “ $x$  has visited Mexico.”

If the domain for  $x$  consists of the students in this class, we can translate this first statement as  $\exists x M(x)$ .

However, if we are interested in people other than those in this class,

“There is a person  $x$  having the properties that  $x$  is a student in this class and  $x$  has visited Mexico.”

We introduce  $S(x)$  to represent

“ $x$  is a student in this class.” Our solution becomes  $\exists x(S(x) \wedge M(x))$ .

[Caution! Our statement cannot be expressed as  $\exists x(S(x) \rightarrow M(x))$ , which is true when there is someone not in the class because, in that case, for such a person  $x$ ,  $S(x) \rightarrow M(x)$  becomes either  $F \rightarrow T$  or  $F \rightarrow F = t$ , both of which are true.]

# Home task

- “For every  $x$  in this class,  $x$  has the property that  $x$  has visited Mexico or  $x$  has visited Canada.”

# Translating from English to Logic

**Example 2:** Translate the following sentence into predicate logic: “Some student in this class has taken a course in Java.”

**Solution:**

First decide on the domain  $U$ .

**Solution 1:** If  $U$  is all students in this class, translate as

$$\exists x J(x)$$

**Solution 1:** But if  $U$  is all people, then translate as

$$\exists x (S(x) \wedge J(x))$$

$\exists x (S(x) \rightarrow J(x))$  is not correct. What does it mean?

if we are interested in people other than those in this class, we look at the statement a little differently. Our statement can be expressed as “There is a person  $x$  having the properties that  $x$  is a student in this class and  $x$  has studied Java.”

In this case, the domain for the variable  $x$  consists of all people. We introduce  $S(x)$  to represent

“ $x$  is a student in this class.” Our solution becomes  $\exists x(S(x) \wedge J(x))$  because the statement is that there is a person  $x$  who is a student in this class and who has studied Java.

[Caution! Our statement cannot be expressed as  $\exists x(S(x) \rightarrow J(x))$ , which is true when there is someone not in the class because, in that case, for such a person  $x$ ,  $S(x) \rightarrow J(x)$  becomes either  $F \rightarrow T$  or  $F \rightarrow F$ , both of which are true.]

# Thinking about Quantifiers as Conjunctions and Disjunctions

- If the domain is finite, a universally quantified proposition is equivalent to a conjunction of propositions without quantifiers and an existentially quantified proposition is equivalent to a disjunction of propositions without quantifiers.
- If  $U$  consists of the integers 1, 2, and 3:

$$\forall x P(x) \equiv P(1) \wedge P(2) \wedge P(3)$$

$$\exists x P(x) \equiv P(1) \vee P(2) \vee P(3)$$

- Even if the domains are infinite, you can still think of the quantifiers in this fashion, but the equivalent expressions without quantifiers will be infinitely long.

# Negating Quantified Expressions

- Consider  $\forall x J(x)$   
“Every student in your class has taken a course in Java.”  
Here  $J(x)$  is “ $x$  has taken a course in Java” and  
the domain is students in your class.
- Negating the original statement gives “It is not  
the case that every student in your class has  
taken a course in Java.” This implies that  
“There is a student in your class who has not  
taken a course in Java.”  
Symbolically  $\neg \forall x J(x)$  and  $\exists x \neg J(x)$  are  
equivalent

# Negating Quantified Expressions (continued)

- Now Consider  $\exists x J(x)$   
“There is a student in this class who has taken a course in Java.”  
Where  $J(x)$  is “x has taken a course in Java.”
- Negating the original statement gives “It is not the case that there is a student in this class who has taken Java.” This implies that “Every student in this class has not taken a course in Java”  
Symbolically  $\neg \exists x J(x)$  and  $\forall x \neg J(x)$  are equivalent

# De Morgan's Laws for Quantifiers

- The rules for negating quantifiers are:

TABLE 2 De Morgan's Laws for Quantifiers.

Negation	Equivalent Statement	When Is Negation True?	When False?
$\neg\exists x P(x)$	$\forall x \neg P(x)$	For every $x$ , $P(x)$ is false.	There is an $x$ for which $P(x)$ is true.
$\neg\forall x P(x)$	$\exists x \neg P(x)$	There is an $x$ for which $P(x)$ is false.	$P(x)$ is true for every $x$ .

- The reasoning in the table shows that:

$$\neg\forall x P(x) \equiv \exists x \neg P(x)$$

$$\neg\exists x P(x) \equiv \forall x \neg P(x)$$

- These are important. You will use these.

# Home task

- What are the negations of the statements “There is an honest politician” and “All Americans eat cheeseburgers”?
- Show that  $\neg \forall x(P(x) \rightarrow Q(x))$  and  $\exists x(P(x) \wedge \neg Q(x))$  are logically equivalent.

# Examples of negation

Example 1: What is the negation of the statement "All dogs bark"?

Solution:

All dogs bark

Some dogs do not bark



$\forall \text{ dogs } d, d \text{ barks}$



$\exists \text{ dogs } d, d \text{ does not bark}$



$\neg (\forall \text{ dogs } d, d \text{ barks}) \rightarrow \exists \text{ dogs } d, \neg(d \text{ barks})$

Example 2: What is the negation of the statement "Most cars are inexpensive"?

Solution:

Most cars are inexpensive



$\exists \text{ cars } c, c \text{ is inexpensive}$

Not all cars are inexpensive



$\forall \text{ cars } c, c \text{ is not inexpensive}$



$\neg (\exists \text{ cars } c, c \text{ is inexpensive}) \rightarrow \forall \text{ cars } c, \neg(c \text{ is inexpensive})$

# Translation from English to Logic

## Examples:

1. “Some student in this class has visited Mexico.”

**Solution:** Let  $M(x)$  denote “ $x$  has visited Mexico” and  $S(x)$  denote “ $x$  is a student in this class,” and  $U$  be all people.

$$\exists x (S(x) \wedge M(x))$$

2. “Every student in this class has visited Canada or Mexico.”

**Solution:** Add  $C(x)$  denoting “ $x$  has visited Canada.”

$$\forall x (S(x) \rightarrow (M(x) \vee C(x)))$$

# Some Fun with Translating from English into Logical Expressions

- $U = \{\text{Nitwits, Blubbers, Oddments}\}$

$F(x)$ :  $x$  is a Nitwit

$S(x)$ :  $x$  is a Blubber

$T(x)$ :  $x$  is a Oddment

Translate “Everything is a Nitwit”

**Solution:**  $\forall x F(x)$

# Translation (cont)

- $U = \{\text{Nitwits, Blubbers, Oddments}\}$

$F(x)$ :  $x$  is a Nitwit

$S(x)$ :  $x$  is a Blubber

$T(x)$ :  $x$  is a Oddment

“Nothing is a Blubber.”

**Solution:**  $\neg \exists x S(x)$  What is this equivalent to?

**Solution:**  $\forall x \neg S(x)$

# Translation (cont)

- $U = \{\text{Nitwits, Blubbers, Oddments}\}$

$F(x)$ :  $x$  is a Nitwit

$S(x)$ :  $x$  is a Blubber

$T(x)$ :  $x$  is a Oddment

“All Nitwits are Blubbers.”

**Solution:**  $\forall x (F(x) \rightarrow S(x))$

# Translation (cont)

- $U = \{\text{Nitwits, Blubbers, Oddments}\}$

$F(x)$ :  $x$  is a Nitwit

$S(x)$ :  $x$  is a Blubber

$T(x)$ :  $x$  is a Oddment

“Some Nitwits are Oddments.”

**Solution:**  $\exists x (F(x) \wedge T(x))$

# Translation (cont)

- $U = \{\text{Nitwits, Blubbers, Oddments}\}$

$F(x)$ :  $x$  is a Nitwit

$S(x)$ :  $x$  is a Blubber

$T(x)$ :  $x$  is a Oddment

“No Blubber is a Oddment.”

**Solution:**  $\neg \exists x (S(x) \wedge T(x))$  What is this equivalent to?

**Solution:**  $\forall x (\neg S(x) \vee \neg T(x))$

# Translation (cont)

- $U = \{\text{Nitwits, Blubbers, Oddments}\}$

$F(x)$ :  $x$  is a Nitwit

$S(x)$ :  $x$  is a Blubber

$T(x)$ :  $x$  is a Oddment

“If any Nitwit is a Blubber then it is also a Oddment.”

**Solution:**  $\forall x ((F(x) \wedge S(x)) \rightarrow T(x))$

# System Specification Example

- Predicate logic is used for specifying properties that systems must satisfy.
- For example, translate into predicate logic:
  - “Every mail message larger than one megabyte will be compressed.”
  - “If a user is active, at least one network link will be available.”
- Decide on predicates and domains (left implicit here) for the variables:
  - Let  $L(m, y)$  be “Mail message  $m$  is larger than  $y$  megabytes.”
  - Let  $C(m)$  denote “Mail message  $m$  will be compressed.”
  - Let  $A(u)$  represent “User  $u$  is active.”
  - Let  $S(n, x)$  represent “Network link  $n$  is state  $x$ .  
$$\forall m(L(m, 1) \rightarrow C(m))$$
- Now we have:  
$$\exists u A(u) \rightarrow \exists n S(n, \text{available})$$

# Lewis Carroll Example



Charles Lutwidge  
Dodgson  
(AKA Lewis Carroll)  
(1832-1898)

- The first two are called *premises* and the third is called the *conclusion*.
  1. “All lions are fierce.”
  2. “Some lions do not drink coffee.”
  3. “Some fierce creatures do not drink coffee.”
- Here is one way to translate these statements to predicate logic. Let  $P(x)$ ,  $Q(x)$ , and  $R(x)$  be the propositional functions “ $x$  is a lion,” “ $x$  is fierce,” and “ $x$  drinks coffee,” respectively.
  1.  $\forall x (P(x) \rightarrow Q(x))$
  2.  $\exists x (P(x) \wedge \neg R(x))$
  3.  $\exists x (Q(x) \wedge \neg R(x))$
- Later we will see how to prove that the conclusion follows from the premises.

# Nested Quantifiers

Section 1.5

# Section Summary

- Nested Quantifiers
- Order of Quantifiers
- Translating from Nested Quantifiers into English
- Translating Mathematical Statements into Statements involving Nested Quantifiers.
- Translated English Sentences into Logical Expressions.
- Negating Nested Quantifiers.

# Nested Quantifiers

- Nested quantifiers are often necessary to express the meaning of sentences in English as well as important concepts in computer science and mathematics.

**Example:** “Every real number has an inverse” is

$$\forall x \exists y(x + y = 0)$$

where the domains of  $x$  and  $y$  are the real numbers.

- We can also think of within the scope of a quantifier as nested propositional functions:

$\forall x \exists y(x + y = 0)$  can be viewed as  $\forall x Q(x)$  where  $Q(x)$  is  $\exists y P(x, y)$  where  $P(x, y)$  is  $(x + y = 0)$

# Understanding Statements Involving Nested Quantifiers

Translate into English the statement

$$1. \forall x \exists y (x + y = 0)$$

says that for every real number  $x$  there is a real number  $y$  such that  $x + y = 0$ .

This states that every real number has an additive inverse.

$$2. \forall x \forall y \forall z (x + (y + z) = (x + y) + z)$$

is the associative law for addition of real numbers.

$$3. \forall x \forall y ((x > 0) \wedge (y < 0) \rightarrow (xy < 0)),$$

where the domain for both variables consists of all real numbers.

**Solution:** This statement says that for every real number  $x$  and for every real number  $y$ , if  $x > 0$  and  $y < 0$ , then  $xy < 0$ .

That is, this statement says that for real numbers  $x$  and  $y$ , if  $x$  is positive and  $y$  is negative, then  $xy$  is negative.

This can be stated more succinctly as “The product of a positive real number and a negative real number is always a negative real number.”

# Thinking of Nested Quantification

- Nested Loops

**Single quantifier: If at every step  $P(x)$  is true, then  $\forall x P(x)$  is true.**

- To see if  $\forall x \forall y P(x,y)$  is true, loop through the values of  $x$ :
  - At each step, loop through the values for  $y$ .
  - If for some pair of  $x$  and  $y$ ,  $P(x,y)$  is false, then  $\forall x \forall y P(x,y)$  is false and both the outer and inner loop terminate.  
 $\forall x \forall y P(x,y)$  is true if the outer loop ends after stepping through each  $x$ .
- To see if  $\forall x \exists y P(x,y)$  is true, loop through the values of  $x$ :
  - At each step, loop through the values for  $y$ .
  - The inner loop ends when a pair  $x$  and  $y$  is found such that  $P(x, y)$  is true.
  - If no  $y$  is found such that  $P(x, y)$  is true the outer loop terminates as  $\forall x \exists y P(x,y)$  has been shown to be false.  
 $\forall x \exists y P(x,y)$  is true if the outer loop ends after stepping through each  $x$ .
- If the domains of the variables are infinite, then this process can not actually be carried out.

# Order of Quantifiers

Many mathematical statements involve multiple quantifications of propositional functions involving more than one variable. It is important to note that the order of the quantifiers is important, unless all the quantifiers are universal quantifiers or all are existential quantifiers.

## Examples:

1. Let  $P(x,y)$  be the statement " $x + y = y + x$ ." Assume that  $U$  is the real numbers. Then  $\forall x \forall y P(x,y)$  and  $\forall y \forall x P(x,y)$  have the same truth value.  
The statement  $\forall y \forall x P(x, y)$  says "For all real numbers  $y$ , for all real numbers  $x$ ,  $x + y = y + x$ ."

This has the same meaning as the statement "For all real numbers  $x$ , for all real numbers  $y$ ,  $x + y = y + x$ ."

That is,  $\forall x \forall y P(x, y)$  and  $\forall y \forall x P(x, y)$  have the same meaning, and both are true.

# Order of Quantifiers

- Let  $Q(x,y)$  be the statement " $x + y = 0$ ." Assume that  $U$  is the real numbers. Then  $\forall x \exists y P(x,y)$  is true, but  $\exists y \forall x P(x,y)$  is false.

No matter what value of  $y$  is chosen, there is only one value of  $x$  for which  $x + y = 0$ . Because there is no real number  $y$  such that  $x + y = 0$  for all real numbers  $x$ , the statement  $\exists y \forall x Q(x, y)$  is false.

# Questions on Order of Quantifiers

**Example 1:** Let  $U$  be the real numbers,  
Define  $P(x,y) : x \cdot y = 0$

What is the truth value of the following:

1.  $\forall x \forall y P(x,y)$

**Answer:** False

2.  $\forall x \exists y P(x,y)$

**Answer:** True

3.  $\exists x \forall y P(x,y)$

**Answer:** True

4.  $\exists x \exists y P(x,y)$

**Answer:** True

# Questions on Order of Quantifiers

**Example 2:** Let  $U$  be the real numbers,  
Define  $P(x,y) : x / y = 1$

What is the truth value of the following:

1.  $\forall x \forall y P(x,y)$

**Answer:** False

2.  $\forall x \exists y P(x,y)$

**Answer:** True

3.  $\exists x \forall y P(x,y)$

**Answer:** False

4.  $\exists x \exists y P(x,y)$

**Answer:** True

# Quantifications of Two Variables

Statement	When True?	When False
$\forall x \forall y P(x, y)$ $\forall y \forall x P(x, y)$	$P(x, y)$ is true for every pair $x, y$ .	There is a pair $x, y$ for which $P(x, y)$ is false.
$\forall x \exists y P(x, y)$	For every $x$ there is a $y$ for which $P(x, y)$ is true.	There is an $x$ such that $P(x, y)$ is false for every $y$ .
$\exists x \forall y P(x, y)$	There is an $x$ for which $P(x, y)$ is true for every $y$ .	For every $x$ there is a $y$ for which $P(x, y)$ is false.
$\exists x \exists y P(x, y)$ $\exists y \exists x P(x, y)$	There is a pair $x, y$ for which $P(x, y)$ is true.	$P(x, y)$ is false for every pair $x, y$

# Translating Nested Quantifiers into English

**Example 1:** Translate the statement

$$\forall x (C(x) \vee \exists y (C(x) \wedge F(x, y)))$$

where  $C(x)$  is “ $x$  has a computer,” and  $F(x, y)$  is “ $x$  and  $y$  are friends,” and the domain for both  $x$  and  $y$  consists of all students in your school.

**Solution:** Every student in your school has a computer or has a friend who has a computer.

# Translating Mathematical Statements into Predicate Logic

**Example :** Translate “The sum of two positive integers is always positive” into a logical expression.

**Solution:**

1. Rewrite the statement to make the implied quantifiers and domains explicit:  
“For every two integers, if these integers are both positive, then the sum of these integers is positive.”

2. Introduce the variables  $x$  and  $y$ , and specify the domain, to obtain:  
“For all positive integers  $x$  and  $y$ ,  $x + y$  is positive.”

3. The result is:

$$\forall x \forall y ((x > 0) \wedge (y > 0) \rightarrow (x + y > 0))$$

where the domain of both variables consists of all integers

OR

Simply I can write  $\forall x \forall y (x+y > 0)$

# Translating English into Logical Expressions Example

**Example:** Use quantifiers to express the statement “There is a woman who has taken a flight on every airline in the world.”

**Solution:**

1. Let  $P(w,f)$  be “ $w$  has taken  $f$  ” and  $Q(f,a)$  be “ $f$  is a flight on  $a$  .”
2. The domain of  $w$  is all women, the domain of  $f$  is all flights, and the domain of  $a$  is all airlines.
3. Then the statement can be expressed as:

$$\exists w \exists f \forall a (P(w,f) \wedge Q(f,a))$$

# Questions on Translation from English

Choose the obvious predicates and express in predicate logic.

**Example 1:** “Everybody loves somebody.”

**Solution:**  $\forall x \exists y L(x,y)$

**Example 2:** “There is someone who is loved by everyone.”

**Solution:**  $\exists y \forall x L(x,y)$

**Example 3:** “There is someone who loves someone.”

**Solution:**  $\exists x \exists y L(x,y)$

**Example 4:** “Everyone loves himself”

**Solution:**  $\forall x L(x,x)$

**Example 5:** “Brothers are siblings.”

**Solution:**  $\forall x \forall y (B(x,y) \rightarrow S(x,y))$

**Example 6:** “Siblinghood is symmetric.”

**Solution:**  $\forall x \forall y (S(x,y) \rightarrow S(y,x))$

# Negating Nested Quantifiers

Example 1: Recall the logical expression developed three slides back:

$$\exists w \forall a \exists f (P(w,f) \wedge Q(f,a))$$

Part 1: Use quantifiers to express the statement that “There does not exist a woman who has taken a flight on every airline in the world.”

$$\text{Solution: } \neg \exists w \forall a \exists f (P(w,f) \wedge Q(f,a))$$

Part 2: Now use De Morgan’s Laws to move the negation as far inwards as possible.

Solution:

1.  $\neg \exists w \forall a \exists f (P(w,f) \wedge Q(f,a))$        $\neg \forall x P(x) \equiv \exists x \neg P(x)$
2.  $\forall w \neg \forall a \exists f (P(w,f) \wedge Q(f,a))$  by De Morgan’s for  $\exists$        $\neg \exists x P(x) \equiv \forall x \neg P(x)$
3.  $\forall w \exists a \neg \exists f (P(w,f) \wedge Q(f,a))$  by De Morgan’s for  $\forall$
4.  $\forall w \exists a \forall f \neg (P(w,f) \wedge Q(f,a))$  by De Morgan’s for  $\exists$
5.  $\forall w \exists a \forall f (\neg P(w,f) \vee \neg Q(f,a))$  by De Morgan’s for  $\wedge$ .

Part 3: Can you translate the result back into English?

Solution:

“For every woman there is an airline such that for all flights, this woman has not taken that flight or that flight is not on this airline”

# Some Questions about Quantifiers (optional)

- Can you switch the order of quantifiers?
  - Is this a valid equivalence?  $\forall x \forall y P(x, y) \equiv \forall y \forall x P(x, y)$   
**Solution:** Yes! The left and the right side will always have the same truth value. The order in which  $x$  and  $y$  are picked does not matter.  $\forall x \exists y P(x, y) \equiv \exists y \forall x P(x, y)$
  - Is this a valid equivalence?  
**Solution:** No! The left and the right side may have different truth values for some propositional functions for  $P$ . Try “ $x + y = 0$ ” for  $P(x, y)$  with  $U$  being the integers. The order in which the values of  $x$  and  $y$  are picked does matter.
- Can you distribute quantifiers over logical connectives?
  - Is this a valid equivalence?  
**Solution:** Yes! The left and the right side will always have the same truth value no matter what propositional functions are denoted by  $P(x)$  and  $Q(x)$ .
  - Is this a valid equivalence?  
**Solution:** No! The left and the right side may have different truth values. Pick “ $x$  is a fish” for  $P(x)$  and “ $x$  has scales” for  $Q(x)$  with the domain of discourse being all animals. Then the left side is false, because there are some fish that do not have scales. But the right side is true since not all animals are fish.

# The Foundations: Logic and Proofs

Chapter 1, Part III: Proofs

# Summary

- Valid Arguments and Rules of Inference
- Proof Methods
- Proof Strategies

# **Rules of Inference**

**Section 1.6**

# Section Summary

- Valid Arguments
- Inference Rules for Propositional Logic
- Using Rules of Inference to Build Arguments
- Rules of Inference for Quantified Statements
- Building Arguments for Quantified Statements

# Rule of inference in propositional logic

- **Premise:** is a proposition on the basis of which we would able to draw a conclusion. You can think of premise as an evidence or assumption.
- Therefore, initially we assume something is true and on the basis of that assumption, we draw some conclusion.
- **Conclusion:** is a proposition that is reached from the given set of premises.
- You can think of it as the result of the assumptions that we made in an argument.

if premise then conclusion

**Argument:** Sequences of statements that ends with a conclusion

OR

It is a set of one or more premises and a conclusion.

# Valid Argument

- Valid Argument: an argument is said to be valid if and only if it is not possible to make all premises true and a conclusion false.

Example 1 of an argument:

P1: “ If I love cat then I love dog”.

P2: “ I love cat”.

Conclusion: Therefore “ I love dog”.

Example 2 of an argument:

P1: “ If I love cat then I love dog”.

P2: “ I love dog”.

Conclusion: Therefore “ I love cat”.

# Valid Arguments

- We will show how to construct valid arguments in two stages; first for propositional logic and then for predicate logic. The rules of inference are the essential building block in the construction of valid arguments.

## 1. Propositional Logic

Inference Rules

## 2. Predicate Logic

Inference rules for propositional logic plus additional inference rules to handle variables and quantifiers.

# Valid Arguments (Propositional Logic)

- “If you have a current password, then you can log onto the network.”
  - “You have a current password.”
  - Therefore, “You can log onto the network.”
- 
- Sequence of propositions (argument) is valid; the conclusion must be true when both premise are true.

# Argument Form

- p: “You have a current password”
- q: “You can log onto the network.”
- the argument has the form

$$p \rightarrow q$$

$$p$$

$$\therefore q$$

the statement  $((p \rightarrow q) \wedge p) \rightarrow q$  is a tautology

# Argument Form

- p: “You have access to the network”
- q: “You can change your grade”
- the argument has the form

“If you have access to the network, then you can change your grade.”

“You have access to the network.”

∴ “You can change your grade.”

- The argument is valid, but if premise is false, we cannot conclude that the conclusion is true.

# Arguments in Propositional Logic

- A *argument* in propositional logic is a sequence of propositions. All but the final proposition are called *premises*. The last statement is the *conclusion*.
- The argument is valid if the premises imply the conclusion. An *argument form* is an argument that is valid no matter what propositions are substituted into its propositional variables.
- If the premises are  $p_1, p_2, \dots, p_n$  and the conclusion is  $q$  then  $(p_1 \wedge p_2 \wedge \dots \wedge p_n) \rightarrow q$  is a tautology.
- Inference rules are all argument simple argument forms that will be used to construct more complex argument forms.

# Rule of inference

- Rule of inference are the template for constructing valid arguments.  
This will help you to construct more complicated argument.
- Inference : Deriving conclusion from evidence.

**TABLE 1** Rules of Inference.

<i>Rule of Inference</i>	<i>Tautology</i>	<i>Name</i>
$\begin{array}{c} p \\ p \rightarrow q \\ \hline \therefore q \end{array}$	$(p \wedge (p \rightarrow q)) \rightarrow q$	Modus ponens
$\begin{array}{c} \neg q \\ p \rightarrow q \\ \hline \therefore \neg p \end{array}$	$(\neg q \wedge (p \rightarrow q)) \rightarrow \neg p$	Modus tollens
$\begin{array}{c} p \rightarrow q \\ q \rightarrow r \\ \hline \therefore p \rightarrow r \end{array}$	$((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$	Hypothetical syllogism
$\begin{array}{c} p \vee q \\ \neg p \\ \hline \therefore q \end{array}$	$((p \vee q) \wedge \neg p) \rightarrow q$	Disjunctive syllogism

**TABLE 1** Rules of Inference.

<i>Rule of Inference</i>	<i>Tautology</i>	<i>Name</i>
$\begin{array}{c} p \\ \hline \therefore p \vee q \end{array}$	$p \rightarrow (p \vee q)$	Addition
$\begin{array}{c} p \wedge q \\ \hline \therefore p \end{array}$	$(p \wedge q) \rightarrow p$	Simplification
$\begin{array}{c} p \\ q \\ \hline \therefore p \wedge q \end{array}$	$((p) \wedge (q)) \rightarrow (p \wedge q)$	Conjunction
$\begin{array}{c} p \vee q \\ \neg p \vee r \\ \hline \therefore q \vee r \end{array}$	$((p \vee q) \wedge (\neg p \vee r)) \rightarrow (q \vee r)$	Resolution

# Rules of Inference for Propositional Logic: *Modus Ponendo Ponens*

$$\frac{P \rightarrow Q, P}{\therefore Q}$$

- Latin for “the way that affirms by affirming”
- **Implication elimination**
- Abbreviated to **MP** or ***modus ponens***

- whenever  $p \rightarrow q$  is true
- and p is true,
- q must also be true.

$p$	$q$	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

# Modus Ponens

$$\frac{p \rightarrow q}{\begin{array}{c} p \\ \hline \therefore q \end{array}}$$

**Corresponding Tautology:**  
 $(p \wedge (p \rightarrow q)) \rightarrow q$

## Example:

Let  $p$  be “It is snowing.”

Let  $q$  be “I will study discrete math.”

“If it is snowing, then I will study discrete math.”  
“It is snowing.”

“Therefore , I will study discrete math.”

# Modus Ponens

- State which rule of inference is the basis of the following argument:
- “If it snows today, then we will go skiing”. “It is snowing today”, “therefore we will go skiing”.
- Let p be the proposition “It is snowing today” and q the proposition “We will go skiing” Then this argument is of the form
  - p  $\square$  q
  - p
  - $\therefore$  q
- This is an argument that uses the Modus Ponens rule.

# **Modus Tollendo Tollens**

$$\frac{P \rightarrow Q, \neg Q}{\therefore \neg P}$$

- Latin for "the way that denies by denying"
- **Denying the consequent**
- ***Abbreviated to modus tollens***
- in every instance in which
- $p \rightarrow q$  is true and  $q$  is false,
- $p$  must also be false.

p	q	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

# Modus Tollens

$$\begin{array}{c} p \rightarrow q \\ \hline \neg q \\ \hline \therefore \neg p \end{array}$$

**Corresponding Tautology:**  
 $(\neg p \wedge (p \rightarrow q)) \rightarrow \neg q$

## Example:

Let  $p$  be “it is snowing.”

Let  $q$  be “I will study discrete math.”

“If it is snowing, then I will study discrete math.”  
“I will not study discrete math.”

“Therefore , it is not snowing.”

# Hypothetical Syllogism

$$\begin{array}{c} p \rightarrow q \\ q \rightarrow r \\ \hline \therefore p \rightarrow r \end{array}$$

**Corresponding Tautology:**  
 $((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$

## Example:

Let  $p$  be “it snows.”

Let  $q$  be “I will study discrete math.”

Let  $r$  be “I will get an A.”

“If it snows, then I will study discrete math.”

“If I study discrete math, I will get an A.”

“Therefore , If it snows, I will get an A.”

# Hypothetical Syllogism

- State which rule of inference is used in the argument:
- If it rains today, then we will not have a barbecue today. If we do not have a barbecue today, then we will have a barbecue tomorrow. Therefore, if it rains today, then we will have a barbecue tomorrow.

$$p \rightarrow q$$

$$q \rightarrow r$$

$$\therefore p \rightarrow r$$

- Hence, this argument is a hypothetical syllogism.

# Disjunctive Syllogism

$$\frac{p \vee q}{\neg p} \therefore q$$

**Corresponding Tautology:**  
 $(\neg p \wedge (p \vee q)) \rightarrow q$

## Example:

Let  $p$  be “I will study discrete math.”

Let  $q$  be “I will study English literature.”

“I will study discrete math or I will study English literature.”

“I will not study discrete math.”

“Therefore , I will study English literature.”

# Addition

$$\frac{p}{\therefore p \vee q}$$

**Corresponding Tautology:**  
 $p \rightarrow(p \vee q)$

## Example:

Let  $p$  be “I will study discrete math.”

Let  $q$  be “I will visit Las Vegas.”

“I will study discrete math.”

“Therefore, I will study discrete math or I will visit Las Vegas.”

# Addition Rule

- State which rule of inference is the basis of the following argument:
- “It is below freezing now. Therefore, it is either below freezing or raining now.”
- Let p be the proposition “It is below freezing now” and q the proposition “It is raining now.” Then this argument is of the form

$$\begin{array}{c} p \\ \therefore p \vee q \end{array}$$

- This is an argument that uses the addition rule.

# Simplification

$$\frac{p \wedge q}{\therefore q}$$

**Corresponding Tautology:**  
 $(p \wedge q) \rightarrow p$

## Example:

Let  $p$  be “I will study discrete math.”

Let  $q$  be “I will study English literature.”

“I will study discrete math and English literature”

“Therefore, I will study discrete math.”

# Simplification Rule

- State which rule of inference is the basis of the following argument:
- “It is below freezing and raining now. Therefore, it is below freezing now.”
- Let p be the proposition “It is below freezing now,” and let q be the proposition “It is raining now.” This argument is of the form
  - $$\begin{array}{c} p \wedge q \\ \therefore p \end{array}$$
- This argument uses the simplification rule.

# Conjunction

$$\begin{array}{c} p \\ q \\ \hline \therefore p \wedge q \end{array}$$

**Corresponding Tautology:**  
 $((p) \wedge (q)) \rightarrow (p \wedge q)$

## Example:

Let  $p$  be “I will study discrete math.”

Let  $q$  be “I will study English literature.”

“I will study discrete math.”

“I will study English literature.”

“Therefore, I will study discrete math and I will study English literature.”

# Resolution

Resolution plays an important role in AI and is used in Prolog.

$$\frac{\neg p \vee r \\ p \vee q}{\therefore q \vee r}$$

**Corresponding Tautology:**  
 $((\neg p \vee r) \wedge (p \vee q)) \rightarrow (q \vee r)$

## Example:

Let  $p$  be “I will study discrete math.”

Let  $r$  be “I will study English literature.”

Let  $q$  be “I will study databases.”

“I will not study discrete math or I will study English literature.”

“I will study discrete math or I will study databases.”

“Therefore, I will study databases or I will English literature.”

# Examples

1. State which rule of inference is the basis of the following argument: “It is below freezing now.

Therefore, it is below freezing or raining now.”

2. State which rule of inference is the basis of the following argument: “It is below freezing and raining now.

Therefore, it is below freezing now.”



3. State which rule of inference is used in the argument:

If it rains today, then we will not have a barbecue today.

If we do not have a barbecue today, then we will have a barbecue tomorrow.

Therefore, if it rains today, then we will have a barbecue tomorrow.

4. Use resolution to show that the hypotheses “Jasmine is skiing or it is not snowing” and “It is snowing or Bart is playing hockey” imply that “Jasmine is skiing or Bart is playing hockey.”



- Show that the premises  $(p \wedge q) \vee r$  and  $r \rightarrow s$  imply the conclusion  $p \vee s$ .

# Using the Rules of Inference to Build Valid Arguments

- A *valid argument* is a sequence of statements. Each statement is either a premise or follows from previous statements by rules of inference. The last statement is called conclusion.
- A valid argument takes the following form:

$$\begin{array}{c} S_1 \\ S_2 \\ \vdots \\ S_n \\ \therefore C \end{array}$$

# Fallacies

- Arguments are based on tautologies.
- Fallacies are based on contingencies.
- The proposition  $((p \rightarrow q) \wedge q) \rightarrow p$  is not a tautology, because it is false when  $p$  is false and  $q$  is true.
- This type of incorrect reasoning is called the **fallacy of affirming the conclusion**.
- Example:
- If you do every problem in this book, then you will learn discrete mathematics. You learned discrete mathematics. Therefore, you did every problem in this book.

# Fallacy of denying the hypothesis.

- The proposition  $((p \rightarrow q) \wedge \neg p) \rightarrow \neg q$  is not a tautology, because it is false when  $p$  is false and  $q$  is true.
- Is it correct to assume that you did not learn discrete mathematics if you did not do every problem in the book, assuming that if you do every problem in this book, then you will learn discrete mathematics?
- It is possible that you learned discrete mathematics even if you did not do every problem in this book. This incorrect argument is of the form  $p \rightarrow q$  and  $\neg p$  imply  $\neg q$ , which is an example of the fallacy of denying the hypothesis.

# Check validity of the argument

- Show that the following argument is valid. If today is Tuesday ,then I have a test in mathematics or economics. Today is Tuesday and my economics professor is sick .Therefore I have a test in Mathematics.

Solution: Let T: today is Tuesday.

M: I have a test in mathematics.

E: I have a test in economics.

S: Economics professor is sick

## Using Rules of Inference

# Fallacies

- Are the following arguments correct?
  - Example 1 (Fallacy of affirming the conclusion)

### Hypothesis

- If you success, you work hard
- You work hard

$$p \rightarrow q$$

$$q$$

---

$$\therefore p$$



### Conclusion

- You success

- Example 2 (Fallacy of denying the hypothesis)

### Hypothesis

- If you success, you work hard
- You do not success

$$p \rightarrow q$$

$$\neg p$$

---

$$\therefore \neg q$$



### Conclusion

- You do not work hard

V. Tell the validity of each argument by choosing one of the following:

**MP- Modus Ponens**

**MT- Modus Tollens**

**LS- Law of Syllogism (Hypothetical)**

**LC- Law of Contrapositive**

**NVC- No Valid Conclusion**

a. \_\_\_\_\_

b. \_\_\_\_\_

c. \_\_\_\_\_

d. \_\_\_\_\_

$$p \rightarrow \neg r$$

$$s \rightarrow t$$

$$r \rightarrow \neg t$$

$$r \rightarrow p$$

$$\neg p$$

$$\neg t$$

$$\neg t \rightarrow v$$

$$p$$

$$\therefore r$$

$$\therefore \neg s$$

$$\therefore r \rightarrow v$$

$$\therefore r$$

e. \_\_\_\_\_

f. \_\_\_\_\_

g. \_\_\_\_\_

h. \_\_\_\_\_

$$\neg w \rightarrow \neg s$$

$$t \rightarrow (r \rightarrow s)$$

$$r \rightarrow q$$

$$\neg w$$

$$\neg(r \rightarrow s)$$

$$p \rightarrow \neg r$$

$$s \rightarrow q$$

$$\therefore \neg s$$

$$\therefore \neg t$$

$$\therefore r \rightarrow \neg p$$

$$\therefore r \rightarrow s$$

V. Tell the validity of each argument by choosing one of the following:

**MP- Modus Ponens**

**MT- Modus Tollens**

**NVC- No Valid Conclusion**

**LS- Law of Syllogism (Hypothetical)**

**LC- Law of Contrapositive**

a. NVC

$$\begin{array}{c} p \cancel{\rightarrow} \sim r \\ \sim p \\ \hline \therefore r \end{array}$$

b. M.T

$$\begin{array}{c} (s \rightarrow t) \quad \checkmark \\ \wedge (\sim t) \\ \hline \therefore \boxed{\sim s} \end{array}$$

c. LS (H.S)

$$\begin{array}{c} r \rightarrow \cancel{s} \\ \cancel{A} \rightarrow v \\ \hline \therefore r \rightarrow v \end{array}$$

d. NVC

$$\begin{array}{c} r \rightarrow p \\ p \\ \hline \therefore r \end{array}$$

e. M.P

$$\begin{array}{c} \sim w \rightarrow \sim s \\ \sim w \\ \hline \therefore \sim s \end{array}$$

f. M.T

$$\begin{array}{c} t \rightarrow (r \rightarrow s) \quad \checkmark \\ \sim(r \rightarrow s) \\ \hline \therefore \sim t \end{array}$$

g. L.C

$$\begin{array}{c} p \rightarrow \sim r \\ \hline \therefore r \rightarrow \sim p \end{array}$$

h. N.V.C

$$\begin{array}{c} r \rightarrow q \\ s \rightarrow q \\ \cancel{s \rightarrow q} \\ \hline \therefore \cancel{r \rightarrow s} \end{array}$$

# Comparison between Inference and Equivalence

- **Inference ( $p \rightarrow q$ )**
    - Meaning:  
If  $p$ , then  $q$
    - $p \rightarrow q$  does **not** mean  $q \rightarrow p$
    - Either inference or equivalence rules can be used
    - $p \leftrightarrow q$  implies  $p \rightarrow q$
    - $\Rightarrow$  is used in proof
  - **Equivalence ( $\leftrightarrow$ )** is a **more restrictive** relation than Inference ( $\rightarrow$ )
- **Equivalence ( $p \leftrightarrow q$ )**
    - Meaning:  
 $p$  is equal to  $q$
    - $p \leftrightarrow q$  means  $q \leftrightarrow p$
    - Only equivalence rules can be used
    - $p \leftrightarrow q$  can be proved by showing  $p \rightarrow q$  and  $q \rightarrow p$
    - $\Leftrightarrow$  is used in proof

# Using Rules of Inference

- Example 1:

- Given:

- It is not sunny this afternoon and it is colder than yesterday.
    - We will go swimming only if it is sunny
    - If we do not go swimming, then we will take a canoe trip
    - If we take a canoe trip, then we will be home by sunset
  - Can these propositions lead to the **conclusion**  
**"We will be home by sunset"** ?

- Let p: It is sunny this afternoon  
q: It is colder than yesterday  
r: We go swimming  
s: We take a canoe trip  
t: We will be home by sunset

$\neg p \wedge q$  ■ It is **not** sunny this afternoon **and** it is colder than yesterday

$r \rightarrow p$  ■ We will go swimming **only if** it is sunny

$\neg r \rightarrow s$  ■ **If** we do **not** go swimming, **then** we will take a canoe trip

$s \rightarrow t$  ■ **If** we take a canoe trip, **then** we will be home by sunset

---

$t$  ■ We will be home by sunset

# Using Rules of Inference

	Step	Reason
Hypothesis:	1. $\neg p \wedge q$	Premise
$\neg p \wedge q$	2. $\neg p$	Simplification using (1)
$r \rightarrow p$	3. $r \rightarrow p$	Premise
$\neg r \rightarrow s$	4. $\neg r$	Modus tollens using (2) and (3)
$s \rightarrow t$	5. $\neg r \rightarrow s$	Premise
	6. $s$	Modus ponens using (4) and (5)
Conclusion:	7. $s \rightarrow t$	Premise
	8. $t$	Modus ponens using (6) and (7)

**t**

Therefore, the propositions can lead to the conclusion  
We will be home by sunset

# Using Rules of Inference

- Or, another presentation method:

Hypothesis:

$$\neg p \wedge q$$

$$r \rightarrow p$$

$$\neg r \rightarrow s$$

$$s \rightarrow t$$

$$(\neg p \wedge q) \wedge (r \rightarrow p) \wedge (\neg r \rightarrow s) \wedge (s \rightarrow t)$$

$$\Rightarrow \neg p \wedge (r \rightarrow p) \wedge (\neg r \rightarrow s) \wedge (s \rightarrow t) \text{ By Simplification}$$

$$\Rightarrow \neg r \wedge (\neg r \rightarrow s) \wedge (s \rightarrow t) \text{ By Modus Tollens}$$

$$\Rightarrow s \wedge (s \rightarrow t) \text{ By Modus Ponens}$$

Conclusion:

$$t$$

$$\Rightarrow t \text{ By Modus Ponens}$$

## EXAMPLE #2

- **Given:**
  - If you send me an e-mail message, then I will finish writing the program
  - If you do not send me an e-mail message, then I will go to sleep early
  - If I go to sleep early, then I will wake up feeling refreshed
- Can these propositions lead to the **conclusion**  
**"If I do not finish writing the program, then I will wake up feeling refreshed."**

Let

- p: you send me an e-mail message
- q: I will finish writing the program
- r: I will go to sleep early
- s: I will wake up feeling refreshed

$p \rightarrow q$

- If you send me an e-mail message,  
*then* I will finish writing the program

$\neg p \rightarrow r$

- If you do **not** send me an e-mail message,  
*then* I will go to sleep early
- If I go to sleep early, *then* I will wake up  
feeling refreshed

$r \rightarrow s$

- 
- If I do **not** finish writing the program,  
*then* I will wake up feeling refreshed

Hypothesis:

$$p \rightarrow q$$

$$\neg p \rightarrow r$$

$$r \rightarrow s$$

Conclusion:

$$\neg q \rightarrow s$$

	<b>Step</b>	<b>Reason</b>
	1. $p \rightarrow q$	Premise
	2. $\neg q \rightarrow \neg p$	Contrapositive of (1)
	3. $\neg p \rightarrow r$	Premise
	4. $\neg q \rightarrow r$	Hypothetical Syllogism using (2) and (3)
	5. $r \rightarrow s$	Premise
	6. $\neg q \rightarrow s$	Hypothetical Syllogism using (4) and (5)

Therefore, the propositions can lead to the conclusion

If I do not finish writing the program,  
then I will wake up feeling refreshed

- Or, another presentation method:

Hypothesis:

$$p \rightarrow q$$

$$\neg p \rightarrow r$$

$$r \rightarrow s$$

Conclusion:


$$\neg q \rightarrow s$$

$$\underline{(p \rightarrow q) \wedge (\neg p \rightarrow r) \wedge (r \rightarrow s)}$$

$$\Leftrightarrow \underline{(\neg q \rightarrow \neg p) \wedge (\neg p \rightarrow r)} \wedge (r \rightarrow s) \quad \text{Contrapositive}$$

$$\Rightarrow \underline{(\neg q \rightarrow r) \wedge (r \rightarrow s)} \quad \text{By Hypothetical Syllogism}$$

$$\Rightarrow \underline{(\neg q \rightarrow s)} \quad \text{By Hypothetical Syllogism}$$



Rule of inference;

If it does not rain or if it is not foggy ,then sailing race will be held and the life saving demonstration will go on” ,” if the sailing race is held ,then the trophy will be awarded” and “ The trophy was not allowed” conclusion: “It rained”.

# Valid Arguments

**Example 3:** From the single proposition

$$p \wedge (p \rightarrow q)$$

Show that  $q$  is a conclusion.

**Solution:**

Step	Reason
1. $p \wedge (p \rightarrow q)$	Premise
2. $p$	Conjunction using (1)
3. $p \rightarrow q$	Conjunction using (1)
4. $q$	Modus Ponens using (2) and (3)

# Valid Arguments (Predicate Logic)

**TABLE 2** Rules of Inference for Quantified Statements.

<i>Rule of Inference</i>	<i>Name</i>
$\begin{array}{c} \forall x P(x) \\ \hline \therefore P(c) \end{array}$	Universal instantiation
$\begin{array}{c} P(c) \text{ for an arbitrary } c \\ \hline \therefore \forall x P(x) \end{array}$	Universal generalization
$\begin{array}{c} \exists x P(x) \\ \hline \therefore P(c) \text{ for some element } c \end{array}$	Existential instantiation
$\begin{array}{c} P(c) \text{ for some element } c \\ \hline \therefore \exists x P(x) \end{array}$	Existential generalization

# Universal Instantiation

- Universal instantiation is the rule of inference used to conclude that  $P(c)$  is true, where  $c$  is a particular member of the domain, given the premise  $\forall x P(x)$ .
- Universal instantiation is used when we conclude from the statement “All women are wise” that “Lisa is wise,” where Lisa is a member of the domain of all women.

# Universal Generalization

- Universal generalization is the rule of inference that states that  $\forall x P(x)$  is true, given the premise that  $P(c)$  is true for all elements  $c$  in the domain.
- Universal generalization is used when we show that  $\forall x P(x)$  is true by taking an arbitrary element  $c$  from the domain and showing that  $P(c)$  is true.
- Universal generalization is used implicitly in many proofs in mathematics and is seldom mentioned explicitly. However, the error of adding unwarranted assumptions about the arbitrary element  $c$  when universal generalization is used is all too common in incorrect reasoning.

# Existential Instantiation

- Existential instantiation is the rule that allows us to conclude that there is an element  $c$  in the domain for which  $P(c)$  is true if we know that  $\exists x P(x)$  is true.
- We cannot select an arbitrary value of  $c$  here, but rather it must be a  $c$  for which  $P(c)$  is true.
- Usually we have no knowledge of what  $c$  is, only that it exists. Because it exists, we may give it a name ( $c$ ) and continue our argument.

# Existential Generalization

- Existential generalization is the rule of inference that is used to conclude that  $\exists x P(x)$  is true when a particular element c with  $P(c)$  true is known.
- That is, if we know one element c in the domain for which  $P(c)$  is true, then we know that  $\exists x P(x)$  is true.

# Examples Valid Arguments (Predicate Logic)

## EXAMPLE 12

Extra  
Examples ➤

Show that the premises “Everyone in this discrete mathematics class has taken a course in computer science” and “Marla is a student in this class” imply the conclusion “Marla has taken a course in computer science.”

*Solution:* Let  $D(x)$  denote “ $x$  is in this discrete mathematics class,” and let  $C(x)$  denote “ $x$  has taken a course in computer science.” Then the premises are  $\forall x(D(x) \rightarrow C(x))$  and  $D(\text{Marla})$ . The conclusion is  $C(\text{Marla})$ .

The following steps can be used to establish the conclusion from the premises.

Step	Reason
1. $\forall x(D(x) \rightarrow C(x))$	Premise
2. $D(\text{Marla}) \rightarrow C(\text{Marla})$	Universal instantiation from (1)
3. $D(\text{Marla})$	Premise
4. $C(\text{Marla})$	Modus ponens from (2) and (3)



# Examples Valid Arguments (Predicate Logic)

**EXAMPLE 13** Show that the premises “A student in this class has not read the book,” and “Everyone in this class passed the first exam” imply the conclusion “Someone who passed the first exam has not read the book.”


*Solution:* Let  $C(x)$  be “ $x$  is in this class,”  $B(x)$  be “ $x$  has read the book,” and  $P(x)$  be “ $x$  passed the first exam.” The premises are  $\exists x(C(x) \wedge \neg B(x))$  and  $\forall x(C(x) \rightarrow P(x))$ . The conclusion is  $\exists x(P(x) \wedge \neg B(x))$ . These steps can be used to establish the conclusion from the premises.

Step	Reason
1. $\exists x(C(x) \wedge \neg B(x))$	Premise
2. $C(a) \wedge \neg B(a)$	Existential instantiation from (1)
3. $C(a)$	Simplification from (2)
4. $\forall x(C(x) \rightarrow P(x))$	Premise
5. $C(a) \rightarrow P(a)$	Universal instantiation from (4)
6. $P(a)$	Modus ponens from (3) and (5)
7. $\neg B(a)$	Simplification from (2)
8. $P(a) \wedge \neg B(a)$	Conjunction from (6) and (7)
9. $\exists x(P(x) \wedge \neg B(x))$	Existential generalization from (8)



# **Basic Structures: Sets and Functions**

**Chapter 2**



**Sets**

**Section 2.1**


# Section Summary

- Definition of sets
- Describing Sets
  - Roster Method
  - Set-Builder Notation
- Some Important Sets in Mathematics
- Empty Set and Universal Set
- Subsets and Set Equality
- Cardinality of Sets
- Tuples
- Cartesian Product

# Introduction

- Sets are one of the basic building blocks for the types of objects considered in discrete mathematics.
  - Important for counting.
  - Programming languages have set operations.
- Set theory is an important branch of mathematics.
  - Many different systems of axioms have been used to develop set theory.
  - Here we are not concerned with a formal set of axioms for set theory. Instead, we will use what is called naïve set theory.

# Sets



- A *set* is an unordered collection of objects.
  - the students in this class
  - the chairs in this room
- The objects in a set are called the *elements*, or *members* of the set. A set is said to *contain* its elements.
- The notation  $a \in A$  denotes that  $a$  is an element of the set  $A$ .
- If  $a$  is not a member of  $A$ , write  $a \notin A$

# Describing a Set: Roster Method

- $S = \{a,b,c,d\}$
- Order not important

$$S = \{a,b,c,d\} = \{b,c,a,d\}$$

- Each distinct object is either a member or not; listing more than once does not change the set.

$$S = \{a,b,c,d\} = \{a,b,c,b,c,d\}$$

- Ellipses (...) may be used to describe a set without listing all of the members when the pattern is clear.

$$S = \{a,b,c,d, \dots, z\}$$

# Roster Method

- Set of all vowels in the English alphabet:

$$V = \{a, e, i, o, u\}$$

- Set of all odd positive integers less than 10:

$$O = \{1, 3, 5, 7, 9\}$$

- Set of all positive integers less than 100:

$$S = \{1, 2, 3, \dots, 99\}$$

- Set of all integers less than 0:

$$S = \{\dots, -3, -2, -1\}$$

# Some Important Sets

**N** = *natural numbers* =  $\{0, 1, 2, 3, \dots\}$

**Z** = *integers* =  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$

**Z<sup>+</sup>** = *positive integers* =  $\{1, 2, 3, \dots\}$

**R** = set of *real numbers*

**R<sup>+</sup>** = set of *positive real numbers*

**C** = set of *complex numbers*.

**Q** = set of rational numbers

# Set-Builder Notation

- Specify the property or properties that all members must satisfy:

$$S = \{x \mid x \text{ is a positive integer less than } 100\}$$

$$O = \{x \mid x \text{ is an odd positive integer less than } 10\}$$

$$O = \{x \in \mathbf{Z}^+ \mid x \text{ is odd and } x < 10\}$$

- A predicate may be used:

$$S = \{x \mid P(x)\}$$

- Example:  $S = \{x \mid \text{Prime}(x)\}$



- Positive rational numbers:

$$\mathbf{Q}^+ = \{x \in \mathbf{R} \mid x = p/q, \text{ for some positive integers } p, q\}$$

# Universal Set and Empty Set

- The *universal set*  $U$  is the set containing everything currently under consideration.
  - Sometimes implicit
  - Sometimes explicitly stated.
  - Contents depend on the context.
- The empty set is the set with no elements. Symbolized  $\emptyset$ , but  $\{\}$  also used.

Venn Diagram



John Venn (1834-1923)  
Cambridge, UK

# Some things to remember

- Sets can be elements of sets.

$$\{\{1,2,3\}, a, \{b,c\}\}$$

$$\{\mathbf{N}, \mathbf{Z}, \mathbf{Q}, \mathbf{R}\}$$

- The empty set is different from a set containing the empty set.

$$\emptyset \neq \{ \emptyset \}$$

# Set Equality


**Definition:** Two sets are *equal* if and only if they have the same elements.

- Therefore if  $A$  and  $B$  are sets, then  $A$  and  $B$  are equal if and only if  $\forall x(x \in A \leftrightarrow x \in B)$ .
- We write  $A = B$  if  $A$  and  $B$  are equal sets.

$$\{1,3,5\} = \{3, 5, 1\}$$

$$\{1,5,5,5,3,3,1\} = \{1,3,5\}$$

# Subsets



**Definition:** The set  $A$  is a *subset* of  $B$ , if and only if every element of  $A$  is also an element of  $B$ .

- The notation  $A \subseteq B$  is used to indicate that  $A$  is a subset of the set  $B$ .
  - $A \subseteq B$  holds if and only if  $\forall x(x \in A \rightarrow x \in B)$
1. Because  $a \in \emptyset$  is always false,  $\emptyset \subseteq S$ , for every set  $S$ .
  2. Because  $a \in S \rightarrow a \in S$ ,  $S \subseteq S$ , for every set  $S$ .

# Showing a Set is or is not a Subset of Another Set

- **Showing that A is a Subset of B:** To show that  $A \subseteq B$ , show that if  $x$  belongs to  $A$ , then  $x$  also belongs to  $B$ .
- **Showing that A is not a Subset of B:** To show that  $A$  is not a subset of  $B$ ,  $A \not\subseteq B$ , find an element  $x \in A$  with  $x \notin B$ . (Such an  $x$  is a counterexample to the claim that  $x \in A$  implies  $x \in B$ .)

## Examples:

1. The set of all computer science majors at your school is a subset of all students at your school.
2. The set of integers with squares less than 100 is not a subset of the set of nonnegative integers.

# Another look at Equality of Sets

- Recall that two sets  $A$  and  $B$  are *equal*, denoted by  $A = B$ , iff
$$\forall x(x \in A \leftrightarrow x \in B)$$
- Using logical equivalences we have that  $A = B$  iff
$$\forall x[(x \in A \rightarrow x \in B) \wedge (x \in B \rightarrow x \in A)]$$
- This is equivalent to

$$A \subseteq B \quad \text{and} \quad B \subseteq A$$


# Proper Subsets

**Definition:** If  $A \subseteq B$ , but  $A \neq B$ , then we say  $A$  is a *proper subset* of  $B$ , denoted by  $A \subset B$ . If  $A \subset B$ , then

$$\forall x(x \in A \rightarrow x \in B) \wedge \exists x(x \in B \wedge x \notin A)$$

is true.

Venn Diagram



# Set Cardinality

**Definition:** If there are exactly  $n$  distinct elements in  $S$  where  $n$  is a nonnegative integer, we say that  $S$  is *finite*. Otherwise it is *infinite*.

**Definition:** The *cardinality* of a finite set  $A$ , denoted by  $|A|$ , is the number of (distinct) elements of  $A$ .

**Examples:**

1.  $|\emptyset| = 0$
2. Let  $S$  be the letters of the English alphabet. Then  $|S| = 26$
3.  $|\{1,2,3\}| = 3$
4.  $|\{\emptyset\}| = 1$
5. The set of integers is infinite.

# Power Sets

**Definition:** The set of all subsets of a set  $A$ , denoted  $P(A)$ , is called the *power set* of  $A$ .

**Example:** If  $A = \{a,b\}$  then

$$P(A) = \{\emptyset, \{a\}, \{b\}, \{a,b\}\}$$

- If a set has  $n$  elements, then the cardinality of the power set is  $2^n$ . (In Chapters 5 and 6, we will discuss different ways to show this.)

# Tuples

- The *ordered n-tuple*  $(a_1, a_2, \dots, a_n)$  is the ordered collection that has  $a_1$  as its first element and  $a_2$  as its second element and so on until  $a_n$  as its last element.
- Two n-tuples are equal if and only if their corresponding elements are equal.
- 2-tuples are called *ordered pairs*.
- The ordered pairs  $(a, b)$  and  $(c, d)$  are equal if and only if  $a = c$  and  $b = d$ .

# Cartesian Product



René Descartes  
(1596-1650)

**Definition:** The *Cartesian Product* of two sets  $A$  and  $B$ , denoted by  $A \times B$  is the set of ordered pairs  $(a,b)$  where  $a \in A$  and  $b \in B$ .

$$A \times B = \{(a, b) | a \in A \wedge b \in B\}$$

**Example:**

$$A = \{a, b\} \quad B = \{1, 2, 3\}$$

$$A \times B = \{(a, 1), (a, 2), (a, 3), (b, 1), (b, 2), (b, 3)\}$$

- **Definition:** A subset  $R$  of the Cartesian product  $A \times B$  is called a *relation* from the set  $A$  to the set  $B$ .  
(Relations will be covered in depth in Chapter 9. )

# Cartesian Product

**Definition:** The cartesian products of the sets  $A_1, A_2, \dots, A_n$ , denoted by  $A_1 \times A_2 \times \dots \times A_n$ , is the set of ordered  $n$ -tuples  $(a_1, a_2, \dots, a_n)$  where  $a_i$  belongs to  $A_i$  for  $i = 1, \dots, n$ .

$$A_1 \times A_2 \times \dots \times A_n = \{(a_1, a_2, \dots, a_n) | a_i \in A_i \text{ for } i = 1, 2, \dots, n\}$$

**Example:** What is  $A \times B \times C$  where  $A = \{0, 1\}$ ,  $B = \{1, 2\}$  and  $C = \{0, 1, 2\}$

**Solution:**  $A \times B \times C = \{(0, 1, 0), (0, 1, 1), (0, 1, 2), (0, 2, 0), (0, 2, 1), (0, 2, 2), (1, 1, 0), (1, 1, 1), (1, 1, 2), (1, 2, 0), (1, 2, 1), (1, 2, 2)\}$

# Truth Sets of Quantifiers

- Given a predicate  $P$  and a domain  $D$ , we define the *truth set* of  $P$  to be the set of elements in  $D$  for which  $P(x)$  is true. The truth set of  $P(x)$  is denoted by
$$\{x \in D | P(x)\}$$
- Example:** The truth set of  $P(x)$  where the domain is the integers and  $P(x)$  is “ $|x| = 1$ ” is the set  $\{-1, 1\}$

# **Set Operations**

**Section 2.2**


# Section Summary

- Set Operations
  - Union
  - Intersection
  - Complementation
  - Difference
- More on Set Cardinality
- Set Identities
- Proving Identities
- Membership Tables

# Boolean Algebra

- Propositional calculus and set theory are both instances of an algebraic system called a *Boolean Algebra*. This is discussed in Chapter 12.
- The operators in set theory are analogous to the corresponding operator in propositional calculus.
- As always there must be a universal set  $U$ . All sets are assumed to be subsets of  $U$ .

# Union




- **Definition:** Let  $A$  and  $B$  be sets. The *union* of the sets  $A$  and  $B$ , denoted by  $A \cup B$ , is the set:

$$\{x | x \in A \vee x \in B\}$$

- **Example:** What is  $\{1,2,3\} \cup \{3, 4, 5\}$ ?

**Solution:**  $\{1,2,3,4,5\}$

Venn Diagram for  $A \cup B$



# Intersection

- **Definition:** The *intersection* of sets  $A$  and  $B$ , denoted by  $A \cap B$ , is

$$\{x | x \in A \wedge x \in B\}$$

- Note if the intersection is empty, then  $A$  and  $B$  are said to be *disjoint*.

- **Example:** What is?  $\{1,2,3\} \cap \{3,4,5\}$  ?


**Solution:**  $\{3\}$

- **Example:** What is?

$$\{1,2,3\} \cap \{4,5,6\} ?$$

**Solution:**  $\emptyset$

Venn Diagram for  $A \cap B$



# Complement

**Definition:** If  $A$  is a set, then the complement of the  $A$  (with respect to  $U$ ), denoted by  $\bar{A}$  is the set  $U - A$


$$\bar{A} = \{x \in U \mid x \notin A\}$$

(The complement of  $A$  is sometimes denoted by  $A^c$ .)

**Example:** If  $U$  is the positive integers less than 100, what is the complement of  $\{x \mid x > 70\}$

Solution:  $\{x \mid x \leq 70\}$

Venn Diagram for Complement




# Difference

- **Definition:** Let  $A$  and  $B$  be sets. The *difference* of  $A$  and  $B$ , denoted by  $A - B$ , is the set containing the elements of  $A$  that are not in  $B$ . The difference of  $A$  and  $B$  is also called the complement of  $B$  with respect to  $A$ .

$$A - B = \{x \mid x \in A \wedge x \notin B\} = A \cap \complement B$$


Venn Diagram for  $A - B$



# The Cardinality of the Union of Two Sets

Inclusion-Exclusion

$$|A \cup B| = |A| + |B| - |A \cap B|$$



Venn Diagram for  $A, B, A \cap B, A \cup B$

**Example:** Let  $A$  be the math majors in your class and  $B$  be the CS majors. To count the number of students who are either math majors or CS majors, add the number of math majors and the number of CS majors, and subtract the number of joint CS/math majors.

We will return to this principle in Chapter 6 and Chapter 8 where we will derive a formula for the cardinality of the union of  $n$  sets, where  $n$  is a positive integer.

# Review Questions

**Example:**  $U = \{0,1,2,3,4,5,6,7,8,9,10\}$   $A = \{1,2,3,4,5\}$ ,  $B = \{4,5,6,7,8\}$

1.  $A \cup B$

**Solution:**  $\{1,2,3,4,5,6,7,8\}$

2.  $A \cap B$

**Solution:**  $\{4,5\}$

3.  $\bar{A}$

**Solution:**  $\{0,6,7,8,9,10\}$

4.

**Solution:**  $\{0,1,2,3,9,10\}$

5.  $A - B$

**Solution:**  $\{1,2,3\}$

6.  $B - A$

**Solution:**  $\{6,7,8\}$

# Symmetric Difference (*optional*)

**Definition:** The *symmetric difference* of A and B, denoted by  $A \oplus B$  is the set


$$\text{Example: } (A - B) \cup (B - A)$$

$$U = \{0,1,2,3,4,5,6,7,8,9,10\}$$

$$A = \{1,2,3,4,5\} \quad B = \{4,5,6,7,8\}$$

What is:

- **Solution:**  $\{1,2,3,6,7,8\}$



Venn Diagram

# Set Identities

- Identity laws

$$A \cup \emptyset = A \quad A \cap U = A$$

- Domination laws

$$A \cup U = U \quad A \cap \emptyset = \emptyset$$

- Idempotent laws

$$A \cup A = A \quad A \cap A = A$$

- Complementation law

$$\overline{(\overline{A})} = A$$

*Continued on next slide* □

# Set Identities

- Commutative laws

$$A \cup B = B \cup A$$

$$A \cap B = B \cap A$$

- Associative laws

$$A \cup (B \cup C) = (A \cup B) \cup C$$

$$A \cap (B \cap C) = (A \cap B) \cap C$$

- Distributive laws

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

*Continued on next slide* □

# Set Identities

- De Morgan's laws

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

- Absorption laws

$$A \cup (A \cap B) = A$$

$$A \cap (A \cup B) = A$$

- Complement laws

$$A \cup \overline{A} = U$$

$$A \cap \overline{A} = \emptyset$$

# Proving Set Identities

- Different ways to prove set identities:
  1. Prove that each set (side of the identity) is a subset of the other.
  2. Use set builder notation and propositional logic.
  3. Membership Tables: Verify that elements in the same combination of sets always either belong or do not belong to the same side of the identity. Use 1 to indicate it is in the set and a 0 to indicate that it is not.

# Proof of Second De Morgan Law

**Example:** Prove that  $\overline{A \cap B} = \overline{A} \cup \overline{B}$

**Solution:** We prove this identity by showing that:

$$1) \overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$$

and

$$2) \overline{A} \cup \overline{B} \subseteq \overline{A \cap B}$$

*Continued on next slide* □

# Proof of Second De Morgan Law

These steps show that:

$$\overline{A \cap B} \subseteq \overline{A} \cup \overline{B}$$

$$x \in \overline{A \cap B}$$

by assumption

$$x \notin A \cap B$$

defn. of complement

$$\neg((x \in A) \wedge (x \in B))$$

defn. of intersection

$$\neg(x \in A) \vee \neg(x \in B)$$

1st De Morgan Law for Prop Logic

$$x \notin A \vee x \notin B$$

defn. of negation

$$x \in \overline{A} \vee x \in \overline{B}$$

defn. of complement

$$x \in \overline{A} \cup \overline{B}$$

defn. of union

*Continued on next slide* □

# Proof of Second De Morgan Law

These steps show that:

$$\overline{A \cup B} \subseteq \overline{A \cap B}$$

$$x \in \overline{A \cup B}$$

by assumption

$$(x \in \overline{A}) \vee (x \in \overline{B})$$

defn. of union

$$(x \notin A) \vee (x \notin B)$$

defn. of complement

$$\neg(x \in A) \vee \neg(x \in B)$$

defn. of negation

$$\neg((x \in A) \wedge (x \in B))$$

by 1st De Morgan Law for Prop Logic

$$\neg(x \in A \cap B)$$

defn. of intersection

$$x \in \overline{A \cap B}$$

defn. of complement



# Set-Builder Notation: Second De Morgan Law

$$\begin{aligned}\overline{A \cap B} &= \{x | x \notin A \cap B\} && \text{by defn. of complement} \\ &= \{x | \neg(x \in (A \cap B))\} && \text{by defn. of does not belong symbol} \\ &= \{x | \neg(x \in A \wedge x \in B)\} && \text{by defn. of intersection} \\ &= \{x | \neg(x \in A) \vee \neg(x \in B)\} && \text{by 1st De Morgan law} \\ &&& \text{for Prop Logic} \\ &= \{x | x \notin A \vee x \notin B\} && \text{by defn. of not belong symbol} \\ &= \{x | x \in \overline{A} \vee x \in \overline{B}\} && \text{by defn. of complement} \\ &= \{x | x \in \overline{A} \cup \overline{B}\} && \text{by defn. of union} \\ &= \overline{A} \cup \overline{B} && \text{by meaning of notation}\end{aligned}$$



# Membership Table

**Example:** Construct a membership table to show that the distributive law holds.

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

**Solution:**

A	B	C	$B \cap C$	$A \cup (B \cap C)$	$A \cup B$	$A \cup C$	$(A \cup B) \cap (A \cup C)$
1	1	1	1	1	1	1	1
1	1	0	0	1	1	1	1
1	0	1	0	1	1	1	1
1	0	0	0	1	1	1	1
0	1	1	1	1	1	1	1
0	1	0	0	0	1	0	0
0	0	1	0	0	0	1	0
0	0	0	0	0	0	0	0

# Functions

Section 2.3


# Section Summary

- Definition of a Function.
  - Domain, Codomain
  - Image, Pre image
- Injection, Surjection, Bijection
- Inverse Function
- Function Composition
- Graphing Functions
- Floor, Ceiling, Factorial

# Functions

**Definition:** Let  $A$  and  $B$  be nonempty sets. A *function*  $f$  from  $A$  to  $B$ , denoted  $f: A \rightarrow B$  is an assignment of each element of  $A$  to exactly one element of  $B$ . We write  $f(a) = b$  if  $b$  is the unique element of  $B$  assigned by the function  $f$  to the element  $a$  of  $A$ .

- Functions are sometimes called *mappings* or *transformations*.



# Functions

- A function  $f: A \rightarrow B$  can also be defined as a subset of  $A \times B$  (a relation). This subset is restricted to be a relation where no two elements of the relation have the same first element.
- Specifically, a function  $f$  from  $A$  to  $B$  contains one, and only one ordered pair  $(a, b)$  for every element  $a \in A$ .

$$\forall x[x \in A \rightarrow \exists y[y \in B \wedge (x, y) \in f]]$$


and

$$\forall x, y_1, y_2[((x, y_1) \in f \wedge (x, y_2)) \rightarrow y_1 = y_2]$$

# Functions

Given a function  $f: A \rightarrow B$ :

- We say  $f$  maps  $A$  to  $B$  or  $f$  is a *mapping* from  $A$  to  $B$ .
- $A$  is called the *domain* of  $f$ .
- $B$  is called the *codomain* of  $f$ .
- If  $f(a) = b$ ,
  - then  $b$  is called the *image* of  $a$  under  $f$ .
  - $a$  is called the *preimage* of  $b$ .
- The range of  $f$  is the set of all images of points in  $A$  under  $f$ . We denote it by  $f(A)$ .
- Two functions are *equal* when they have the same domain, the same codomain and map each element of the domain to the same element of the codomain.



# Equal Functions


- Two functions are **equal** when they
  - have the same domain,
  - have the same codomain,
  - map each element of their common domain to the same element in their common codomain.
- If we change either the domain or the codomain of a function, then we obtain a different function.
- If we change the mapping of elements, then we also obtain a different function.

# Representing Functions

- Functions may be specified in different ways:
  - An explicit statement of the assignment.  
Students and grades example.
  - A formula.  
 $f(x) = x + 1$
  - A computer program.
    - A Java program that when given an integer  $n$ , produces the  $n$ th Fibonacci Number (covered in the next section and also in Chapter 5).

# Activity Time

What are the domain, codomain, and range of the following function



**FIGURE 1** Assignment of Grades in a Discrete Mathematics Class.

# Solution

- Let  $G$  be the function that assigns a grade to a student in our discrete mathematics class.
- Note that  $G(\text{Adams}) = A$ , for instance.
- The domain of  $G$  is the set  $\{\text{Adams}, \text{Chou}, \text{Goodfriend}, \text{Rodriguez}, \text{Stevens}\}$ ,
- The codomain is the set  $\{A, B, C, D, F\}$ .
- The range of  $G$  is the set  $\{A, B, C, F\}$ ,

# Questions

$f(a) = ? \quad z$

The image of d is ?  $\quad z$


The domain of f is ?  $\quad A$

The codomain of f is ?  $\quad B$

The preimage of y is ?  $\quad b$

$f(A) = ?$

The preimage(s) of z is (are) ?  $\quad \{a,c,d\}$




# Question on Functions and Sets

- If  $f : A \rightarrow B$  and  $S$  is a subset of  $A$ , then

$$f(S) = \{f(s) | s \in S\}$$



$f \{a,b,c,\}$  is ?     $\{y,z\}$

$f \{c,d\}$  is ?     $\{z\}$




# Injections

**Definition:** A function  $f$  is said to be *one-to-one* , or *injective*, if and only if  $f(a) = f(b)$  implies that  $a = b$  for all  $a$  and  $b$  in the domain of  $f$ . A function is said to be an *injection* if it is one-to-one.(every element of codomain is the image of at most one element of its domain).




# Surjections

**Definition:** A function  $f$  from  $A$  to  $B$  is called *onto* or *surjective*, if and only if for every element  $b \in B$  there is an element  $a \in A$  with  $f(a) = b$ . A function  $f$  is called a *surjection* if it is onto.(every element of  $B$  is mapped by atleast one element of  $A$ ).







# Bijections

**Definition:** A function  $f$  is a *one-to-one correspondence*, or a *bijection*, if it is both one-to-one and onto (surjective and injective).



# Question





**FIGURE 5** Examples of Different Types of Correspondences.

# real-valued

- A function is called **real-valued** if its codomain is the set of real numbers.
- A function is called **integer-valued** if its codomain is the set of integer numbers.
- Let  $f_1$  and  $f_2$  be functions from  $A$  to  $\mathbf{R}$ .
- Then  $f_1 + f_2$  and  $f_1f_2$  are also functions from  $A$  to  $\mathbf{R}$  defined for all  $x \in A$  by
  - $(f_1 + f_2)(x) = f_1(x) + f_2(x),$
  - $(f_1f_2)(x) = f_1(x)f_2(x).$

**EXAMPLE 6** Let  $f_1$  and  $f_2$  be functions from  $\mathbf{R}$  to  $\mathbf{R}$  such that  $f_1(x) = x^2$  and  $f_2(x) = x - x^2$ . What are the functions  $f_1 + f_2$  and  $f_1 f_2$ ?

*Solution:* From the definition of the sum and product of functions, it follows that

$$(f_1 + f_2)(x) = f_1(x) + f_2(x) = x^2 + (x - x^2) = x$$

and

$$(f_1 f_2)(x) = x^2(x - x^2) = x^3 - x^4.$$

# Showing that $f$ is one-to-one or onto

Suppose that  $f : A \rightarrow B$ .

*To show that  $f$  is injective* Show that if  $f(x) = f(y)$  for arbitrary  $x, y \in A$  with  $x \neq y$ , then  $x = y$ .

*To show that  $f$  is not injective* Find particular elements  $x, y \in A$  such that  $x \neq y$  and  $f(x) = f(y)$ .

*To show that  $f$  is surjective* Consider an arbitrary element  $y \in B$  and find an element  $x \in A$  such that  $f(x) = y$ .

*To show that  $f$  is not surjective* Find a particular  $y \in B$  such that  $f(x) \neq y$  for all  $x \in A$ .

# Showing that $f$ is one-to-one or onto

**Example 1:** Let  $f$  be the function from  $\{a,b,c,d\}$  to  $\{1,2,3\}$  defined by  $f(a) = 3$ ,  $f(b) = 2$ ,  $f(c) = 1$ , and  $f(d) = 3$ . Is  $f$  an onto function?

# Showing that $f$ is one-to-one or onto

**Example 1:** Let  $f$  be the function from  $\{a,b,c,d\}$  to  $\{1,2,3\}$  defined by  $f(a) = 3$ ,  $f(b) = 2$ ,  $f(c) = 1$ , and  $f(d) = 3$ . Is  $f$  an onto function?

**Solution:** Yes,  $f$  is onto since all three elements of the codomain are images of elements in the domain. If the codomain were changed to  $\{1,2,3,4\}$ ,  $f$  would not be onto.

## Example 2:

Determine whether the function  $f(x) = x^2$  from the set of integers to the set of integers is one-to-one.

The function  $f(x) = x^2$  is not one-to-one because, for instance,  $f(1) = f(-1) = 1$ , but  $1 \neq -1$ .

# The *identity function*

- Let  $A$  be a set. The *identity function* on  $A$  is the function  $\iota_A : A \rightarrow A$ , where


$$\iota_A(x) = x$$

- for all  $x \in A$ .
- The function  $\iota_A$  is one-to-one and onto, so it is a bijection.



# Inverse Functions

**Definition:** Let  $f$  be a bijection from  $A$  to  $B$ . Then the *inverse* of  $f$ , denoted  $f^{-1}$  is the function from  $B$  to  $A$  defined as  $f^{-1}(y) = x$  iff  $f(x) = y$

No inverse exists unless  $f$  is a bijection. Why?



# Inverse Functions



# Questions

**Example 1:** Let  $f$  be the function from  $\{a,b,c\}$  to  $\{1,2,3\}$  such that  $f(a) = 2$ ,  $f(b) = 3$ , and  $f(c) = 1$ . Is  $f$  invertible and if so what is its inverse?

**Solution:** The function  $f$  is invertible because it is a one-to-one correspondence. The inverse function  $f^1$  reverses the correspondence given by  $f$ , so  $f^1(1) = c$ ,  $f^1(2) = a$ , and  $f^1(3) = b$ .

# Questions

**Example 2:** Let  $f: \mathbf{Z} \rightarrow \mathbf{Z}$  be such that  $f(x) = x + 1$ . Is  $f$  invertible, and if so, what is its inverse?

**Solution:** The function  $f$  is invertible because it is a one-to-one correspondence. The inverse function  $f^{-1}$  reverses the correspondence so  $f^{-1}(y) = y - 1$ .

# Questions


**Example 3:** Let  $f: \mathbf{R} \rightarrow \mathbf{R}$  be such that  $f(x) = x^2$  Is  $f$  invertible, and if so, what is its inverse?

**Solution:** The function  $f$  is not invertible because it is not one-to-one .



# Composition

- **Definition:** Let  $f: B \rightarrow C$ ,  $g: A \rightarrow B$ . The *composition of f with g*, denoted  $f \circ g$ , is the function from  $A$  to  $C$  defined by

$$f \circ g(x) = f(g(x))$$



# Composition



# Composition

**Example 1:** If  $f(x) = x^2$  and  $g(x) = 2x + 1$  ,  
then

$$f(g(x)) = (2x + 1)^2$$

and

$$g(f(x)) = 2x^2 + 1$$

# Composition Questions

**Example 2:** Let  $g$  be the function from the set  $\{a,b,c\}$  to itself such that  $g(a) = b$ ,  $g(b) = c$ , and  $g(c) = a$ . Let  $f$  be the function from the set  $\{a,b,c\}$  to the set  $\{1,2,3\}$  such that  $f(a) = 3$ ,  $f(b) = 2$ , and  $f(c) = 1$ .

What is the composition of  $f$  and  $g$ , and what is the composition of  $g$  and  $f$ .

**Solution:** The composition  $f \circ g$  is defined by

$$f \circ g (a) = f(g(a)) = f(b) = 2.$$

$$f \circ g (b) = f(g(b)) = f(c) = 1.$$

$$f \circ g (c) = f(g(c)) = f(a) = 3.$$

Note that  $g \circ f$  is not defined, because the range of  $f$  is not a subset of the domain of  $g$ .

# Composition Questions

**Example 2:** Let  $f$  and  $g$  be functions from the set of integers to the set of integers defined by  $f(x) = 2x + 3$  and  $g(x) = 3x + 2$ .

What is the composition of  $f$  and  $g$ , and also the composition of  $g$  and  $f$ ?


**Solution:**

$$f \circ g (x) = f(g(x)) = f(3x + 2) = 2(3x + 2) + 3 = 6x + 7$$


$$g \circ f (x) = g(f(x)) = g(2x + 3) = 3(2x + 3) + 2 = 6x + 11$$

# Graphs of Functions

- Let  $f$  be a function from the set  $A$  to the set  $B$ . The *graph* of the function  $f$  is the set of ordered pairs  $\{(a,b) \mid a \in A \text{ and } f(a) = b\}$ .



Graph of  $f(n) = 2n + 1$   
from  $Z$  to  $Z$



Graph of  $f(x) = x^2$   
from  $Z$  to  $Z$

# Some Important Functions

- The *floor* function, denoted

$$f(x) = \lfloor x \rfloor$$

is the largest integer less than or equal to  $x$ .

- The *ceiling* function, denoted



$$f(x) = \lceil x \rceil$$

is the smallest integer greater than or equal to  $x$

**Example:**  $\lceil 3.5 \rceil = 4$        $\lfloor 3.5 \rfloor = 3$

$$\lceil -1.5 \rceil = -1 \quad \lfloor -1.5 \rfloor = -2$$

# Floor and Ceiling Functions



Graph of (a) Floor and (b) Ceiling Functions

# Floor and Ceiling Functions

**TABLE 1** Useful Properties of the Floor and Ceiling Functions.

( $n$  is an integer,  $x$  is a real number)

(1a)  $\lfloor x \rfloor = n$  if and only if  $n \leq x < n + 1$

(1b)  $\lceil x \rceil = n$  if and only if  $n - 1 < x \leq n$

(1c)  $\lfloor x \rfloor = n$  if and only if  $x - 1 < n \leq x$

(1d)  $\lceil x \rceil = n$  if and only if  $x \leq n < x + 1$

(2)  $x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$

(3a)  $\lfloor -x \rfloor = -\lceil x \rceil$

(3b)  $\lceil -x \rceil = -\lfloor x \rfloor$

(4a)  $\lfloor x + n \rfloor = \lfloor x \rfloor + n$

(4b)  $\lceil x + n \rceil = \lceil x \rceil + n$

# Proving Properties of Functions

**Example:** Prove that  $x$  is a real number, then

$$\lfloor 2x \rfloor = \lfloor x \rfloor + \lfloor x + 1/2 \rfloor$$

**Solution:** Let  $x = n + \varepsilon$ , where  $n$  is an integer and  $0 \leq \varepsilon < 1$ .

*Case 1:*  $\varepsilon < \frac{1}{2}$

- $2x = 2n + 2\varepsilon$  and  $\lfloor 2x \rfloor = 2n$ , since  $0 \leq 2\varepsilon < 1$ .
- $\lfloor x + 1/2 \rfloor = n$ , since  $x + \frac{1}{2} = n + (1/2 + \varepsilon)$  and  $0 \leq \frac{1}{2} + \varepsilon < 1$ .
- Hence,  $\lfloor 2x \rfloor = 2n$  and  $\lfloor x \rfloor + \lfloor x + 1/2 \rfloor = n + n = 2n$ .

*Case 2:*  $\varepsilon \geq \frac{1}{2}$

- $2x = 2n + 2\varepsilon = (2n + 1) + (2\varepsilon - 1)$  and  $\lfloor 2x \rfloor = 2n + 1$ , since  $0 \leq 2\varepsilon - 1 < 1$ .
- $\lfloor x + 1/2 \rfloor = \lfloor n + (1/2 + \varepsilon) \rfloor = \lfloor n + 1 + (\varepsilon - 1/2) \rfloor = n + 1$  since  $0 \leq \varepsilon - 1/2 < 1$ .
- Hence,  $\lfloor 2x \rfloor = 2n + 1$  and  $\lfloor x \rfloor + \lfloor x + 1/2 \rfloor = n + (n + 1) = 2n + 1$ .



# Factorial Function

**Definition:**  $f: \mathbf{N} \rightarrow \mathbf{Z}^+$ , denoted by  $f(n) = n!$  is the product of the first  $n$  positive integers when  $n$  is a nonnegative integer.

$$f(n) = 1 \cdot 2 \cdots (n-1) \cdot n, \quad f(0) = 0! = 1$$

**Examples:**

$$f(1) = 1! = 1$$

$$f(2) = 2! = 1 \cdot 2 = 2$$

$$f(6) = 6! = 1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 = 720$$

$$f(20) = 2,432,902,008,176,640,000.$$

Stirling's Formula:

$$n! \sim \sqrt{2\pi n} (n/e)^n$$

$$f(n) \sim g(n) \doteq \lim_{n \rightarrow \infty} f(n)/g(n) = 1$$

# *Increasing/ decreasing functions*

- A function  $f$  whose domain and codomain are subsets of the set of real numbers is called *increasing* if  $f(x) \leq f(y)$ , and *strictly increasing* if  $f(x) < f(y)$ , whenever  $x < y$  and  $x$  and  $y$  are in the domain of  $f$ .
- Similarly,  $f$  is called *decreasing* if  $f(x) \geq f(y)$ , and *strictly decreasing* if  $f(x) > f(y)$ , whenever  $x < y$  and  $x$  and  $y$  are in the domain of  $f$ .

# *Increasing/ decreasing functions*

- A function  $f$  is
  - increasing if  $\forall x \forall y (x < y \rightarrow f(x) \leq f(y))$ ,
  - strictly increasing if  $\forall x \forall y (x < y \rightarrow f(x) < f(y))$ ,
  - decreasing if  $\forall x \forall y (x < y \rightarrow f(x) \geq f(y))$ ,
  - strictly decreasing if  $\forall x \forall y (x < y \rightarrow f(x) > f(y))$ ,

where the universe of discourse is the domain of  $f$ .

# Relationship with one-to-one

- a function that is either strictly increasing or strictly decreasing must be one-to-one.
- a function that is increasing, but not strictly increasing, or decreasing, but not strictly decreasing, is not one-to-one.