# Feynman diagrams

#### Alexander Huss

July 21, 2025

#### **Contents**

1	Introduction	1
2	Drell-Yan	1
3	Gluon scattering	2

### 1 Introduction

Let's have some fun with Feynman diagrams; we'll use the tool qgraf for their generation. If you want to reproduce the results in this document, you can get the source from here: qgraf (note that you'll need a Fortran compiler for this).

## 2 Drell-Yan

Let's start simple and see how we can generate the diagrams for the Drell-Yan process that we already know. The example files are located in ../../tools/qgraf/dy and the program is controlled using the input file qgraf.dat. In this example, the Feynman rules are encoded in a simple model file dyqcd and the output style is set by custom.sty. We can run qgraf in that directory to generate the diagrams

#loops	v-degrees		#diagrams	
0				
	_	4^1	 0	
	3^2	-	 2	

total = 2 connected diagrams

This will generate the output inside the file qlist.out:

```
*--#[ d1:
a(1) := (-1)*
  in[-1](quark(p1))*
  in[-3](antiquark(p2))*
out[-2](l_plus(q1))*
 out[-4](1_minus(q2))*
prop[1,2](photon(-p1-p2))*
vrtx[-3,-1,1] (antiquark(p2), quark(p1), photon(-p1-p2))*
vrtx[-4,-2,2](l_plus(-q2),l_minus(-q1),photon(p1+p2));
*--#] d1:
*--#[ d2:
a(2) := (-1)*
  in[-1](quark(p1))*
  in[-3](antiquark(p2))*
 out[-2](l_plus(q1))*
 out[-4](1_minus(q2))*
prop[1,2](Z(-p1-p2))*
vrtx[-3,-1,1] (antiquark(p2),quark(p1),Z(-p1-p2))*
vrtx[-4,-2,2](l_plus(-q2),l_minus(-q1),Z(p1+p2));
*--#] d2:
* end
```

which are precisely the two s-channel diagrams we already know for this process.

## 3 Gluon scattering

We will now consider the case of  $2 \to n$  gluon scattering,  $gg \to g \dots g$ . To make this more generic, we have a template file from which we generate an input file for qgraf to automatically excract the number of diagrams qgraf has generated for us.

```
#> create an input file from the tempalte
out_string=""
sep=""
for i in $(seq 1 ${ngluons}); do
    out_string="${out_string}${sep} gluon"
    sep=","
done
sed -e "s/&out&/${out_string}/g" -e "s/&loops&/${nloop}/g" qgraf.template > qgraf.dat
#> run qgraf and look for the line
#> `total = <N> connected diagrams`
#> to extract the number of diagrams qgraf has generated
```

```
if [[ -f qlist.out ]]; then rm qlist.out; fi
qgraf | awk '$1~/^total$/&&$(NF)~/^diagrams$/{print $3}'
```

We can then fill this table for different gluon multiplicities at tree and 1-loop level:

n	# diagrams (tree)	# diagrams (1-loop)
2	4	223
3	25	2105
4	220	25120

We can appreciate the rapid increase in the number of diagrams as more legs or loops are added to the diagrams. Note that this only consitites a very naive estimate for the scaling of the complexity as each individual diagram further becomes more complicated to evalute, especially with each additional loop that involves and additional integration that must be performed.

For the tree diagrams, we can write a very simple recursive formula for the number of diagrams:

$$N_{2 \to n}^{\text{diags}} = \mathcal{N}_{n+2}(g, t) \Big|_{\substack{g=1 \\ t=1}}$$

$$\tag{1}$$

where  $\mathcal{N}_{n+2}(g,t)$  is a polynomial that "counts" the number of gluons and three-gluon vertices for each diagram and can be defined recursively:

$$\mathcal{N}_{m+1}(g,t) = \left(g^3 t \frac{\partial}{\partial g} + g \frac{\partial}{\partial t}\right) \mathcal{N}_m(g,t), \qquad \mathcal{N}_3(g,t) = g^3 t \qquad (2)$$

The first term corresponds to an insertion of a new gluon into an existing gluon line, while the second term describes the insertion of a gluon into an existing three-gluon vertex and converting it to a four-gluon vertex.

```
import sympy
g, t = sympy.symbols('g t')
n_poly = g**3 * t
n_legs = 3
for i in range(9):
    n_poly = sympy.diff(n_poly,g) * g**3 * t + sympy.diff(n_poly,t) * g
    n_legs += 1
    print("{}->{} gluon scattering has {} diagrams".format(2,n_legs-2,n_poly.subs([(g,1),(t,1)])))
```

```
2->2 gluon scattering has 4 diagrams
```

<sup>2-&</sup>gt;3 gluon scattering has 25 diagrams

<sup>2-&</sup>gt;4 gluon scattering has 220 diagrams

<sup>2-&</sup>gt;5 gluon scattering has 2485 diagrams

<sup>2-&</sup>gt;6 gluon scattering has 34300 diagrams

<sup>2-&</sup>gt;7 gluon scattering has 559405 diagrams

<sup>2-&</sup>gt;8 gluon scattering has 10525900 diagrams

<sup>2-&</sup>gt;9 gluon scattering has 224449225 diagrams

<sup>2-&</sup>gt;10 gluon scattering has 5348843500 diagrams