

PART-A

Name of the Student: Jay Amin

Signature of the Student:  Roll No.: 230123022

Signature of the Invigilator: 

INSTRUCTIONS

1. Check that you have **10 printed pages**. The PART-A of the examination has **3 questions**, for a total of **20 points**.
2. Attempt **all questions**.
3. There is **no credit** for a solution if the appropriate work is not shown, even if the answer is correct.
4. Write your answers in this booklet and only in the **space marked for answers**. Any answer written in any space other than the designated space will **not be evaluated**.
5. The pages numbered **8 to 10** are kept as **additional pages**. If you cannot able to fit a complete answer in the space provided just after the question, you can use these three pages by referring the page number of additional page in the original space.
6. At the beginning of the examination, a supplementary sheet has been provided only for rough work. Should you need more, please ask the invigilator.
7. No question requires any clarification from the instructors or invigilators, even if a question has an error or incomplete data. The students are advised to write answer according to their understanding or write reasons for why it is not possible to solve partially or completely that question by citing errors/insufficient data.

FOR OFFICE USE ONLY

Question:	1	2	3	Total
Points:	5	6	9	20
Score:	0	0	4	04

QUESTIONS

1. (5 points) Let Y and ϵ are the response variable and the error in a regression problem, respectively. θ_1 and θ_2 are the two parameters. Let $y_i, i = 1, 2, 3$, be the observed values of the response variable. Consider,

$$Y_1 = \theta_1 + \theta_2 + \epsilon_1,$$

$$Y_2 = \theta_1 - 2\theta_2 + \epsilon_2,$$

$$Y_3 = 2\theta_1 - \theta_2 + \epsilon_3,$$

where $E(\epsilon_i) = 0$, and $V(\epsilon_i) = \sigma^2, i = 1, 2, 3; \sigma > 0$. Find the least square estimates (LSE) of θ_1 and θ_2 .

$$\rightarrow Y_1 = \theta_1 + \theta_2 + \epsilon_1,$$

$$Y_2 = \theta_1 - 2\theta_2 + \epsilon_2$$

$$Y_3 = 2\theta_1 - \theta_2 + \epsilon_3$$

$$\rightarrow E(\epsilon_i) = 0 \quad | \quad V(\epsilon_i) = \sigma^2; i = 1, 2, 3.$$

$$\sigma > 0.$$

$$\rightarrow \text{Let } Y = \theta_1 x_1 + \theta_2 x_2 + \epsilon_i.$$

$$\rightarrow \text{for } Y_1 \quad Y_1 = \theta_1 + \theta_2 + \epsilon_1$$

$$x_{11} = 1$$

$$x_{21} = 1$$

$$\rightarrow Y_2 = \theta_1 - 2\theta_2 + \epsilon_1$$

$$x_{12} = 1$$

$$x_{22} = -2$$

$$\rightarrow Y_3 = 2\theta_1 - \theta_2 + \epsilon_3 \Rightarrow \begin{matrix} x_{13} = 2 \\ x_{23} = -1 \end{matrix}$$

$$\rightarrow E[\hat{\theta}_1] = \theta_1$$

$$\rightarrow E[\hat{\theta}_2] = \theta_2$$

$$\rightarrow E(z^2) = \text{Var}(z) + (E(z))^2$$

$$\rightarrow \text{LSE}(\theta_1) = \text{Var}(\theta_1) + [E(\theta_1)]^2$$

$$\text{Var}(\theta_1) = \sigma^2 - E[\sum y_i^2] + \dots$$

$$\rightarrow E[\sum y_i] \Rightarrow S_{n, m_2}$$

$$\rightarrow \text{LSE}(\theta_1) = \sigma^2 - S_{n, m_2} + \frac{4}{3} S_{n, n_1}$$

similarly for the least square estimator of

(50)

$$E[\hat{\theta}_2] = \theta_2$$

$$\text{LSE}(\theta_2) = \text{Var}(\theta_2) + [E(\theta_2)]^2$$

$$\text{Var}(\theta_2) = \sigma^2 - E[\sum y_i^2]$$

$$\Rightarrow \text{LSE}(\theta_2) = \sigma^2 - S_{n_2, 4} + \frac{2}{3} S_{n_2, n_2}$$

2. Let U_1 and U_2 be two independent uniform random variables on the interval $(0, 1)$. Suppose that $\mathbf{X} = (X_1, X_2, X_3)'$, where $X_1 = U_1$, $X_2 = U_2$, and $X_3 = U_1 + U_2$.

(a) (3 points) Compute the correlation matrix ρ of $\mathbf{X}$.

(b) (3 points) Find the variance of first principal component based on the correlation matrix that you find in part (a).

(9) $\rightarrow \mathbf{X} = (X_1, X_2, X_3)'$
 $X_1 = U_1 \mid X_2 = U_2 \mid X_3 = U_1 + U_2$



$$\mathbf{X} = \begin{pmatrix} U_1 \\ U_2 \\ U_1 + U_2 \end{pmatrix}$$

$f(\mathbf{X}) =$ ~~scribbles~~


~~scribbles~~ Hat matrix:-

$$H = \mathbf{X} (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$$

$$\hat{\mathbf{X}} = H \mathbf{X}$$

$$= H (X_1, X_2, X_3)'$$

$$=$$



$$\mathbf{X} = \begin{bmatrix} U_1 \\ U_2 \\ U_1 + U_2 \end{bmatrix}$$

$$\mathbf{X}^T = \begin{bmatrix} U_1 & U_2 & U_1 + U_2 \end{bmatrix}$$

$$\mathbf{X}^T \mathbf{X} = \mathbf{X}^2$$

$$\Rightarrow H = \mathbf{X} (\mathbf{X}^2)^{-1} \mathbf{X}^T$$

Consider the system of linear equations

$$\begin{cases} x + 2y + 3z = 1 \\ 2x + 3y + 4z = 2 \\ 3x + 4y + 5z = 3 \end{cases}$$

$$\begin{bmatrix} 1 & 2 & 3 & 1 \\ 2 & 3 & 4 & 2 \\ 3 & 4 & 5 & 3 \end{bmatrix} \xrightarrow{R_2 - 2R_1, R_3 - 3R_1} \begin{bmatrix} 1 & 2 & 3 & 1 \\ 0 & -1 & -2 & 0 \\ 0 & -2 & -4 & 0 \end{bmatrix}$$

which is the same as the system

$$\begin{cases} x + 2y + 3z = 1 \\ -y - 2z = 0 \\ -2y - 4z = 0 \end{cases}$$

the second equation gives $y = -2z$. Substituting this into the first equation, we get

$$\begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 + R_2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 - R_2} \begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 + R_2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 - R_2} \begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 + R_2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \xrightarrow{R_1 - R_2} \begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

3. Let the matrix of distance be

$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & & & \\ 1 & 0 & & \\ 11 & 2 & 0 & \\ 5 & 3 & 3.5 & 0 \end{bmatrix} \end{matrix}$$

(a) (6 points) Cluster the four items using each of the following procedures:

- Complete linkage hierarchical procedure.
- Average linkage hierarchical procedure.

(b) (3 points) Draw the dendrograms and compare the results obtained using previous two procedures.

(i)

$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & & & \\ 1 & 0 & & \\ 11 & 2 & 0 & \\ 5 & 3 & 3.5 & 0 \end{bmatrix} \end{matrix}$$

Complete linkage hierarchical procedure, distance.
As uses the ~~concept~~ ~~of~~ minimum of the element in the matrix and links it.

$$\begin{matrix} & \begin{matrix} 1 & 2 & 3 & 4 \end{matrix} \\ \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & & & \\ \textcircled{1} & 0 & & \\ 11 & 2 & 0 & \\ 5 & 3 & 3.5 & 0 \end{bmatrix} \end{matrix}$$

$$\begin{matrix} & \begin{matrix} (21) & 3 & 4 \end{matrix} \\ \begin{matrix} (21) \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & & \\ 11 & 0 & \\ 5 & \textcircled{3.5} & 0 \end{bmatrix} \end{matrix}$$

$$\begin{matrix} & \begin{matrix} (21) & 3 & 4 \end{matrix} \\ \begin{matrix} (21) \\ 3 \\ 4 \end{matrix} & \begin{bmatrix} 0 & & \\ 11 & 0 & \\ 5 & \textcircled{3.5} & 0 \end{bmatrix} \end{matrix} \Rightarrow \begin{matrix} & \begin{matrix} (2143) \end{matrix} \\ \begin{matrix} (2143) \end{matrix} & \begin{bmatrix} 0 \end{bmatrix} \end{matrix}$$

$$\begin{matrix} & \begin{matrix} (21) & (43) \end{matrix} \\ \begin{matrix} (21) \\ (43) \end{matrix} & \begin{bmatrix} 0 & \\ \textcircled{11} & 0 \end{bmatrix} \end{matrix}$$

→ ~~For~~ For linkage, the complete linkage procedure, first links the variables with minimum distance, and as we move forward linking the other variables.

(ii) Average linkage hierarchical procedure:-
→ It takes the average distance b/w two variables and link them.

$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 0 & & & \\ 1 & 0 & & \\ 11 & 2 & 0 & \\ 5 & 3 & 3.5 & 0 \end{bmatrix}
 \end{array}
 \Rightarrow
 \begin{array}{c}
 \begin{matrix} (2,1) & (4,3) \end{matrix} \\
 \begin{matrix} (2,1) \\ (4,3) \end{matrix} \begin{bmatrix} 0 & & \\ 0 & & \\ 11 & & 0 \end{bmatrix}
 \end{array}$$



$$\Rightarrow d(n,y) = \frac{\sum (x_i - y_i)}{n}$$

→ For linkage, ~~average~~ 2 clusters has selected and then linked using their min. average distance.

03

ADDITIONAL PAGES FOR ANSWERS

Q (b) Dendrogram for Complete linkage hierarchical procedure:-



→ Dendrogram for Average linkage hierarchical procedure:-

