

PART-A

Name of the Student: Nelamab debnath

Signature of the Student: Nelamab debnath Roll No.: 230123041

Signature of the Invigilator: Ganra K. Mohy

INSTRUCTIONS

1. Check that you have **10 printed pages**. The PART-A of the examination has **3 questions**, for a total of **20 points**.
2. Attempt **all questions**.
3. There is **no credit** for a solution if the appropriate work is not shown, even if the answer is correct.
4. Write your answers in this booklet and only in the **space marked for answers**. Any answer written in any space other than the designated space will **not be evaluated**.
5. The pages numbered **8 to 10** are kept as **additional pages**. If you cannot able to fit a complete answer in the space provided just after the question, you can use these three pages by referring the page number of additional page in the original space.
6. At the beginning of the examination, a supplementary sheet has been provided only for rough work. Should you need more, please ask the invigilator.
7. No question requires any clarification from the instructors or invigilators, even if a question has an error or incomplete data. The students are advised to write answer according to their understanding or write reasons for why it is not possible to solve partially or completely that question by citing errors/insufficient data.

FOR OFFICE USE ONLY

Question:	1	2	3	Total
Points:	5	6	9	20
Score:	5	1	9	15

QUESTIONS

1. (5 points) Let Y and ϵ are the response variable and the error in a regression problem, respectively. θ_1 and θ_2 are the two parameters. Let $y_i, i = 1, 2, 3$, be the observed values of the response variable. Consider,

$$Y_1 = \theta_1 + \theta_2 + \epsilon_1,$$

$$Y_2 = \theta_1 - 2\theta_2 + \epsilon_2,$$

$$Y_3 = 2\theta_1 - \theta_2 + \epsilon_3,$$

where $E(\epsilon_i) = 0$, and $V(\epsilon_i) = \sigma^2, i = 1, 2, 3; \sigma > 0$. Find the least square estimates (LSE) of θ_1 and θ_2 .

Let us assume the three responses are $y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$

We know, for $y = X\beta + \epsilon$, the LSE value of β is

$$\hat{\beta} = (X^T X)^{-1} \cdot X^T y.$$

$$X = \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 2 & -1 \end{bmatrix}_{3 \times 2}, \quad X^T = \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & -1 \end{bmatrix}_{2 \times 3}$$

$$X^T X = \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 2 & -1 \end{bmatrix} = \begin{bmatrix} 6 & -3 \\ -3 & 6 \end{bmatrix}$$

$$(X^T X)^{-1} = \begin{bmatrix} 6 & -3 \\ -3 & 6 \end{bmatrix}^{-1} = \begin{bmatrix} 2/9 & 1/9 \\ 1/9 & 2/9 \end{bmatrix}_{2 \times 2}.$$

$$\therefore \hat{\beta} = (X^T X)^{-1} \cdot X^T y$$

$$\hat{\beta} = \begin{bmatrix} 2/9 & 1/9 \\ 1/9 & 2/9 \end{bmatrix}_{2 \times 2} \cdot \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & -1 \end{bmatrix}_{2 \times 3} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}_{3 \times 1}$$

$$2) \beta = \begin{bmatrix} 3/0 & 0 & 3/0 \\ 3/0 & -3/0 & 0 \end{bmatrix}_{2 \times 3} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}_{3 \times 1}$$

$$\Rightarrow \beta = \begin{bmatrix} 1/3 & 0 & 1/3 \\ 1/3 & -1/3 & 0 \end{bmatrix}_{2 \times 3} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}_{3 \times 1}$$

$$\Rightarrow [\hat{\theta}_1, \hat{\theta}_2]_{2 \times 1} = \begin{bmatrix} \frac{y_1 + y_3}{3}, \frac{y_1 - y_2}{3} \end{bmatrix}$$

$$\therefore \hat{\theta}_1 = \frac{y_1 + y_3}{3}, \quad \hat{\theta}_2 = \frac{y_1 - y_2}{3}$$

05

LS of θ_1 & θ_2 .

2. Let U_1 and U_2 be two independent uniform random variables on the interval $(0, 1)$. Suppose that $\mathbf{X} = (X_1, X_2, X_3)'$, where $X_1 = U_1$, $X_2 = U_2$, and $X_3 = U_1 + U_2$.
- (a) (3 points) Compute the correlation matrix ρ of $\mathbf{X}$.
- (b) (3 points) Find the variance of first principal component based on the correlation matrix that you find in part (a).

$$(a): \text{var}(X_1) = E[(U - E(U))^2] = \int_0^1 \left(u - \frac{1}{2}\right)^2 \cdot \text{den} \\ = \frac{1}{3} - \frac{1}{2} + \frac{1}{4} = \frac{1}{12}$$

$$\text{var}(X_2) = \frac{1}{12}$$

$$\text{var}(X_1 + X_2) =$$

$$\text{var}(X_3) = \text{var}(U_1 + U_2) = \text{var}(U_1) + \text{var}(U_2) \quad \left[\begin{array}{l} U_1 \& U_2 \\ \text{are independent} \end{array} \right] \\ = \frac{2}{12} = \frac{1}{6}$$

$$\text{cov}(X_1, X_2) = E[(X_1 - E(X_1))(X_2 - E(X_2))] \\ = \int_0^1 \int_0^1 \left(u - \frac{1}{2}\right) \left(y - \frac{1}{2}\right) \text{den} \\ = \frac{1}{12} \cdot \frac{1}{12} = \frac{1}{144} \quad \text{X}$$

$$\text{cov}(X_1, X_3) = \text{cov}(X_1, U_1 + U_2) = \text{cov}(X_1, U_1) + \text{cov}(X_1, U_2)$$

$$\left\{ \begin{array}{l} \text{cov}(X_1, U_1) = \text{cov}(X_1, X_1) \\ \text{cov}(X_1, U_2) = \text{cov}(X_1, X_2) \end{array} \right\} = \text{var}(X_1) + \text{cov}(X_1, X_2) = \frac{1}{12} + \frac{1}{144} = \frac{13}{144}$$

$$\therefore \text{cov}(X_2, X_3) = 13/144$$

$\therefore$ Correlation Matrix

$$\rho = \begin{matrix} & \begin{matrix} X_1 & X_2 & X_3 \end{matrix} \\ \begin{matrix} X_1 \\ X_2 \\ X_3 \end{matrix} & \begin{bmatrix} 1 & 1/12 & 13/12\sqrt{2} \\ 1/12 & 1 & 13/12\sqrt{2} \\ 13/12\sqrt{2} & 13/12\sqrt{2} & 1 \end{bmatrix} \end{matrix} = \begin{matrix} & \begin{matrix} X_1 & X_2 & X_3 \end{matrix} \\ \begin{matrix} X_1 \\ X_2 \\ X_3 \end{matrix} & \begin{bmatrix} 1 & 0.083 & 0.76 \\ 0.083 & 1 & 0.766 \\ 0.766 & 0.766 & 1 \end{bmatrix} \end{matrix}$$

$$\text{corr}(X_1, X_3) = \frac{13/144}{\sqrt{1/12} \cdot \sqrt{1/6}} = \frac{13/144}{\sqrt{2}/12} = \frac{13}{12\sqrt{2}}$$

②: (b): first principal component $PC_1 = e_1^T \cdot X$
 $e_1 \rightarrow$ largest value corresponding eigen vector.

$$\text{var}(PC_1) = \text{var}(e_1^T X) = \underline{e_1^T \cdot \text{var}(X) \cdot e_1}$$

variance of largest eigen vector $\rightarrow$ largest eigen vector of P .

$$\begin{vmatrix} 1-\lambda & 0.083 & 0.766 \\ 0.083 & 1-\lambda & 0.766 \\ 0.766 & 0.766 & 1-\lambda \end{vmatrix} = 0$$

$$\Rightarrow (1-\lambda) \left[\underbrace{(1-\lambda)^2}_{1-2\lambda+\lambda^2} - 0.5867 \right] \\ - 0.083 (0.083(1-\lambda) - 0.5867) \\ + 0.766 (0.08357 - (1-\lambda) \cdot 766) = 0$$

$$\Rightarrow -\lambda^3 + \lambda^2 + 0.8196\lambda - 0.078204 = 0$$

$$\Rightarrow \lambda = \underline{0.109}, \quad 0.4459, \quad 0.4454$$

λ_1
largest eigen value

01 $\Rightarrow \underline{\underline{\lambda = 0.109}}$

3. Let the matrix of distance be

	1	2	3	4
1	0			
2	1	0		
3	11	2	0	
4	5	3	3.5	0

(a) (6 points) Cluster the four items using each of the following procedures:

- Complete linkage hierarchical procedure.
- Average linkage hierarchical procedure.

(b) (3 points) Draw the dendrograms and compare the results obtained using previous two procedures.

(a): (i.) Complete linkage:

$$d_{u,w} = \max_{u \in U, v \in V} \{d_{u,v}\}$$

$i=0$

	1	2	3	4
1	0			
2	1	0		
3	11	2	0	
4	5	3	3.5	0

→ minimum → 1 → {1 & 2}

∴ clusters: {12}, {3}, {4}

$$\& d_{\{12\}, \{3\}} = \max\{11, 2\} = 11.$$

$$d_{\{12\}, \{4\}} = \max\{5, 3\} = 5, \quad d_{3,4} = 3.5$$

$i=1$

	{12}	3	4
{12}	0		
3	11	0	
4	5	3.5	0

→ minimum distance = 3.5 → {3 & 4}

clusters: {12}, {34}

$$\therefore d_{\{12\}, \{34\}} = \max\{11, 2, 5, 3.5\} = 11$$

$$\begin{array}{c} \{12\} \quad \{34\} \\ \{12\} \begin{bmatrix} 0 & \\ & \end{bmatrix} \\ \{34\} \begin{bmatrix} 11 & 0 \end{bmatrix} \end{array} \rightarrow \text{min distance} = 11 \rightarrow \{1234\}.$$

clustering complete //

(a): (ii) Average linkage:

$$d_{uw} = \frac{\left(\sum_{\substack{u \in U \\ w \in W}} d_{u,w} \right)}{\underbrace{|U| \cdot |W|}_{\text{total no. of linkages between } U \text{ \& } W}}$$

$$\begin{array}{c} 1 \quad 2 \quad 3 \quad 4 \\ \begin{bmatrix} 0 & & & \\ 1 & 0 & & \\ 11 & 2 & 0 & \\ 5 & 3 & 3.5 & 0 \end{bmatrix} \end{array} \rightarrow \text{min distance} = 1 \rightarrow \{12\}.$$

clusters formed $\rightarrow \{12\}, \{3\}, \{4\}.$

$$d_{\{12\}, \{3\}} = \frac{11+2}{2} = 6.5, \quad d_{3,4} = 3.5$$

$$d_{\{12\}, \{4\}} = \frac{5+3}{2} = 4.$$

$$\begin{array}{c} \{12\} \quad \{34\} \\ \{12\} \begin{bmatrix} 0 & & \\ 6.5 & 0 & \\ 4 & 3.5 & 0 \end{bmatrix} \\ \{34\} \end{array} \rightarrow \text{min distance} = 3.5 \rightarrow \{34\}.$$

clusters: $\{12\}, \{34\}.$

$$d_{\{12\}, \{34\}} = \frac{11+2+5+3}{4} = \frac{21}{4} = 5.25$$



(Q6 Ans continued)

(a) (ii): {12} {34}

(06) $\begin{matrix} \{2\} \\ \{34\} \end{matrix} \begin{bmatrix} 0 \\ 5.25 \end{bmatrix} \begin{matrix} \text{min distance} \rightarrow 5.25 \\ \rightarrow \end{matrix} \begin{matrix} \{123\} \\ \{1234\} \end{matrix}$

 $\therefore$ clusters $\rightarrow \{123\} \{1234\} \rightarrow$ clustering complete.

(a) (b): Complete dendrogram:

average linkage:

{12} & {34} merge at same distance in both processes but the final cluster {1234} is formed earlier in the average linkage.

