

PART-A

Name of the Student: Samir Raj
Signature of the Student: Samir Roll No.: 230123056
Signature of the Invigilator: Ganar Ku. Meher

INSTRUCTIONS

1. Check that you have **10 printed pages**. The PART-A of the examination has **3 questions**, for a total of **20 points**.
2. Attempt **all questions**.
3. There is **no credit** for a solution if the appropriate work is not shown, even if the answer is correct.
4. Write your answers in this booklet and only in the **space marked for answers**. Any answer written in any space other than the designated space will **not be evaluated**.
5. The pages numbered **8 to 10** are kept as **additional pages**. If you cannot able to fit a complete answer in the space provided just after the question, you can use these three pages by referring the page number of additional page in the original space.
6. At the beginning of the examination, a supplementary sheet has been provided only for rough work. Should you need more, please ask the invigilator.
7. No question requires any clarification from the instructors or invigilators, even if a question has an error or incomplete data. The students are advised to write answer according to their understanding or write reasons for why it is not possible to solve partially or completely that question by citing errors/insufficient data.

FOR OFFICE USE ONLY

Question:	1	2	3	Total
Points:	5	6	9	20
Score:	00	0	9	09

QUESTIONS

1. (5 points) Let Y and ϵ are the response variable and the error in a regression problem, respectively. θ_1 and θ_2 are the two parameters. Let $y_i, i = 1, 2, 3$, be the observed values of the response variable. Consider,

$$Y_1 = \theta_1 + \theta_2 + \epsilon_1,$$

$$Y_2 = \theta_1 - 2\theta_2 + \epsilon_2,$$

$$Y_3 = 2\theta_1 - \theta_2 + \epsilon_3,$$

where $E(\epsilon_i) = 0$, and $V(\epsilon_i) = \sigma^2, i = 1, 2, 3; \sigma > 0$. Find the least square estimates (LSE) of θ_1 and θ_2 .

$$\epsilon_1 = Y_1 - \theta_1 - \theta_2$$

$$\epsilon_2 = Y_2 - \theta_1 + 2\theta_2 \rightarrow (3)$$

$$\epsilon_3 = Y_3 - 2\theta_1 + \theta_2 \rightarrow (2)$$

$$Y_3 - Y_2 - Y_1 = \epsilon_3 - \epsilon_2 - \epsilon_1$$

$$E(\epsilon_i) = 0$$

$$\Rightarrow \frac{\epsilon_1 + \epsilon_2 + \epsilon_3}{3} = 0$$

$$\Rightarrow \epsilon_1 + \epsilon_2 + \epsilon_3 = Y_1 + Y_2 + Y_3 - 4\theta_1 + 2\theta_2$$

$$\Rightarrow (\epsilon_1 + \epsilon_2 + \epsilon_3)^2 = (Y_1 + Y_2 + Y_3 - 4\theta_1 + 2\theta_2)^2$$

Differentiate w.r.t to θ_1

$$\Rightarrow 0 = 2(Y_1 + Y_2 + Y_3 - 4\hat{\theta}_1 + 2\hat{\theta}_2)(-4)$$

Differentiate w.r.t θ_2

$$\Rightarrow 0 = 2(Y_1 + Y_2 + Y_3 - 4\hat{\theta}_1 + 2\hat{\theta}_2)(2)$$

$$\Rightarrow Y_1 + Y_2 + Y_3 - 4\theta_1 + 2\theta_2 = 0 \rightarrow (4)$$

$$\epsilon_1^2 = (Y_1 - \theta_1 - \theta_2)^2$$

Differentiate w.r.t θ_1

$$\Rightarrow 0 = 2(Y_1 - \theta_1 - \theta_2)(-1) \Rightarrow \hat{\theta}_1 = Y_1 - \theta_2$$

Similarly for eqn ① and ③

$$\theta_1 = Y_2 + 2\theta_2$$

$$\theta_1 = \frac{Y_3 + \theta_2}{2}$$

$$\Rightarrow \hat{\theta}_1 = \frac{Y_1 - \theta_2 + Y_2 + 2\theta_2 + \frac{Y_3 + \theta_2}{2}}{3}$$

$$\Rightarrow \hat{\theta}_1 = \frac{2Y_1 + 2Y_2 + Y_3 - 2\hat{\theta}_2 + 4\hat{\theta}_2 + \hat{\theta}_2}{6}$$

$$\Rightarrow \hat{\theta}_1 = \frac{2Y_1 + 2Y_2 + Y_3 + 3\hat{\theta}_2}{6} \rightarrow \text{⑤}$$

same as

from ④

$$\hat{\theta}_1 = \frac{Y_1 + Y_2 + Y_3 + 2\hat{\theta}_2}{4}$$

Equate

$$\Rightarrow \frac{Y_1 + Y_2 + Y_3 + 2\hat{\theta}_2}{4} = \frac{2Y_1 + 2Y_2 + Y_3 + 3\hat{\theta}_2}{6}$$

$$\Rightarrow 3Y_1 + 3Y_2 + 3Y_3 + 6\hat{\theta}_2 = 4Y_1 + 4Y_2 + 2Y_3 + 6\hat{\theta}_2$$

$$\Rightarrow Y_1 + Y_2 - Y_3 = 0$$

$$\text{Var}(\varepsilon_i) = E(\varepsilon_i^2) - (E(\varepsilon_i))^2 = E(\varepsilon_i^2)$$

$$\Rightarrow \sigma^2 = \frac{(Y_1 - \theta_1 - \theta_2)^2 + (Y_2 - \theta_1 + 2\theta_2)^2 + (Y_3 - 2\theta_1 + 2\theta_2)^2}{3}$$

Differentiate w.r.t θ_1

$$\Rightarrow 6\hat{\theta}_1 = Y_1 + Y_2 + 2Y_3 + 5\hat{\theta}_2$$

Equate with equation

$$\Rightarrow \hat{\theta}_2 = \frac{Y_1 + Y_2 - Y_3}{4}$$

⑤

$$\Rightarrow \hat{\theta}_1 = \frac{3Y_1 + 3Y_2 + Y_3}{8}$$

LSE of $\theta_1 = \hat{\theta}_1$
LSE of $\theta_2 = \hat{\theta}_2$

2. Let U_1 and U_2 be two independent uniform random variables on the interval $(0, 1)$. Suppose that $\mathbf{X} = (X_1, X_2, X_3)'$, where $X_1 = U_1$, $X_2 = U_2$, and $X_3 = U_1 + U_2$.

(a) (3 points) Compute the correlation matrix ρ of $\mathbf{X}$.

(b) (3 points) Find the variance of first principal component based on the correlation matrix that you find in part (a).

a)

$$\text{Cor}(X_1, X_2) = 0 \quad (\text{Random } U_1 \text{ and } U_2)$$

$$\text{Cor}(X_2, X_3) = \frac{1}{2} \quad (X_3 = U_1 + U_2, \text{ Half by } X_2 = U_2)$$

$$\text{Cor}(X_1, X_3) = \frac{1}{2} \quad (X_3 = U_1 + U_2, \text{ Half by } X_1 = U_1)$$

Correlation matrix of $\mathbf{X}$,

$$\rho = \begin{bmatrix} 1 & 0 & 1/2 \\ 0 & 1 & 1/2 \\ 1/2 & 1/2 & 1 \end{bmatrix}$$

b)

$$\begin{aligned} \text{Var}(\mathbf{X}) &= \text{Var}(X_1) + \text{Var}(X_2) + \text{Var}(X_3) \\ &\quad + \text{Cor}(X_1, X_2) + \text{Cor}(X_2, X_3) \\ &\quad + \text{Cor}(X_1, X_3) \end{aligned}$$

~~Correlation matrix~~

$$\text{Correlation}(X_1, X_2) = \frac{\text{Covariance}(X_1, X_2)}{\sqrt{\text{Var}(X_1)} \sqrt{\text{Var}(X_2)}}$$

$$\Rightarrow \text{Var}(\mathbf{X}) = 1 + 1 + \frac{1}{2} + 0 + \frac{1}{2}(\sqrt{1})(\sqrt{\frac{1}{2}}) + \frac{1}{2}(\sqrt{1})(\sqrt{\frac{1}{2}})$$

$$\begin{aligned}
 \Rightarrow \text{Var}(X) &= \frac{5}{2} + \frac{1}{2\sqrt{2}} + \frac{1}{2\sqrt{2}} \\
 &= \frac{5}{2} + \frac{\sqrt{2}}{4} + \frac{\sqrt{2}}{4} \\
 &= \frac{5}{2} + \frac{\sqrt{2}}{2} = \frac{5+\sqrt{2}}{2}
 \end{aligned}$$

$$\Rightarrow \text{Var}(X) = \frac{5+\sqrt{2}}{2} = 3.2071$$

$$\begin{aligned}
 \text{Var}(U(a,b)) &= \frac{1}{b-a}, \quad \text{Var}(X_1) = \text{Var}(X_2) = \frac{1}{1-0} \\
 \Rightarrow \text{Var}(X_3) &= \frac{1}{2-0} = \frac{1}{2} = 1
 \end{aligned}$$

Variance of first principal component,

~~$$\begin{aligned}
 &= \frac{\text{Var}(X_1)}{\text{Var}(X)} = \frac{1}{\frac{5+\sqrt{2}}{2}} = 0.311807 \\
 &= 31.1807\%
 \end{aligned}$$~~

~~$$\begin{aligned}
 &= \frac{\frac{1}{2} + \frac{1}{2}(1)(\frac{\sqrt{2}}{2})}{\frac{5+\sqrt{2}}{2}} = \frac{1 + \frac{\sqrt{2}}{2}}{5+\sqrt{2}} = 0.22048
 \end{aligned}$$~~

By X_1

$$\begin{aligned}
 &= \frac{1 + \frac{1}{2}\sqrt{1}\left(\frac{\sqrt{2}}{2}\right)^{\frac{1}{2}}}{\frac{5+\sqrt{2}}{2}} = 0.366927 \\
 &= 36.6927\%
 \end{aligned}$$

3. Let the matrix of distance be

	1	2	3	4
1	0			
2	1	0		
3	11	2	0	
4	5	3	3.5	0

(a) (6 points) Cluster the four items using each of the following procedures:

- Complete linkage hierarchical procedure.
- Average linkage hierarchical procedure.

(b) (3 points) Draw the dendrograms and compare the results obtained using previous two procedures.

3) a) i)

$d_{uv} = \max(d_{uw}, d_{vw})$

Initial matrix:

	1	2	3	4
1	0			
2	1	0		
3	11	2	0	
4	5	3	3.5	0

Step 1: $\max(11, 5) = 11$, $\max(5, 3) = 5$. Merge (1, 2) and (3, 4).

Matrix after first merge:

	(1,2)	(3,4)
(1,2)	0	
(3,4)	11	0

Step 2: $\max(11, 5) = 11$. Merge ((1,2), (3,4)).

Final matrix:

	((1,2), (3,4))
((1,2), (3,4))	0

Clusters: $(1), (2), (3), (4) \xrightarrow{(1,2)} (1,2), (3), (4) \xrightarrow{(3,4)} (1,2), (3,4) \xrightarrow{11} ((1,2), (3,4))$

ii) $d_{uv} = \frac{\sum_i d_{ik}}{N_i N_k}$

Initial matrix:

	1	2	3	4
1	0			
2	1	0		
3	11	2	0	
4	5	3	3.5	0

Step 1: Merge (1, 2) and (3, 4).

Matrix after first merge:

	(1,2)	(3,4)
(1,2)	0	
(3,4)	6.5	0

$d_{(1,2)(3)} = \frac{11+2}{2 \times 1} = \frac{13}{2} = 6.5$

$d_{(1,2)(4)} = \frac{5+3}{2 \times 1} = \frac{8}{2} = 4$



Step 2: Merge ((1,2), (3,4)).

$$\begin{array}{ccc}
 (1,2,3,4) & (1,2,3,4) & (1,2) \quad (3,4) \\
 (1,2,3,4) \begin{bmatrix} 0 \end{bmatrix} & \leftarrow & \begin{bmatrix} 0 & \\ (5.25) & 0 \end{bmatrix}
 \end{array}$$

$$\begin{aligned}
 d_{(1,2)(3,4)} &= \frac{11+5+2+3}{2 \times 2} \\
 &= \frac{21}{4} = 5.25
 \end{aligned}$$

Clusters : $(1,2), (3,4) \xrightarrow[\text{@ } 1]{(1,2)} (1,2), (3), (4) \xrightarrow[\text{@ } 3.5]{(3,4)} (1,2), (3,4) \xrightarrow[\text{@ } 5.25]{(1,2,3,4)} (1,2,3,4)$

b) i) Dendrogram for complete linkage ii) Dendrogram for average linkage



Ans we can see according to complete linkage hierarchical procedure cluster $(1,2)$ and $(3,4)$ are less similar to each other compared to the inference of average linkage hierarchical procedure.

Average linkage if $(3,4)$ are similar then $(1,2,3,4)$ are also something near that

Complete linkage if $(3,4)$ are similar still $(1,2,3,4)$ has ~~good~~ difference. Formation of cluster $(1,2)$ and $(3,4)$ is similar in both process.

ADDITIONAL PAGES FOR ANSWERS

Rough for Question 1
Continued calculation

$$\sigma^2 = \frac{(Y_1 - \theta_1 - \theta_2)^2 + (Y_2 - \theta_1 + 2\theta_2)^2 + (Y_3 - 2\theta_1 + 2\theta_2)^2}{3}$$

$$\Rightarrow 0 = -2(Y_1 - \theta_1 - \theta_2) - 2(Y_2 - \theta_1 + 2\theta_2) - 4(Y_3 - 2\theta_1 + 2\theta_2)$$

$$\Rightarrow 6\theta_1 = Y_1 + Y_2 + 2Y_3 + 5\theta_2$$

~~and~~ Equate

$$\frac{Y_1 + Y_2 + 2Y_3 + 5\theta_2}{6} = \frac{Y_1 + Y_2 + Y_3 + 2\theta_2}{4}$$

$$\Rightarrow 4\theta_2 = Y_1 + Y_2 - Y_3$$

Put in eqn (4)

$$\theta_1 = \frac{Y_1 + Y_2 + 2Y_3 + \frac{5}{4}(Y_1 + Y_2 - Y_3)}{6}$$

$$= \frac{9Y_1 + 9Y_2 + 3Y_3}{24} = \frac{3Y_1 + 3Y_2 + Y_3}{8}$$

