

PART-A

Name of the Student: Vedant Sunil Vanjari
Signature of the Student: Vanjari Roll No.: 230123069
Signature of the Invigilator: Shan

INSTRUCTIONS

1. Check that you have **10 printed pages**. The PART-A of the examination has **3 questions**, for a total of **20 points**.
2. Attempt **all questions**.
3. There is **no credit** for a solution if the appropriate work is not shown, even if the answer is correct.
4. Write your answers in this booklet and only in the **space marked for answers**. Any answer written in any space other than the designated space will **not be evaluated**.
5. The pages numbered **8 to 10** are kept as **additional pages**. If you cannot able to fit a complete answer in the space provided just after the question, you can use these three pages by referring the page number of additional page in the original space.
6. At the beginning of the examination, a supplementary sheet has been provided only for rough work. Should you need more, please ask the invigilator.
7. No question requires any clarification from the instructors or invigilators, even if a question has an error or incomplete data. The students are advised to write answer according to their understanding or write reasons for why it is not possible to solve partially or completely that question by citing errors/insufficient data.

FOR OFFICE USE ONLY

Question:	1	2	3	Total
Points:	5	6	9	20
Score:	3	0	9	12

QUESTIONS

1. (5 points) Let Y and ϵ are the response variable and the error in a regression problem, respectively. θ_1 and θ_2 are the two parameters. Let $y_i, i = 1, 2, 3$, be the observed values of the response variable. Consider,

$$Y_1 = \theta_1 + \theta_2 + \epsilon_1,$$

$$Y_2 = \theta_1 - 2\theta_2 + \epsilon_2,$$

$$Y_3 = 2\theta_1 - \theta_2 + \epsilon_3,$$

where $E(\epsilon_i) = 0$, and $V(\epsilon_i) = \sigma^2, i = 1, 2, 3; \sigma > 0$. Find the least square estimates (LSE) of θ_1 and θ_2 .

Let

$$Q(\theta_1, \theta_2) = \sum \epsilon_i^2$$

$$= \sum_{i=1}^3 (y_i - x_{i1}\theta_1 - x_{i2}\theta_2)^2$$

$$Q(\theta_1, \theta_2) = (Y - X^T B)^T (Y - X^T B)$$

where $Y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ $X = \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 2 & -1 \end{bmatrix}$

$$Q(\theta_1, \theta_2) = (Y - XB)^T (Y - XB)$$

where $Y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ $X = \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 2 & -1 \end{bmatrix}$ $B = \begin{bmatrix} \theta_1 \\ \theta_2 \end{bmatrix}$

$$\therefore Q(\beta) = y^T y - \beta^T x^T y - y^T x \beta + \beta^T x^T x \beta$$

now,

$\beta^T x^T y$ is scalar

$$\therefore (\beta^T x^T y)^T = y^T x \beta$$

and transpose of a scalar is equal to itself

$$\therefore Q(\beta) = y^T y - 2\beta^T x^T y + \beta^T x^T x \beta$$

now, we want to minimize this for LSE's

$$\Rightarrow \hat{\beta} = (x^T x)^{-1} x^T y$$

$$\therefore x^T x = \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & -2 \\ 2 & -1 \end{bmatrix} = \begin{bmatrix} 6 & -3 \\ -3 & 4 \end{bmatrix}$$

~~$$(x^T x)^{-1} = \frac{1}{15} \begin{bmatrix} 4 & 3 \\ 3 & 6 \end{bmatrix}$$~~

$$(x^T x)^{-1} = \frac{1}{15} \begin{bmatrix} 4 & 3 \\ 3 & 6 \end{bmatrix}$$

$$\therefore \hat{\beta} = \frac{1}{15} \begin{bmatrix} 4 & 3 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 1 & -2 & -1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

$$= \frac{1}{15} \begin{bmatrix} 7 & -2 & 5 \\ 9 & -9 & 0 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$$

03

→ These are LSE's of θ_1 & θ_2 resply.

$$\hat{\theta}_1 = \frac{1}{15} [7y_1 - 2y_2 + 5y_3]$$

$$\hat{\theta}_2 = \frac{1}{15} [9y_1 - 9y_2]$$

2. Let U_1 and U_2 be two independent uniform random variables on the interval $(0, 1)$. Suppose that $\mathbf{X} = (X_1, X_2, X_3)'$, where $X_1 = U_1$, $X_2 = U_2$, and $X_3 = U_1 + U_2$.

(a) (3 points) Compute the correlation matrix ρ of $\mathbf{X}$.

(b) (3 points) Find the variance of first principal component based on the correlation matrix that you find in part (a).

(a) Let the correlation matrix of $\mathbf{X}$ be ρ .

$$\Rightarrow \rho = \begin{bmatrix} \text{Corr}(X_1, X_1) & \text{Corr}(X_1, X_2) & \text{Corr}(X_1, X_3) \\ \text{Corr}(X_2, X_1) & \text{Corr}(X_2, X_2) & \text{Corr}(X_2, X_3) \\ \text{Corr}(X_3, X_1) & \text{Corr}(X_3, X_2) & \text{Corr}(X_3, X_3) \end{bmatrix}$$

$$\text{Now, } \text{Corr}(X_i, X_j) = \frac{\text{Cov}(X_i, X_j)}{\sqrt{\text{Var}(X_i)} \sqrt{\text{Var}(X_j)}}$$

$$\text{Var}(X_1) = \frac{1}{12}, \quad \text{Var}(X_2) = \frac{1}{12}, \quad \text{Var}(X_3) = \frac{1}{6}$$

$$\text{Cov}(X_1, X_2) = \text{Cov}(X_2, X_1) = \frac{1}{12}$$

$$\text{Cov}(X_1, X_3) = \text{Cov}(X_3, X_1) = \frac{1}{6}$$

$$\text{Cov}(X_2, X_3) = \text{Cov}(X_3, X_2) = \frac{1}{6}$$

$$\rho = \begin{bmatrix} 1 & \frac{1/12}{\sqrt{1/12}\sqrt{1/12}} & \frac{1/6}{\sqrt{1/6}\sqrt{1/12}} \\ \frac{1/12}{\sqrt{1/12}\sqrt{1/12}} & 1 & \frac{1/6}{\sqrt{1/6}\sqrt{1/12}} \\ \frac{1/6}{\sqrt{1/6}\sqrt{1/12}} & \frac{1/6}{\sqrt{1/6}\sqrt{1/12}} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & \sqrt{2} \\ 1 & 1 & \sqrt{2} \\ \sqrt{2} & \sqrt{2} & 1 \end{bmatrix}$$

(b) Now, we find eigenvectors.

$$P - \lambda I = 0$$

$\Rightarrow$ from this we get.

$$\lambda = 0$$

$$\lambda = \frac{3 \pm \sqrt{17}}{2}$$

highest eigenvalue = $\lambda_1 = \frac{3 + \sqrt{17}}{2}$

$\therefore$ let e_1 be the eigenvector corresponding to λ_1 .

$$y = e_1^T x$$

~~$\therefore$~~
now

$$\text{Var}(y) = \cancel{e_1^T} x e$$

$$\text{Var}(y) = e_1^T P e$$

this is the variance of first principal component

3. Let the matrix of distance be

$$\begin{array}{c}
 \begin{matrix} & 1 & 2 & 3 & 4 \end{matrix} \\
 \begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 0 & & & \\ 1 & 0 & & \\ 11 & 2 & 0 & \\ 5 & 3 & 3.5 & 0 \end{bmatrix}
 \end{array}$$

- (a) (6 points) Cluster the four items using each of the following procedures:
- Complete linkage hierarchical procedure.
 - Average linkage hierarchical procedure.
- (b) (3 points) Draw the dendrograms and compare the results obtained using previous two procedures.

① we take min^m distance i.e. btwn 1 & 2.

∴ new matrix.

$$\begin{array}{c}
 \begin{matrix} & (12) & 3 & 4 \end{matrix} \\
 \begin{matrix} (12) \\ 3 \\ 4 \end{matrix} \begin{bmatrix} 0 & & \\ 11 & 0 & \\ 5 & 3.5 & 0 \end{bmatrix}
 \end{array}$$

As here we are seeing complete linkage hierarchical procedure,

$$d_{(u,v),w} = \max\{d_{u,w}, d_{v,w}\}$$

for u, v in same group.

now, again we cluster using minimum distance.

$$\begin{array}{cc} (12) & (34) \\ (12) & \begin{bmatrix} 0 & \\ & \end{bmatrix} \\ (34) & \begin{bmatrix} 11 & 0 \end{bmatrix} \end{array}$$

now this will cluster again

$$\begin{array}{c} (1234) \\ (1234) \begin{bmatrix} 0 \end{bmatrix} \end{array}$$

(ii) Average linkage hierarchical procedure.

$$\begin{array}{cc} & \begin{array}{cc} (12) & 3 & 4 \end{array} \\ \begin{array}{c} (12) \\ 3 \\ 4 \end{array} & \begin{bmatrix} 0 & & \\ 6.5 & 0 & \\ 4 & 3.5 & 0 \end{bmatrix} \end{array}$$

$$\text{new values} = \arg d_{(u,v),w} \quad u,v \in G$$

Ob^{eg.}

$$d_{(12),3} = \frac{d_{1,3} + d_{1,2}}{2} = \frac{11 + 2}{2} = 6.5$$

Now, next step.


$$\begin{array}{cc} (12) & (34) \\ (12) & \begin{bmatrix} 0 & \\ & \end{bmatrix} \\ (34) & \begin{bmatrix} 5.25 & 0 \end{bmatrix} \end{array} \Rightarrow (1234) \begin{bmatrix} 0 \end{bmatrix}$$

$$\begin{aligned} d_{(12),(34)} &= \frac{d_{1,3} + d_{1,4} + d_{2,3} + d_{2,4}}{4} \\ &= \frac{11 + 5 + 2 + 3}{4} \\ &= 5.25 \end{aligned}$$


Q3.

(b) Drawing dendrograms

for (i)



for (ii)



We can see that, in the average linkage hierarchical procedure, the highest value of distance is less than that of in complete linkage.

OB

