

PART-B

Name of the Student: Samsiddhi Dasgupta
Signature of the Student: [Signature] Roll No.: 230123057
Signature of the Invigilator: [Signature]

INSTRUCTIONS

1. Check that you have **10 printed pages**. The PART-B of the examination has **4 questions**, for a total of **20 points**.
2. Attempt **all questions**.
3. There is **no credit** for a solution if the appropriate work is not shown, even if the answer is correct.
4. Write your answers in this booklet and only in the **space marked for answers**. Any answer written in any space other than the designated space will **not be evaluated**.
5. The pages numbered **8 to 10** are kept as **additional pages**. If you cannot able to fit a complete answer in the space provided just after the question, you can use these three pages by referring the page number of additional page in the original space.
6. At the beginning of the examination, a supplementary sheet has been provided only for rough work. Should you need more, please ask the invigilator.
7. No questions require any clarification from the instructor or invigilators, even if a question has an error or incomplete data. The students are advised to write answer according to their understanding or write reasons for why it is not possible to solve partially or completely that question by citing errors/insufficient data.

FOR OFFICE USE ONLY

Question:	1	2	3	4	Total
Points:	6	7	5	2	20
Score:	0	5	5	2	12

QUESTIONS

1. (6 points) Let

$$S = \begin{pmatrix} S_{yy} & S_{yX} \\ S_{yX} & S_{XX} \end{pmatrix}$$

be the partitioned sample covariance matrix of $Y_{1 \times 1}$ and $X_{q \times 1}$ based on a centered (with respect to sample mean) sample of size n . Prove that the squared sample first canonical correlation r_1^2 is identical to the coefficient of determination (R^2) obtained from the multiple linear regression of Y on the vector X .

when finding sample ^{just} canonical correlation
we are trying to maximize the correlation
between 2 qty $\rightarrow$ linear comb. of X , and
linear combination of Y

essentially

$$\max_{\text{Corr}} (bY, a_1X_1 + a_2X_2 + \dots + a_qX_q)$$

on performing multiple linear regression of Y on X
if we keep the coefficients of X_i as obtained
 ~~$a = (X^T X)^{-1} X^T Y$~~ in the first canonical variable
(and multiply with a factor and add bias)

$$R^2 = \frac{SS_{\text{Reg}}}{SS_{\text{T}}} \rightarrow \text{proportion of variance explained by the regression}$$

$$= 1 - \frac{SS_{\text{Res}}}{SS_{\text{T}}}$$

so if we use the a , we will be able to have
 $R^2 = r_1^2$

while performing multiple linear regression we
minimize SS_{Res} or maximize R^2

and as we know from canonical variables,
there is no other linear combination of X which has

higher correlation with y than the first canonical variable x_1 .

So the result obtained by MLR is same as just canonical variable

00

2. Suppose you are performing a factor analysis on a 4×4 sample covariance matrix $\mathbf{S}$. The eigenvalues of $\mathbf{S}$ are $\lambda_1 = 0.8$, $\lambda_2 = 2.1$, $\lambda_3 = 4.2$, and $\lambda_4 = 0.2$. The corresponding normalized eigenvectors are $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$, and $\mathbf{e}_4$. You decide to extract $m = 2$ factors using the principal component method.
- (2 points) Write down the explicit mathematical expression for the estimated factor loading matrix $\tilde{\mathbf{L}}$ in terms of the given eigenvalues and eigenvectors.
 - (1 point) Let the first row of $\tilde{\mathbf{L}}$ be $\tilde{\mathbf{l}}_1^T = [0.8, -0.4]$. Calculate the estimated communality $\tilde{h}_1^2$ for the first variable.
 - (4 points) Prove that the sum of all estimated communalities $\sum_{i=1}^p \tilde{h}_i^2$ is exactly equal to the sum of the largest $m = 2$ eigenvalues of $\mathbf{S}$.

$$a) \quad \tilde{\mathbf{L}} = [\sqrt{\lambda_{(1)}} \mathbf{e}_{(1)} \mid \sqrt{\lambda_{(2)}} \mathbf{e}_{(2)} \mid \dots \mid \sqrt{\lambda_{(m)}} \mathbf{e}_{(m)}]$$

in our case $m = 2$ so ~~so~~

$$\tilde{\mathbf{L}} = [\sqrt{\lambda_{(1)}} \mathbf{e}_{(1)} \mid \sqrt{\lambda_{(2)}} \mathbf{e}_{(2)}]$$

* where $\lambda_{(i)}$ is the i^{th} largest eigenvalue and $\mathbf{e}_{(i)}$ is the eigenvector associated with $\lambda_{(i)}$

$$b) \quad \tilde{h}_1^2 = \lambda_{(1)}^2 + \lambda_{(2)}^2 = 0.8^2 + (-0.4)^2 = \underline{\underline{0.8}}$$

$$c) \quad \sum_{i=1}^p \tilde{h}_i^2 = \sum_{j=1}^p \sum_{i=1}^p \lambda_{(j)}^2 = \text{tr}(\mathbf{L} \mathbf{L}^T)$$

$$\mathbf{L} = [\sqrt{\lambda_{(1)}} \mathbf{e}_{(1)} \mid \sqrt{\lambda_{(2)}} \mathbf{e}_{(2)}]$$

$$\mathbf{L}^T = \begin{bmatrix} \sqrt{\lambda_{(1)}} \mathbf{e}_{(1)}^T \\ \sqrt{\lambda_{(2)}} \mathbf{e}_{(2)}^T \end{bmatrix}$$

$$\text{on multiplying } \mathbf{L}^T \mathbf{L} = \lambda_{(1)} \mathbf{e}_{(1)}^T \mathbf{e}_{(1)} + \lambda_{(2)} \mathbf{e}_{(2)}^T \mathbf{e}_{(2)} \\ = \underline{\underline{\lambda_{(1)} + \lambda_{(2)}}}$$

~~sum of 2 largest eigenvalues of S~~

on multiplying

$$LL^T = \begin{bmatrix} \cancel{d_1 e_{11} e_{11}^T} & \cancel{\sqrt{d_1 d_2} e_{11} e_{12}^T} \\ \cancel{\sqrt{d_1 d_2} e_{12} e_{11}^T} & \cancel{d_2 e_{22} e_{22}^T} \end{bmatrix} \begin{bmatrix} d_1 e_{11} e_{11}^T + d_2 e_{12} e_{12}^T \\ 0 \end{bmatrix}$$

$$\text{tr}(LL^T) = \text{tr}(L^T L) \quad \left(\begin{array}{l} \text{due to cyclic properties} \\ \text{of trace} \end{array} \right)$$

$$\Rightarrow \text{tr}(LL^T) = \text{tr}(L^T L)$$

$$\text{now } L^T L = \begin{bmatrix} \sqrt{d_1} e_{11}^T e_{11} + d_2 e_{12}^T e_{12} \\ 0 \end{bmatrix}$$

~~$\frac{1}{2} (d_1 + d_2)$~~
on multiplication we get

$$L^T L = \begin{bmatrix} d_{11} e_{11}^T e_{11} & \sqrt{d_{11} d_{22}} e_{11}^T e_{12} \\ \sqrt{d_{21} d_{12}} e_{12}^T e_{11} & d_{22} e_{12}^T e_{12} \end{bmatrix}$$

$$\begin{aligned} \text{tr}(L^T L) &= d_{11} e_{11}^T e_{11} + d_{22} e_{12}^T e_{12} \\ &= \underline{d_1 + d_2} \quad \left(\begin{array}{l} \text{because } e_1 \text{ and } e_{12} \\ \text{are normalized} \end{array} \right) \end{aligned}$$

04

hence proved

3. (5 points) Let $X_1, X_2, \dots, X_n$ be a random sample of size n from a distribution with mean $\mu \in \mathbb{R}$ and variance μ^2 . Let $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$. Derive a variance-stabilizing transformation $g(\bar{X}_n)$ such that the asymptotic variance of $g(\bar{X}_n)$ does not depend on μ .

by Delta method

$$\sqrt{n}(g(\bar{X}_n) - g(\mu)) \xrightarrow{d} N(0, (g'(\mu))^2 \sigma^2)$$

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

$$\text{mean} = \mu, \quad \text{variance} = \frac{\mu^2}{n}$$

for the variance of $g(\bar{X}_n)$ to be independent of μ

$$(g'(\mu))^2 \frac{\mu^2}{n} = k \xrightarrow{\text{independent of } \mu} g'(\mu) = \frac{c}{\mu} \quad (\text{for some } c)$$


$$\text{OS} \quad g(\mu) = c \ln(\mu) \quad (\text{for some } c)$$

so for $g(\bar{X}_n) = c \ln(\bar{X}_n)$

asymptotic variance of $g(\bar{X}_n)$ doesn't depend on μ

$$(g'(\mu))^2 \sigma^2 = \left(\frac{c}{\mu}\right)^2 \times \frac{\mu^2}{n} = \frac{c^2}{n} \rightarrow \text{independent of } \mu$$

4. (2 points) The figure below displays a Normal Q-Q plot for a dataset of size $n = 50$. Interpret this plot to determine if the data follows a normal distribution. If the normality assumption is violated, specify which class of distributions would more accurately characterize the data.



No, the dataset doesn't follow a normal distribution.
this can be inferred as near the tails the graph is
not linear.

~~The class of distributions which would more accurately
characterize this is uniform distribution.~~

The class of distributions which would more accurately
characterize this is heavy tailed distributions.

02

ADDITIONAL PAGES FOR ANSWERS
